Baseline Assessment – Stream Attributes

Reach S-E78/E82/R1 (Pipeline ROW) Perennial Spread C Webster County, West Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope >4%)
RBP Physical Characteristics Form	√ ·
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A – No habitat; Narrow stream.
Wolman Pebble Count	✓
Reference Reach Software Pebble Count Data	✓
Longitudinal Profile and Cross Sections	✓



Photo Type: DS View of Channel from Edge of TMB Location, Orientation, Photographer Initials: Downstream View of Channel from Edge of Timber Mat Bridge, ABK/AAK/TA



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, ABK/AAK/TA



Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, ABK/AAK/TA



Photo Type: CP, US View Location, Orientation, Photographer Initials: Center ROW, Upstream View, ABK/AAK/TA



Photo Type: CP, DS View Location, Orientation, Photographer Initials: Center ROW, Downstream View, ABK/AAK/TA



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, ABK/AAK/TA



Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, ABK/AAK/TA

 $[&]quot;Q: \label{lem:conditions} \begin{subarray}{l} $$ "Q: \charleston \charge \c$

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

USACE FILE NO./ Project Name:		MOL	JNTAIN VALLEY	PIPELINE		COORDINATES:	Lat.	38.676223	Lon.		-80.477663	WEATHER:	10	00% cloudy	DATE:			
(v2.1, Sept 2015)					(in De	cimal Degrees)											8/10/21	
IMPACT STREAM/SITE ID	AND SITE DES	CRIPTION		LINT to Left Forl	Holly River (S-E7	R_F82_P1\		MITIGATION STREA	M CLASS/SITE ID	AND SITE	DESCRIPTION:				Comments:			
(watershed size {acreage},				OIT to Leit For	Criony River (O-E/	J-L02-1(1)			ed size {acreage}, unaltered						Comments.			
STREAM IMPACT LENGTH:	102	FORM	OF		MIT C	OORDINATES:	Lat.		Lon.			PRECIPITATION PAST 48 HRS	:		Mitigation Length:			
		MITIGAT	ION:	RESTORATION (Levels I-III)	(in De	cimal Degrees)									13.11.13.			
								Column No. 3- N	Mitigation Projected a	at Five Year	9	Column No. 4- Mitigation	Column No. 4- Mitigation Projected at Ten Years Post Completion (Credit)					
Column No. 1- Impact Existing	Condition (Deb	oit)	•	Column No. 2- Mitigation Exist	ing Condition - Bas	eline (Credit)		Pos	t Completion (Credit))								ojected at Matu
Stream Classification:	Peren	nnial	Stream	Classification:				Stream Classification:		0		Stream Classification:	d)	Stream Classification:		0	
Percent Stream Channel Slo	ppe	1.9		Percent Stream Chann	el Slope			Percent Stream	Channel Slope		0	Percent Stream Channe	el Slope	0	Percent Stream Chann	el Slope		0
HGM Score (attach da	ata forms):			HGM Score (at	tach data forms):			HGM Score (attach data forms):				HGM Score (attac	ch data forms):		HGM Score (atta	h data forms	s):	
		Averege				Average					Average			Average				Average
Hydrology		Average	Hydrolo	oav		Average		Hydrology			Average	Hydrology		Average	Hydrology		A	verage
Biogeochemical Cycling		0		chemical Cycling		0		Biogeochemical Cycling			0	Biogeochemical Cycling		0	Biogeochemical Cycling			0
Habitat PART I - Physical, Chemical and	Riological Indica	atore	Habitat	PART I - Physical, Chemic	and Riological In	dicators		Habitat	Chemical and Biolog	rical Indicat	ors	Habitat PART I - Physical, Chemical	and Riological India	ators	Habitat PART I - Physical, Chemical	and Riologic	al Indicators	
PART 1 - Physical, Chemical and	Biological illulca	ators		FART 1 - Filysical, Chemic	ai and Biological in	uicators		FART I - FIIysical,	Chemical and Biolog	jicai iliulcai	ors	PART 1 - Physical, Glienlical	and Biological mulc	ators	PART 1 - Physical, Gliefflical	and Biologica	ai illuicators	
	Points Scale Range	Site Score			Points Scale Range	Site Score			Points Scale	e Range	Site Score		Points Scale Range	Site Score		Points Scale	Range	Site Score
PHYSICAL INDICATOR (Applies to all streams	classifications)		PHYSIC	CAL INDICATOR (Applies to all st	reams classifications)			PHYSICAL INDICATOR (Applies	s to all streams classifica	tions)		PHYSICAL INDICATOR (Applies to all str	reams classifications)		PHYSICAL INDICATOR (Applies to all st	eams classificati	ions)	
USEPA RBP (High Gradient Data Sheet)				RBP (Low Gradient Data She				USEPA RBP (High Gradient D				USEPA RBP (High Gradient Data She			USEPA RBP (High Gradient Data She			
Epifaunal Substrate/Available Cover Epipaddadaaaa	0-20	14 14		unal Substrate/Available Cover	0-20			Epifaunal Substrate/Available Endeddadada				Epifaunal Substrate/Available Cover Epihaddadassa	0-20		Epifaunal Substrate/Available Cover Epifaunal Substrate/Available Cover	0-20		
Embeddedness Velocity/ Depth Regime	0-20 0-20	7		Substrate Characterization Variability	0-20 0-20			Embeddedness Velocity/ Depth Regime	0-20 0-20			Embeddedness Velocity/ Depth Regime	0-20 0-20		Embeddedness Velocity/ Depth Regime	0-20 0-20	-	
Velectify Depart Regime Sediment Deposition	0-20	12		ment Deposition	0-20			4. Sediment Deposition	0-20			4. Sediment Deposition	0-20		Sediment Deposition	0-20	_	
5. Channel Flow Status	0-20	14		nel Flow Status	0-20			5. Channel Flow Status	0-20			5. Channel Flow Status	0-20		5. Channel Flow Status	0-20	1	
6. Channel Alteration	0-20 0-1	17	6. Chan	nel Alteration	0-20			6. Channel Alteration	0-20	0-1		6. Channel Alteration	0-20		Channel Alteration	0-20	0-1	
7. Frequency of Riffles (or bends)	0-20	7	7. Chan	nel Sinuosity	0-20			7. Frequency of Riffles (or bend	s) 0-20			7. Frequency of Riffles (or bends)	0-20		7. Frequency of Riffles (or bends)	0-20		
8. Bank Stability (LB & RB)	0-20	14	8. Bank	Stability (LB & RB)	0-20			8. Bank Stability (LB & RB)	0-20			8. Bank Stability (LB & RB)	0-20		8. Bank Stability (LB & RB)	0-20		
Vegetative Protection (LB & RB)	0-20	20		tative Protection (LB & RB)	0-20			Vegetative Protection (LB & F				Vegetative Protection (LB & RB)	0-20		Vegetative Protection (LB & RB)	0-20		
10. Riparian Vegetative Zone Width (LB & RB)	0-20	8		rian Vegetative Zone Width (LB & F				10. Riparian Vegetative Zone Widtl				10. Riparian Vegetative Zone Width (LB & F			10. Riparian Vegetative Zone Width (LB & F			
Total RBP Score Sub-Total	Suboptimal	127 0.635		BP Score	Poor	0		Total RBP Score	Pi	oor	0	Total RBP Score Sub-Total	Poor	0	Total RBP Score Sub-Total	Po	oor	0
Sub-1 otal CHEMICAL INDICATOR (Applies to Intermitten	nt and Perennial Stre		Sub-Tot CHEMIC	CAL INDICATOR (Applies to Inter	mittent and Perennial S	treams)		Sub-Total CHEMICAL INDICATOR (Applie	es to Intermittent and Pere	ennial Stream	s)	CHEMICAL INDICATOR (Applies to Inter	mittent and Perennial St	reams)	CHEMICAL INDICATOR (Applies to Inter	mittent and Pere	nnial Streams))
WVDEP Water Quality Indicators (General)	1		WVDER	Water Quality Indicators (Ge	neral)			WVDEP Water Quality Indicate	ors (General)			WVDEP Water Quality Indicators (Ger	neral)		WVDEP Water Quality Indicators (Ger	neral)		
Specific Conductivity				c Conductivity	neral)	0		Specific Conductivity	ors (General)			Specific Conductivity	ilei ai)		Specific Conductivity	eraij		
	0-90	70.0			0-90				0-90				0-90			0-90		
<=99 - 90 points	0-90	79.8			0-90				0-90				0-90			0-90		
pH		00	pН			0		pH				pH			рН			
6.0-8.0 = 80 points	0-80	6.48			5-90				5-90	0-1			5-90			5-90	0-1	
0.0-6.0 = 80 points			DO		L			DO.				DO.			DO.	_		
	10-30	0.00	20		10-30				10-30			50	10-30			10-30		
>5.0 = 30 points	10-30	6.63			10-30				10-30				10-30			10-30		
Sub-Total		1	Sub-Tot			0		Sub-Total			0	Sub-Total		0	Sub-Total			0
BIOLOGICAL INDICATOR (Applies to Intermit	tent and Perennial S	Streams)	BIOLOG	GICAL INDICATOR (Applies to Ir	termittent and Perennia	l Streams)		BIOLOGICAL INDICATOR (Ap)	plies to Intermittent and	d Perennial S	treams)	BIOLOGICAL INDICATOR (Applies to I	ntermittent and Perenn	nial Streams)	BIOLOGICAL INDICATOR (Applies to I	ntermittent and	Perennial Str	eams)
WV Stream Condition Index (WVSCI)			WV Stre	eam Condition Index (WVSCI)				WV Stream Condition Index (\				WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)	$\overline{}$		
0	0-100 0-1				0-100 0-1				0-100	0-1			0-100 0-1			0-100	0-1	
Sub-Total		0	Sub-Tot	tal		0		Sub-Total			0	Sub-Total		0	Sub-Total			0
PART II - Index and U	nit Score			DADT II Indo	and Unit Score			DART	II - Index and Unit Co.	ore	II	PART II - Index a	nd Unit Score	П	PART II - Index a	nd Unit Secre		
PART II - IIIdex and Onit Score				PART II - Index and Unit Score			PART II - Index and Unit Score				FANT II - IIIUUX A	na onit ocore		PART II - IIIQEX 8	ia onit ocole			
No. of	lines For	Unit C		1	11:	Limit Comm		1.4.			Hait Can	1.4.	1:	Unit O	1. 3.		-F4 Tr	nit Car
Index	Linear Feet	Unit Score		Index	Linear Feet	Unit Score		Index	Linea	ar Feet	Unit Score	Index	Linear Feet	Unit Score	Index	Linear	r Feet Un	nit Score
0.818	102	83.385		0	0	0		0		0	0	0	0	0	0	0)	0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME UNT to	Left Fork Holly River	LOCATION S-E78/E82/R1 Spre	ead C
STATION#F	RIVERMILE	STREAM CLASS Perennial	
LAT 38.676223 L	ONG -80.477663	COUNTY Webster	
STORET#		AGENCY Potesta	
INVESTIGATORS AK/AK	√TA		
FORM COMPLETED BY	A. Kincaid	DATE 8/10/2021 TIME 1030 AM	REASON FOR SURVEY Preliminary Assessment
WEATHER CONDITIONS	Now	Past 24 hours	Has there been a heavy rain in the last 7 days? Yes No
A SACRA CONTRACTOR OF A SACRA	rain (shower	Steady runny	Air Temperature 75 F 0 C Other Previous heavy rain lasting 20 mins
SITE LOCATION/MAP	Draw a map of the sit	e and indicate the areas sample	ed (or attach a photograph)
	Bielge	Heavy be regulation of the survey of the sur	
STREAM CHARACTERIZATION	Stream Subsystem Perennial Interpretation Stream Origin	ermittent Tidal	Stream Type ✓Coldwater Warmwater Catchment Area km²
	Glacial Non-glacial montane Swamp and bog	Spring-fed	

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Fores	Pasture Industria	duse rcial al	Local Watershed NPS □ No evidence □ Son □ Obvious sources Local Watershed Erosi □ None □ Moderate	ne potential sources
RIPARIA VEGETA (18 meter	N TION buffer)		e the dominant type and s S ant species present Gra	record the do arubs asses domina	minant species present He Grasses He nt	rbaceous
INSTREA FEATURI		Estima Sampli Area in Estima Surface (at thal	ted Stream Width 2.5 ng Reach Area km² (m²x1000) ted Stream Depth e Velocity 0.0 m	m² km²	2202000 12200 (N) 12200 (N)	ly shaded □Shadedm epresented by Stream Run_10% ☑No ☑No
LARGE V DEBRIS	VOODY	LWD Density	m ² of LWDm	1 ² /km ² (LWD/	reach area) N/A	
AQUATIO VEGETA		Domin	e the dominant type and ed emergent Ro ing Algae At	record the do ooted submerge tached Algae	minant species present nt Rooted floating	☐Free floating
WATER (QUALITY	Specific Dissolv pH 6.4	rature 19.5 C c Conductance 79.8 us/cm ed Oxygen 6.63 mg/L 8 SU ity 12.9 ntu strument Used YSI			Chemical Other Globs Flecks
SEDIMEN SUBSTRA		Odors Norm Chen Othe		Petroleum None	are the undersides blac	h are not deeply embedded,
INC		STRATE dd up to	COMPONENTS		ORGANIC SUBSTRATE C	
Substrate Type	Diamet		% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area
Bedrock Boulder	> 256 mm (10")		0	Detritus	sticks, wood, coarse plant materials (CPOM)	10
Cobble Gravel	64-256 mm (2.5 2-64 mm (0.1"-2	"-10")	10	Muck-Mud	black, very fine organic (FPOM)	-
Sand	0.06-2mm (gritt	y)	35	Marl	grey, shell fragments	
Silt	0.004-0.06 mm		5			_
Clay	< 0.004 mm (sli	ck)		1		

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME UNT to Left Fork Holly River	LOCATION S-E78/E82/R1 Spread C								
STATION # RIVERMILE	STREAM CLASS Perennial								
LAT 38.676223 LONG -80.477663	COUNTY Webster								
STORET#	AGENCY Potesta								
INVESTIGATORS AK/AK/TA									
FORM COMPLETED BY A. Kincaid	DATE 8/10/2021 REASON FOR SURVEY Preliminary Assessment								

	Habitat		Condition	ı Category	
	Parameter	Optimal	Suboptimal	Marginal	Poor
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.
1	SCORE 14	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.
ed ir	SCORE 14	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime N/A	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).
ıram	SCORE 7	20 19 18 17 16	15 14 13 12 11	10 9 8 🚺 6	5 4 3 2 1 0
P ₂	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.
	_{SCORE} 12	20 19 18 17 16	15 14 13 🔃 11	10 9 8 7 6	5 4 3 2 1 0
	5. Channel Flow Status N/A	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.
	SCORE 14	20 19 18 17 16	15 🚺 13 12 11	10 9 8 7 6	5 4 3 2 1 0

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	ı Category	
	Parameter	Optimal	Suboptimal	Marginal	Poor
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.
	SCORE 17	20 19 18 17 16	present. 15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
ing reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.
ampl	score 7	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing deuterstand.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional sears.
e eva	SCORE 6	Left Bank 10 9	8 7 6	5 4 3	2 1 0
to b	SCORE 8	Right Bank 10 9	8 7 6	5 4 3	2 1 0
Parameter	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.
	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0
	SCORE 10	Right Bank 10 9	8 7 6	5 4 3	2 1 0
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.
	SCORE 4	Left Bank 10 9	8 7 6	5 4 3	2 1 0
	SCORE 4	Right Bank 10 9	8 7 6	5 4 3	2 1 0

Total Score 127

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STATION #	1.4.1.1.10.1.	eft F	ork He	olly Ri	ver		LO	CATIO	N S-E78	E82/	R1 Sp	oread	С						
STATION#_	R	IVE	RM	ILE_			ST	REAM	CLASS I	Pere	nnia								
LAT 38.676223	_ L	ONO	j -80	47766	3		CO	UNTY	W	ebst	er								
STORET#							AG	ENCY	Potesta	9									
INVESTIGATORS	AK/AK	JΤΑ										1	TO.	NUMBER					
FORM COMPLETE	D BY	Α	. ł	(ir	nca	aic	DA TIM	TE 8/ 1E 10	10/2021 030 AM			1	REAS	SON FOR SURVEY	relimir	nary	Asse	essn	nent
HABITAT TYPES	In [dica Co Sub	ite th obbl merg	ne pe e_ ged N	rcen % Macro	tage 6 D	of each Snags	habita %	t type pr	esen eget	t ated ther	Banl	ks	%	%				
SAMPLE	G	ear	used		D-fi	ame	kic	k-net											
COLLECTION	н	ow v	vere	the	samp	oles c	ollected	1?	wadin	g		fror	n ban	k from boa	ıt				
	∥□	Cob	ble				Snags		n in each □V	eget	ated	Ban	ks	Sand					
GENERAL COMMENTS	В	er	ıth	ics	uı	nak	ole t	o be	e colle	ec	tec	1. k	Νo	habitat. Nar	rov	V S	tre	ear	n.
QUALITATIVE Indicate estimate Dominant									erved, 1	= J	Rare	e, 2	= C	ommon, 3= Abuno	dant,	4 =	=		
Periphyton					0	1	2 3	4		Sli	nes				0	1	2	3	4
Periphyton Filamentous Algae	;				-	-	2 3 2 3					nve	rtebr	ates	-	-	2 2	-	-
Periphyton Filamentous Algae Macrophytes	•				0	1		4			croi	nve	rtebr	ates	0	-	2	-	4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate	ATIO	und	anc	e:	0 0 (AC) 0 = org	1 1 ROB Abso	2 3 2 3 EENTHent/Noms), 3=	4 4 HOS of Obse	erved, ndant (2	Ma Fis 1 = >10	croi n Rar org	e (1 anis	-3 o	rganisms), 2 = Coi , 4 = Dominant (>:	0 0	1 1 n (3	2 2 2	3 3	4
Filamentous Algae Macrophytes	ATIO	und	anc	e:	0 0 (AC) 0 = org	1 1 ROB Abso	2 3 2 3 EENTHent/Noms), 3=	4 4 4 HOS of Obse	erved, ndant (2	Ma Fis 1 = >10	croi n Rar org	e (1 anis	-3 o	rganisms), 2 = Coi , 4 = Dominant (>: Chironomidae	0 0 mmo 50 o	1 1 n (3	2 2 -9 nism	3 3	4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa	ATIO d abo	1 1	2 2	e: 3 3	0 0 (AC) 0 = org:	1 1 ROB Abseanisi	2 3 2 3 EENTHent/Noms), 3= isopte	4 4 4 HOS of Obse = Abuse	erved, and ant (2	Ma Fis 1 = >10	Rar org	e (1 anis	-3 or sms)	rganisms), 2 = Cor , 4 = Dominant (> Chironomidae Ephemeroptera	0 0 0 mmo 50 o	1 1 n (3 rgan	2 2 -9 nism 2 2	3 3 3 3 3	4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes	O 0 0	1 1 1	2 2 2	3 3 3	0 0 0 0 = org:	1 1 ROB Abseanisi An Zy He	2 3 2 3 EENTHent/Nons), 3- isopte- gopter- mipter	4 4 4 HOS at Obse = Abut	erved, and ant (2)	Ma Fis 1 = >10	Rar org	e (1 anis 3 3	-3 or sms)	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria	0 0 0 0	1 1 1 1	2 2 2 2	3 3 3 3	0 0 0 0 = org	An Zy He Co	2 3 2 3 EENTHent/Noms), 3= isopter gopter mipter leopte	4 4 4 HOS of Obse = Abuse ra a a a ra	0 0 0 0	Ma Fis 1 = >10	Rar org	3 3 3 3	-3 or sms)	rganisms), 2 = Cor , 4 = Dominant (> Chironomidae Ephemeroptera	0 0 0 mmo 50 o	1 1 n (3 rgan	2 2 -9 nism 2 2	3 3 3 3 3	4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea	0 0 0 0 0	1 1 1 1	2 2 2 2 2	3 3 3 3	0 0 0 0 = org	And	2 3 2 3 EENTHent/Nons), 3= isopte gopter mipter leopte pidopt	4 4 4 HOS of Obse = Abuse ra a a a ra	0 0 0 0 0	Ma Fis 1 = >10	Rar org	3 3 3 3 3	4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta	0 0 0 0 0 0	1 1 1 1 1	2 2 2 2 2 2	3 3 3 3 3	0 0 0 0 = orgs	Annisi Annisi An Zy He Co Le Sia	2 3 2 3 EENTHent/Nons), 3= isoptement is not included in the content of the content is not included in the content in the content is not included in the content i	4 4 4 HOS of Obse = Abur ra a ra era	0 0 0 0 0	Ma Fis 1 = >10	Rarrorg 2 2 2 2 2 2	e (1 3 3 3 3 3 3	-3 onsms) 4 4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda	0 0 0 0 0 0	1 1 1 1 1 1 1	2 2 2 2 2 2 2 2	3 3 3 3 3 3	0 0 0 = 0 = 0 org: 4 4 4 4 4 4 4 4	And	2 3 2 3 EENTHent/Noms), 3= isopter gopter mipter leopte pidopt lidae rydalid	4 4 4 HOS at Obse = Abut ra at a ra era	0 0 0 0 0 0	Ma Fis 1 = >10	Rarrorg 2 2 2 2 2 2 2 2	3 3 3 3 3 3	-3 or sms) 4 4 4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda	0 0 0 0 0 0	1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3	0 0 0 0 = org	Anna Anna Anna Anna Anna Anna Anna Anna	2 3 2 3 EENTHent/Nons), 3= isopter gopter mipter leopte pidopt didae rydalio bulidae	4 4 4 HOS at Obse = Abut ra at	0 0 0 0 0 0 0	Ma Fis	Rar org 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3	4 4 4 4 4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda Decapoda	0 0 0 0 0 0 0 0	1 1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3	0 0 0 = 0 0 = 0 4 4 4 4 4 4 4 4 4 4 4	Annisi Annisi Annisi Anti- Zy He Co Le Sia Co Tip En	2 3 2 3 EENTHent/Noms), 3= isopter gopter gidopte gido	4 4 4 HOS at Obse = Abut ra at ra era era	0 0 0 0 0 0 0	Ma Fis	Rar org 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3	-3 on 4 4 4 4 4 4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4 4
Filamentous Algae Macrophytes FIELD OBSERV Indicate estimate Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda	0 0 0 0 0 0	1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3	0 0 0 0 = org	Annisi Annisi Annisi Co	2 3 2 3 EENTHent/Nons), 3= isopter gopter mipter leopte pidopt didae rydalio bulidae	4 4 4 HOS At Obse Abut ra a a a a a a a a a a a a a a a a a	0 0 0 0 0 0 0	Ma Fis	Rar org 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3	4 4 4 4 4 4 4 4 4	rganisms), 2 = Con , 4 = Dominant (>) Chironomidae Ephemeroptera Trichoptera	0 0 0 mmo 50 o	1 1 1 m (3 rgan	2 2 2 9 nism 2 2 2	3 3 3 3 3 3	4 4 4 4 4

SITE ID: S-E78/E82/F	31	
DATE: 8/10/21		

Wolman Pebble Count (Reach Wide) (mm) *


Wolmen Peb	ble Count (R	each Wide)	mm							NOTES:
0.175	33	0.50	18	0.63	21	18	0.06	0.60	0.062	
0.175	12	74.0	- 38	1.062	14	4:052	0.062	060	0.662	
0.60	19	31D	33	6:062	18	6.052	zl	0.30	8	
0.75	33	19	jur	4,062	14	4062	22	10	16	
0.175	24	27	89	6-125	٩	2062	26	12	4	
44	152	U	367	0.062	12	6,062	38	15	70	
126	100	42	0,70	0.125	0.25	4362	19	4	6	
156	41	UL	ilia	g	23	4.062	002	0,36	3	
0.125	32	21	105	0.062	.062	4,062	0.062	5	3	
0.175	4	22	34	0.062	4.062	4,062	183	-18	7	

Riffle Pebble	Count		JV -0 2/7	Maria 257	A. T. A.	Programme I			1544	NOTES:
20	30	20	158	158	158	158	158	18	158	
13	18	28	158	158	158	158	158	22	26	collected at rittle X-sec (surveyed)
17	26	62	158	158	158	158	158	28	26	X-Sec (surveyed)
32	16	62	158	128	58	158	20	28	2	
25	19	62	158	158	158	158	27	18	21	
75	26	21	158	158	158	158	18	18	31	
ιC	13	34	158	158	158	158	19	18	31	
14	15-	34	158	158	158	158	26	20	27	
18	12	2	17	33	62	3.2	17	20	24	
ና	15	- 11	17.	33	62	32	27	20	19	


	19	20	27	32	6z	33	17-	11	15	ৰ
NOTES:		if Now Jo			The Part			O PARTY		
*										


Inches	PARTICLE	Millimeters	
	Silt / Clay	< .062	S/C
	Very Fine	.062125	_
	Fine	.125 - 25	S
	Medium	.25 - ,50	S A N
	Çoarse	.50 - 1.0	D
0408	Very Coarse	1.0 - 2	
.0816	Very Fine	2-4	1 4 E
.1622	Fine	4 - 5.7	
.2231	Fine	5.7 - 8	G
.3144	Medium	8-11.3	R
.4463	Medium	11.3 - 16	
.6389	Coarse	16 - 22.6	E
.89 - 1,3	Coarse	22.6 - 32	
1.3 - 1.8	Very Coarse	32 - 45	000
1.8 - 2.5	Very Coarse	45-64	*CX
2.5 - 3.5	Small	64 - 90	HOL
3.5 - 5.0	Small	90 - 128	
5.0 - 7.1	Large	128 - 180	
7.1 - 10.1	Large	180 - 256	5
10.1 - 14,3	Small	256 - 362	8
14.3 - 20	Small	362 - 512	Ŭ
20 - 40	Medium	512 - 1024	SP.
40 - 80	Large-Vry Large	1024 - 2048	R
	Bedrock		BDRK


Bankfull Channel			
Bankfull Channel			
Material Size Rang			
	0.062 23		
very fine sand <u>0.062</u> - 0).125 7		
fine sand 0.125 - 0			
medium sand 0.25 - 0			
coarse sand 0.5 - 1	7		
very coarse sand 1 - 2	}		
very fine gravel 2 - 4			
fine gravel 4 - 6	1		
fine gravel 6 - 8	4		
medium gravel <u>8 - 1</u>			
medium gravel <u>11 - 1</u>			
coarse gravel 16 - 2			
coarse gravel 22 - 3			
very coarse gravel 32 - 4			
very coarse gravel 45 - 6			
small cobble 64 - 9			
medium cobble 90 - 1			
large cobble 128 - 1			
very large cobble 180 - 2			
small boulder <u>256</u> - 3	-		
small boulder <u>362 - 5</u>			
medium boulder 512 - 1			
large boulder 1024 - 2			
very large boulder 2048 - 4	096		
total particle	count: 100		
bedrock			
clay hardpan			
detritus/wood			
artificial			
total	count: 100		
Note:			



S-E78-E82-R1







LEGEND


EXISTING SURVEY-LOCATED THALWEG

1176.87 +EXISTING SURVEYED GROUND SHOT ELEVATION

SURVEY NOTES:


- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS AND COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.

S-E78-E82-R1 BASELINE CROSS-SECTION A



DISTANCE ALONG CROSS-SECTION (FT)

S-E78-E82-R1 BASELINE CROSS-SECTION B



DISTANCE ALONG CROSS-SECTION (FT)

CROSS SECTION LEGEND — EXISTING GRADE

NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

PRE-CROSSING PHOTOS



PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS



PHOTO TAKEN LOOKING UPSTREAM FROM DOWNSTREAM IMPACT LIMITS

POST-CROSSING PHOTOS

PENDING CROSSING

PHOTO TAKEN LOOKING DOWNSTREAM UPSTREAM FROM IMPACT LIMITS

PENDING CROSSING

PHOTO TAKEN LOOKING UPSTREAM FROM UPSTREAM IMPACT LIMITS

PRE-CROSSING

Drawing No

Checked

BB/JLY Approved

NOTED Scale:

SEPT. 2021 Date: