Reach S-L38 (Pipeline ROW) Perennial Spread D Nicholas County, West Virginia

Data	Included
Photos	\checkmark
SWVM Form	\checkmark
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope >4%)
RBP Physical Characteristics Form	\checkmark
Water Quality Data	\checkmark
RBP Habitat Form	\checkmark
RBP Benthic Form	\checkmark
Benthic Identification Sheet	N/A – Low flow
Wolman Pebble Count	\checkmark
Reference Reach Software Pebble Count Data	\checkmark
Longitudinal Profile and Cross Sections	\checkmark



Photo Type: US at DS Edge of ROW

Location, Orientation, Photographer Initials: Downstream Edge of Right of Way, Upstream View, TF/AG/WP/EW



Photo Type: DS at DS Edge of ROW Location, Orientation, Photographer Initials: Downstream Edge of Right of Way, Downstream View, TF/AG/WP/EW



Photo Type: US at Center of ROW Location, Orientation, Photographer Initials: Center of Right of Way, Upstream View, TF/AG/WP/EW



Photo Type: DS at Center of ROW Location, Orientation, Photographer Initials: Center of Right of Way, Downstream View, TF/AG/WP/EW

Stream S-L38 (Pipeline ROW)



Photo Type: US at US Edge of ROW Location, Orientation, Photographer Initials: Upstream Edge of Right of Way, Upstream View, TF/AG/WP/EW



Photo Type: DS at US Edge of ROW Location, Orientation, Photographer Initials: Upstream Edge of Right of Way, Downstream View, TF/AG/WP/EW

"Q:\Charleston\2021 Projects\21-0244- MVP- STREAM AND WETLAND CONDITIONS ASSESSMENT AND SURVEY PLAN\002 - Pre-Crossing Monitoring\Spread D\S-L38"

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017


USACE FILE NO./ Project Name: (v2.1, Sept 2015)		Mountair	n Valley Pipeline	IMPACT COORDINATE: (in Decimal Degrees)	S: Lat.	38.205534	Lon.	-80.718246	WEATHER:		Cle
IMPACT STREAM/SITE II (watershed size {acreage			S-L38 UNT 1	o Riley Branch		MITIGATION STREAM CLAS (watershed size {acr					
STREAM IMPACT LENGTH:	75	FORM OF MITIGATION:	RESTORATION (Levels I-III)	MIT COORDINATES: (in Decimal Degrees)	Lat.		Lon.		PRECIPITATION PAST 48 HRS:		
Column No. 1- Impact Existin	ng Condition (Deb	it)	Column No. 2- Mitigation Existing 0	Condition - Baseline (Credit)		Column No. 3- Mitigation Post Comple	n Projected at Fi etion (Credit)	ve Years	Column No. 4- Mitigation Pro Post Completion		Γen Ye
Stream Classification:	Perer	nnial	Stream Classification:			Stream Classification:		0	Stream Classification:		
Percent Stream Channel S	Blope	4.9	Percent Stream Channel SI	оре		Percent Stream Channe	I Slope	0	Percent Stream Channel S	Slope	
HGM Score (attach	data forms):		HGM Score (attach	data forms):		HGM Score (atta	ach data forms	:	HGM Score (attach	data form	s):
		Average		Average				Average			
Hydrology		g	Hydrology			Hydrology			Hydrology	_	
Biogeochemical Cycling		0	Biogeochemical Cycling	0		Biogeochemical Cycling		0	Biogeochemical Cycling		<u> </u>
Habitat		U	Habitat	V		Habitat		- U	Habitat		<u> </u>
PART I - Physical, Chemical an	d Biological Indica	ators	PART I - Physical, Chemical ar	nd Biological Indicators		PART I - Physical, Chemica	al and Biological	Indicators	PART I - Physical, Chemical an	d Biologica	al Indi
	Points Scale Range	Site Score		Points Scale Range Site Score			Points Scale R	ange Site Score		Points Scale	Range
PHYSICAL INDICATOR (Applies to all stream	ns classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)		PHYSICAL INDICATOR (Applies to all stre	eams classifications)	PHYSICAL INDICATOR (Applies to all stream	ms classifica	tions)
USEPA RBP (High Gradient Data Sheet)			USEPA RBP (Low Gradient Data Sheet)			USEPA RBP (High Gradient Data Shee	et)		USEPA RBP (High Gradient Data Sheet)		
1. Epifaunal Substrate/Available Cover	0-20	10	 Epifaunal Substrate/Available Cover 	0-20		1. Epifaunal Substrate/Available Cover	0-20		 Epifaunal Substrate/Available Cover 	0-20	
2. Embeddedness	0-20	13	2. Pool Substrate Characterization	0-20		2. Embeddedness	0-20		2. Embeddedness	0-20	7
Velocity/ Depth Regime	0-20	9	Pool Variability	0-20		Velocity/ Depth Regime	0-20		Velocity/ Depth Regime	0-20	
4. Sediment Deposition	0-20	9	4. Sediment Deposition	0-20		Sediment Deposition	0-20		4. Sediment Deposition	0-20	
5. Channel Flow Status	0-20 0-1	8	5. Channel Flow Status	0-20 0-1		5. Channel Flow Status	0-20)-1	5. Channel Flow Status	0-20	0-1
6. Channel Alteration	0-20	14	6. Channel Alteration	0-20		6. Channel Alteration	0-20	J- 1	6. Channel Alteration	0-20	0-1
Frequency of Riffles (or bends)	0-20	10	7. Channel Sinuosity	0-20		Frequency of Riffles (or bends)	0-20		Frequency of Riffles (or bends)	0-20	
8. Bank Stability (LB & RB)	0-20	16	Bank Stability (LB & RB)	0-20		8. Bank Stability (LB & RB)	0-20		8. Bank Stability (LB & RB)	0-20	
Vegetative Protection (LB & RB)	0-20	18	Vegetative Protection (LB & RB)	0-20		Vegetative Protection (LB & RB)	0-20		Vegetative Protection (LB & RB)	0-20	
10. Riparian Vegetative Zone Width (LB & RB)		18	Riparian Vegetative Zone Width (LB & RB)	0-20		10. Riparian Vegetative Zone Width (LB & RE			10. Riparian Vegetative Zone Width (LB & RB)		
Total RBP Score	Suboptimal	125	Total RBP Score	Poor 0		Total RBP Score	Poor	0	Total RBP Score	Po	or
Sub-Total		0.625	Sub-Total	0		Sub-Total		0	Sub-Total		
CHEMICAL INDICATOR (Applies to Intermitt	tent and Perennial Str	eams)	CHEMICAL INDICATOR (Applies to Intermitter	nt and Perennial Streams)		CHEMICAL INDICATOR (Applies to Intern	nittent and Perennia	l Streams)	CHEMICAL INDICATOR (Applies to Intermit	tent and Pere	ennial S
WVDEP Water Quality Indicators (Generation	al)		WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (Gen	eral)		WVDEP Water Quality Indicators (Gener	al)	
Specific Conductivity			Specific Conductivity			Specific Conductivity			Specific Conductivity		
	0-90	38.6		0-90			0-90			0-90	
<=99 - 90 points	0.00	30.0		0.00			0.00			0.00	_
рН		48	рН			рН			рН		
	0-80	5.94		5-90 0-1			5-90	0-1		5-90	0-1
5.6-5.9 = 45 points											_
DO			DO			DO			DO		4
>5.0 = 30 points	10-30	6.4		10-30			10-30			10-30	
Sub-Total		0.825	Sub-Total	0		Sub-Total		0	Sub-Total		_
				, v							
BIOLOGICAL INDICATOR (Applies to Intern	nittent and Perennial	Streams)	BIOLOGICAL INDICATOR (Applies to Intermit	ttent and Perennial Streams)		BIOLOGICAL INDICATOR (Applies to In	termittent and Per	ennial Streams)	BIOLOGICAL INDICATOR (Applies to Inte	rmittent and	l Perer
WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)		
0	0-100 0-1			0-100 0-1			0-100	0-1		0-100	0-1
ě		0	Sub Total	0		Sub Total		0	Sub Total		
Sub-Total		U	Sub-Total	U		Sub-Total		U	Sub-Total		
DADT II. Index and	Unit Coore	1	DART II Index and	Unit Seere		DART II. Index	and Unit Coore		DART II Index and	Unit Coorr	

PART II - Index and Unit Score			
Index	Linear Feet	Unit Score	
0.725	75	54.375	

PART II - Index and Unit Score				
Index	Linear Feet	Unit Score		
0	0	0		

PART II - Index and Unit Score				
Index	Linear Feet	Unit Score		
0	0	0		

BIOLOGICAL INDICATOR (Applies to Intermittent and Peren					
WV Stream Condition Index (WVSCI)					
	0-100	0-1			
Sub-Total					
PART II - Index and U	nit Score				
Index	Linear	Feet			
0	0				






PART II - Index and Unit Score			
Index	Linear Feet	Unit Score	
0	0	0	

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME S-L38	LOCATION UNT to Riley Branch, Spread D		
STATION # RIVERMILE	STREAM CLASS Perennial		
LAT <u>38.205534</u> LONG <u>-80.718246</u>	COUNTY Nicholas		
STORET #	AGENCYPotesta/ Edge		
INVESTIGATORS TF, EW, AG, WP			
FORM COMPLETED BY TF, EW	DATE 8-25-21 TIME 1030 REASON FOR SURVEY Preliminary Assessment		



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERSHED FEATURES	Predominant Surrounding Landuse ✓ Forest Commercial Field/Pasture Industrial Agricultural Ø Other Residential	Local Watershed NPS Pollution ☐ No evidence ☑ Some potential sources ☐ Obvious sources ☐ Local Watershed Erosion ☑ None ☐ Moderate ☐ Heavy		
RIPARIAN VEGETATION (18 meter buffer)	Indicate the dominant type and record the dominant species present intervent. the blockberry	inant species present ☐ Grasses ☐ Herbaceous		
INSTREAM FEATURES	Estimated Reach Length \$0.11 m m Estimated Stream Width \$0.11 m m Sampling Reach Area \$300.62 m² Area in km² (m²x1000) km² Estimated Stream Depth \$3.11 m Surface Velocity \$2.29 m/sec (at thalweg) Stream Dry []	Canopy Cover Partly shaded □Shaded Partly open Partly shaded □Shaded High Water Mark \$0.1 Proportion of Reach Represented by Stream Morphology Types Riffle10 % Pool \$00 % Channelized Yes Dam Present Yes No		
LARGE WOODY DEBRIS	LWDm ² Density of LWDm ² /km ² (LWD/ res	ach area)		
AQUATIC VEGETATION	Indicate the dominant type and record the dominant Rooted emergent Rooted submergent Floating Algae Attached Algae Dominant species present Portion of the reach with aquatic vegetation	☐Rooted floating ☐Free floating		
WATER QUALITY US point	Temperature 10 C Specific Conductance 38.6 us/cm Dissolved Oxygen 6.4 mg/L pH 5.00 Turbidity 5.53 ntu WQ Instrument Used 381	Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Globs Slick Sheen Other Water Surface Oils Slick Sheen Other Globs Flecks Unone Other Clear Slightly turbid Opaque Stained		
SEDIMENT/ SUBSTRATE	Odors ✓ Normal Chemical Other Oils ✓ Absent Slight Moderate	Deposits □Paper fiber ☑Sand Sludge □Sawdust □Paper fiber ☑Sand Relict shells □Other		
INORGANIC SU	BSTRATE COMPONENTS C	DRGANIC SUBSTRATE COMPONENTS		

INORGANIC SUBSTRATE COMPONENTS (should add up to 100%)			ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)			
Substrate Type	Diameter	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area	
Bedrock		68	Detritus	sticks, wood, coarse plant	0	
Boulder	> 256 mm (10")	2		materials (CPOM)	0	
Cobble	64-256 mm (2.5"-10")	0	Muck-Mud	black, very fine organic (FPOM)	0	
Gravel	2-64 mm (0.1"-2.5")	0		(FPOM)	0	
Sand	0.06-2mm (gritty)	20	Marl	grey, shell fragments	0	
Silt	0.004-0.06 mm	10				
Clay	< 0.004 mm (slick)	0				

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-L38	LOCATION UNT to Riley Branch		
STATION # RIVERMILE	STREAM CLASS Perennial		
LAT <u>38.205534</u> LONG <u>-80.718246</u>	COUNTY Nicholas		
STORET #	AGENCYPotesta/ Edge		
INVESTIGATORS TF, EW, AG, WP			
FORM COMPLETED BY TF, EW	DATE 8-25-21 TIME 1030 AM PM REASON FOR SURVEY Preliminary Assessment		

	Habitat	Condition Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor	
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.	
	_{SCORE} 10▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0	
ı sampling reach	2. Embeddedness	Gravel, cobble, and Gravel, cobble, and Gravel, cobble, and			Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.	
ted ir	_{SCORE} 13 ▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0	
Parameters to be evaluated in sampling reach	3. Velocity/Depth All four velocity/depth Regime regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). N/A (Slow is < 0.3 m/s, deep is > 0.5 m.)		Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).	
Iram	score 9 💌	20 19 18 17 16	15 14 13 12 11	10 🧐 8 7 6	5 4 3 2 1 0	
P	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.	
	score 9	20 19 18 17 16	15 14 13 12 11	10 🧕 8 7 6	5 4 3 2 1 0	
	5. Channel Flow Status N/A	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.	
	SCORE 8	20 19 18 17 16	15 14 13 12 11	10 9 🚷 7 6	5 4 3 2 1 0	

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Conditi	on Category			
	Parameter	Optimal	Suboptimal	Marginal	Poor		
g reach	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.		
	score 14 ▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water o shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.		
d m be	_{SCORE} 10 ▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing detractment.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion	areas of erosion; high erosion potential during	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.		
2	SCORE 8	Left Bank 10 9	8 7 6	5 4 3	2 1 0		
	SCORE 8	Right Bank 10 9	8 7 6	5 4 3	2 1 0		
rarameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potentia to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- l half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.		
	SCORE 9	Left Bank 10 🥘	8 7 6	5 4 3	2 1 0		
	_{SCORE} 9 ▼ ,	Right Bank 10 🗕 🧕	8 7 6	5 4 3	2 1 0		
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone < meters: little or no riparian vegetation due to human activities.		
	score 9 ▼,	Left Bank 10 🧕	8 7 6	5 4 3	2 1 0		
	SCORE 9 ()	Right Bank 10 🧐	8 7 6	5 4 3	2 1 0		

Total Score 125

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-	_38	LOCATION UNT to Riley B	Branch					
STATION #	RIVERMILE	STREAM CLASS Perennial	STREAM CLASS Perennial					
LAT 38.205534	LONG -80.718246	COUNTY Nicholas						
STORET #		AGENCYPotesta/ Edge						
INVESTIGATORS T			LOT NUMBER					
FORM COMPLETED	^{BY} TF, EW	DATE 8-25-21 TIME 1030	REASON FOR SURVEY Preliminary Assessment					
HABITAT TYPES	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%							
SAMPLE COLLECTION	Gear used D-frame How were the samples coll Indicate the number of jat CobbleSn Submerged Macrophytes	lected? ☐ wading ☐ fi ps/kicks taken in each habitat ty pags ↓ Vegetated B	anksSand					
GENERAL COMMENTS No benthics collected due to sand/BR substrate and low water lev								

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

SITE ID:	5-635	(3)	Riley	Branch	Spread	ſ
17		1. A				

DATE: 8 25/21

COLLECTOR(S): TEEWAG

49	bble Count (F	175	445	-3945	627	FS	160	1.46	IS
6131	28	16	35	TS	15	173	15	15	0
5	50	45	13	FS	2	FS	B	FS.	18
25	91	5	20	8	27	FS	22	- Fis	70
15	ND.	125	15	博	15	45	170	F5)	18
(D)	Gar	FS	FS	195	'es	85	é Si	165	15
FS	ES.	195	51	FS	35	51	SL	ES	- 51
2.51	FS	91	135	FS	-91	150	FS	5T	160
SI	B	- ES	160	55	180	- 26 -	FS.	160	25
270	185	44	200	TE	i Asje	Eg Es	15	FS	15


iffle Peb	ble Count					
	_	 		 	 	
_			2	 	 	


	 	 NOTES:
· · · · · · · · · · · · · · · · · · ·	 	


Inche:	154-67	Millimeters	
	Sinclag	1 (62	\$.0
	Very Fine	282-115	6
	Fine	125 26	S
	Medium	25 - 50	SAND
	Coarse	50 - 1.0	Ð
92 - 58	V#17/20378#	16-2	1
31 - 81	Very Fixe	2 - 4	
16 - 22	2 ₁₀ e	소나는 것	
22 - 31 -	Fine	57.8	ିତ୍ର
31 - 44	Vedura	6.113	.R
447 62	Vedum	112-19	1A
63 - 89	Coarse	16 - 32 6	E
39-10	Coarse	22 8 - 30	9
13-18	Very Coarse	32 - 45	
16-26	Very Uparse	45,764	
25-25	Small	64+30	201
35-65	Sma"	90 - 128	Zel
55-71	Large	125 - 150	3E
7.1-10.5	Large	190 - 256	
10-1-14-5	Small	256 - 362	Ô
14/3 - 20	Smail	362 - 512	Ιų.
22+42	Vedium	512 - 1624	Juchul
40 90	Large-Vry Large	1124 - 2048	, e
	Bedroos		8033

NOTES:

NOTES:








	LEGEND
	STUDY AREA (EASEMENT)
· · ·	EXISTING SURVEY-LOCATED THALWEG
1176.87 +	EXISTING SURVEYED GROUND SHOT ELEVATION

SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON 8-25-2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- COMBINATION WITH SURVEY POINTS AND COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.








CROSS SECTION LEGEND									
EXISTING GRADE									
$\begin{array}{c c} \underline{CROSS \ SECTION} \\ Brite SCALE: & H: 1"=10' \\ CROSS & V: 1"=5' \end{array}$									

NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN



-S-L38 CAD File No.

MBS Drawn

СНН Checked

BB/JLY

Approved

NOTED Scale:

SEPT. 2021 Date:

21-0244-005 Project No.

Z

PHOTO TAKEN LOOKING UPSTREAM FROM UPSTREAM IMPACT LIMITS

Drawing No

PRE-CROSSING