Reach S-A71 (Timber Mat Crossing) Perennial Spread D Nicholas County, West Virginia

| Data | Included |
|--|--|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | N/A – Perennial stream (not shadeable) |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | N/A – No flow |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | N/A – No flow |
| Wolman Pebble Count | \checkmark |
| Reference Reach Software Pebble Count Data | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, KP/SM Lat: 38.321572 Long: -80.670958



Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, KP/SM Lat: 38.321572 Long: -80.670958

Stream S-A71 (Timber Mat Crossing)



Photo Type: US View at Center Location, Orientation, Photographer Initials: Center ROW, Upstream View, KP/SM Lat: 38.321572 Long: -80.670958



Photo Type: DS View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, KP/SM Lat: 38.321572 Long: -80.670958

Spread D

Stream S-A71 (Timber Mat Crossing)

Nicholas County



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, KP/SM Lat: 38.321572 Long: -80.670958



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, KP/SM Lat: 38.321572 Long: -80.670958

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | Mountain | n Valley Pipeline | IMPACT COORDINATES: (in Decimal Degrees) | Lat. | 38.321572 | Lon. | -80.670958 | WEATHER: | 50% | Cloud Cover | DATE: | September | r 4, 2021 |
|--|--------------------------|--|---|------|---|--------------------|-------------|---|--------------------|---|---|--------------------------------------|------------|
| IMPACT STREAM/SITE ID AND SI (watershed size (acreage), unaltered | | S- | A71 | | MITIGATION STREAM CLASS./SI (watershed size (acreage), t | | | | | | Comments: | No flow so readings or were ta | r benthics |
| TREAM IMPACT LENGTH: 2 | 2 FORM OF MITIGATION: | RESTORATION (Levels I-III) | MIT COORDINATES: (in Decimal Degrees) | Lat. | | Lon. | | PRECIPITATION PAST 48 HRS: | | | Mitigation Length: | | |
| Column No. 1- Impact Existing Conditi | on (Debit) | Column No. 2- Mitigation Existing C | ondition - Baseline (Credit) | | Column No. 3- Mitigation Proje Post Completion (| | ears | Column No. 4- Mitigation Proje Post Completion (C | | ars | Column No. 5- Mitigation Project | ed at Maturity (C | redit) |
| tream Classification: | Perennial | Stream Classification: | | | Stream Classification: | | 0 | Stream Classification: | c |) | Stream Classification: | 0 | |
| Percent Stream Channel Slope | 9 | Percent Stream Channel Slo | ope | | Percent Stream Channel Slop | pe | 0 | Percent Stream Channel Sic | pe | 0 | Percent Stream Channel S | lope | 0 |
| HGM Score (attach data form | s): | HGM Score (attach | data forms): | | HGM Score (attach da | ata forms): | | HGM Score (attach da | ta forms): | | HGM Score (attach o | ata forms): | |
| | Average | | Average | | | | Average | | | Average | | | Avera |
| drology ogeochemical Cycling | | Hydrology Biogeochemical Cycling | | | Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling | | |
| abitat | | Habitat | Ť | | Habitat | | | Habitat | | , i i i i i i i i i i i i i i i i i i i | Habitat | - | Ĩ |
| PART I - Physical, Chemical and Biologic | al Indicators | PART I - Physical, Chemical and | d Biological Indicators | | PART I - Physical, Chemical and | Biological Indi | cators | PART I - Physical, Chemical and E | Biological Indic | ators | PART I - Physical, Chemical and | Biological Indica | tors |
| Poleta Scale | Range Site Score | | Points Scale Range Site Score | | | Poinza Scale Range | Site Score | | Points Scale Range | Site Score | | Points Scale Range | Site S |
| YSICAL INDICATOR (Applies to all streams classificat | ions) | PHYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDICATOR (Applies to all streams cl | assifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDICATOR (Applies to all stream | a classifications) | |
| EPA RBP (High Gradient Data Sheet) | | USEPA RBP (Low Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | 1 | |
| pifaunal Substrate/Available Cover 0-20 mbeddedness 0-20 | 0 | 1. Epifaunal Substrate/Available Cover 2. Pool Substrate Characterization | 0-20 | | 1. Epifaunal Substrate/Available Cover 2. Embeddedness | 0-20 | | 1. Epifaunal Substrate/Available Cover | 0-20 | | 1. Epifaunal Substrate/Available Cover 2. Embeddedness | 0-20 | |
| Embeddedness 0-20 /elocity/ Depth Regime 0-20 | 0 | 2. Pool Substrate Characterization 3. Pool Variability | 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | - |
| Sediment Deposition 0-20 | 19 | 4. Sediment Deposition | 0.20 | | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition | 0-20 | |
| Channel Flow Status 0-20 | 0-1 | 5. Channel Flow Status | 0-20 0-1 | | 5. Channel Flow Status | 0-20 0-1 | | 5. Channel Flow Status | 0-20 0-1 | | 5. Channel Flow Status | 0-20 0-1 | |
| channel Alteration 0-20 | 18 | 6. Channel Alteration | 0-20 | | 6. Channel Alteration | 0-20 | | 6. Channel Alteration | 0-20 | | 6. Channel Alteration | 0-20 | |
| requency of Riffles (or bends) 0-20 lank Stability (LB & RB) 0-20 | 18 | 7. Channel Sinuosity 8. Bank Stability (LB & RB) | 0-20 | | 7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB) | 0-20 | | 7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB) | 0-20 | | 7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB) | 0-20 | |
| (egetative Protection (LB & RB) 0-20 | 16 | 9. Vegetative Protection (LB & RB) | 0.20 | | 9. Vegetative Protection (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) | 0-20 | |
| Riparian Vegetative Zone Width (LB & RB) 0-20 | 14 | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | |
| tal RBP Score Marc | | Total RBP Score | Poor 0 | | Total RBP Score | Poor | 0 | Total RBP Score | Poor | 0 | Total RBP Score | Poor | |
| b-Total HEMICAL INDICATOR (Applies to Intermittent and Pere | 0.425 | Sub-Total CHEMICAL INDICATOR (Applies to Intermittent | 0 | | Sub-Total CHEMICAL INDICATOR (Applies to Intermittent a | | 0 | Sub-Total CHEMICAL INDICATOR (Applies to Intermittent | and Barranial Or | 0 | Sub-Total CHEMICAL INDICATOR (Applies to Intermitte | | (|
| | inal Streams) | | | | | ind Perennial Sire | airis) | | | rearris) | | | ams) |
| DEP Water Quality Indicators (General) ecific Conductivity | | WVDEP Water Quality Indicators (General) Specific Conductivity | | | WVDEP Water Quality Indicators (General) Specific Conductivity | | | WVDEP Water Quality Indicators (General) Specific Conductivity | | | WVDEP Water Quality Indicators (General Specific Conductivity | <u> </u> | |
| 0-90 0-90 | | | 0-90 | | | 0-90 | | | 0-90 | | | 0-90 | |
| 0-80 | 0-1 | рН | 5-90 0-1 | | pH | 5-90 0-1 | | рН | 5-90 0-1 | | рН | 5-90 0-1 | |
| 5.6-5.9 = 45 points | | 80 | | | P0 | | | PO | | | 80 | | |
| 10-30 | | 50 | 10-30 | | | 10-30 | | 50 | 10-30 | | 50 | 10-30 | |
| | | 0.1.7.1.1 | 1030 | | 0.1.7.1. | 10-30 | | 0.1.7.1 | 10-30 | | 0.1.7.1. | 10-30 | 0 |
| -Total COGICAL INDICATOR (Applies to Intermittent and Pe | | Sub-Total BIOLOGICAL INDICATOR (Applies to Intermitte | U U | | Sub-Total BIOLOGICAL INDICATOR (Applies to Intermitt | | 0 | Sub-Total BIOLOGICAL INDICATOR (Applies to Intermi | | U (Internet) | Sub-Total BIOLOGICAL INDICATOR (Applies to Interr | | · · |
| | Terma Streams) | | nicand Perenniai Sireanis) | | | ent and Felenin | ai Streams) | | ttent and Feren | ilai Streams) | | | si otrean |
| / Stream Condition Index (WVSCI) 0-100 | 0.1 | WV Stream Condition Index (WVSCI) | 0-100 0-1 | | WV Stream Condition Index (WVSCI) | 0-100 0-1 | | WV Stream Condition Index (WVSCI) | 0-100 0-1 | | WV Stream Condition Index (WVSCI) | 0-100 0-1 | |
| 0 b-Total | 0 | Sub-Total | 0 | | Sub-Total | 0-100 | 0 | Sub-Total | 0-100 0-1 | 0 | Sub-Total | 0-100 0-1 | 0 |
| - 10m | | our rout | v | Ш | 000 1000 | | U U | our i dai | | <u> </u> | Par rola | | |
| PART II - Index and Unit Score | | PART II - Index and | Unit Score | 1 | PART II - Index and U | nit Score | | PART II - Index and Ur | iit Score | | PART II - Index and U | Init Score | |
| Index Linear | r Feet Unit Score | Index | Linear Feet Unit Score | | Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score | Index | Linear Feet | Unit S |
| | | | | | | | | | | | | 4 | |
| | | | | | | | | | | | | | |

| inear Feet | Unit Score | Index | L |
|------------|------------|-------|---|
| 22 | 13.475 | 0 | |
| | | | |


0.613

| nit Score | |
|-------------|------------|
| | |
| Linear Feet | Unit Score |
| 0 | 0 |
| • | • |

| Stream Condition Index (WVSCI) | | | | | | |
|--------------------------------|--------|------|------------|--|--|--|
| | 0-100 | 0-1 | | | | |
| Total | | | 0 | | | |
| | | | | | | |
| PART II - Index and Unit Score | | | | | | |
| Index | Linear | Feet | Unit Score | | | |
| 0 | 0 | | 0 | | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME | REASON FOR SURVEY |



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential Indicate the dominant type and record the domin Trees Shrubs Dominant species present | Grasses Herbaceous |
|--|---|--|
| INSTREAM FEATURES | Estimated Reach Length m Estimated Stream Width m Sampling Reach Area ² Area in km² (m²x1000) km² Estimated Stream Depth m Surface Velocity m/sec (at thalweg) m/sec | Canopy Cover Partly open Partly shaded Shaded High Water Mark m Proportion of Reach Represented by Stream Morphology Types Riffle% Run% Riffle % Root % Root % No No |
| LARGE WOODY DEBRIS AQUATIC VEGETATION | LWDm² Density of LWDm²/km² (LWD/ reac Indicate the dominant type and record the domin Rooted emergent Rooted submergent Floating Algae Attached Algae Dominant species present | ant species present Rooted floating Free floating |
| WATER QUALITY | Temperature0 C Specific Conductance Dissolved Oxygen pH Turbidity WQ Instrument Used | Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Slick Sheen Globs Flecks None Other Turbidity (if not measured) Clear Slightly turbid Clear Slightly turbid Turbid Opaque Stained Other |
| SEDIMENT/ SUBSTRATE | Odors Petroleum Normal Sewage Petroleum Chemical Anaerobic None Other | Deposits Sludge Sawdust Paper fiber Sand Relict shells Other Lpoking at stones which are not deeply embedded, are the undersides black in color? Yes No |

| INORGANIC SUBSTRATE COMPONENTS (should add up to 100%) | | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | |
|---|----------------------|------------------------------------|---|----------------------------|-----------------------------------|--|
| Substrate Type | Diameter | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | | |
| Cobble | 64-256 mm (2.5"-10") | | Muck-Mud | black, very fine organic | | |
| Gravel | 2-64 mm (0.1"-2.5") | | | (FPOM) | | |
| Sand | 0.06-2mm (gritty) | | Marl | grey, shell fragments | | |
| Silt | 0.004-0.06 mm | | | | | |
| Clay | < 0.004 mm (slick) | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME AM PM | REASON FOR SURVEY |

| | Habitat | | Condition | ı Category | | |
|--|---|---|---|---|---|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | |
| ted i | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | |
| uram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Condition | n Category | | | | |
|---|--|--|--|---|--|--|--|
| Parameter | Optimal | Suboptimal | Marginal | Poor | | | |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE(RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME | | LOCATION | | | |
|----------------------|--|--|-------------------|--|--|
| STATION # | _ RIVERMILE | STREAM CLASS | | | |
| LAT | LONG | RIVER BASIN | | | |
| STORET # | | AGENCY | | | |
| INVESTIGATORS | | | LOT NUMBER | | |
| FORM COMPLETED | BY | DATE TIME | REASON FOR SURVEY | | |
| HABITAT TYPES | Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()% | | | | |
| SAMPLE COLLECTION | Indicate the number of jab | lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B | anks Sand | | |
| GENERAL COMMENTS | | | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS


Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)


| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM


| County: | Nicholas | Stream ID: | S-A71 |
|---------------------------|---------------------------------|---------------|---------|
| Stream Name: | UNT to Big Beaver Creek | | |
| HUC Code: Survey Date: | 5050005 9/4/2021 | Basin: | Elk |
| Surveyors: Type: | KP, SM Representative/Riffle | Impact Reach: | 12.19 m |

| | | | LE COUNT | | | | |
|-------------|--------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | ▲ ▼ | 0 | 0.00 | 0.00 |
| | Very Fine | .062125 | | • | 93 | 93.00 | 93.00 |
| | Fine | .12525 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| | Medium | .255 | S A N D | ▲ ▼ | 0 | 0.00 | 93.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .0408 | Very Coarse | 1.0-2 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .0816 | Very Fine | 2 -4 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .1622 | Fine | 4 -5.7 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .2231 | Fine | 5.7 - 8 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .3144 | Medium | 8 -11.3 | GRAVEL | ▲ ▼ | 0 | 0.00 | 93.00 |
| .4463 | Medium | 11.3 - 16 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .6389 | Coarse | 16 -22.6 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | ▲ ▼ | 0 | 0.00 | 93.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | ▲ ▼ | 1 | 1.00 | 94.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ▲ ▼ | 1 | 1.00 | 95.00 |
| 3.5 - 5.0 | Small | 90 - 128 | CODDIE | ▲ ▼ | 1 | 1.00 | 96.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | ▲ ▼ | 0 | 0.00 | 96.00 |
| 7.1 - 10.1 | Large | 180 - 256 | | ▲ ▼ | 1 | 1.00 | 97.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | ▲ ▼ | 0 | 0.00 | 97.00 |
| 14.3 - 20 | Small | 362 - 512 | | ▲ ▼ | 0 | 0.00 | 97.00 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | ▲ ▼ | 2 | 2.00 | 99.00 |
| 40 - 80 | Large | 1024 -2048 | | ▲ ▼ | 0 | 0.00 | 99.00 |
| 80 - 160 | Vry Large | 2048 -4096 | 1 | ▲ ▼ | 1 | 1.00 | 100.00 |
| | Bedrock | | BDRK | • • | 0 | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |
| | Total Tally: | | | | | | |






File C/Users/shite/Teins Tech, Inc/Waltur, David - Sent 00-14-2021/VEI/2021-00-04 - S-A71_S-A71 BINUD SITEMU TOPO_UP 116.1/S-A71 - MP 116.1 - 22/34.4mg Pet David Three Sen 26, 2021 - Beltom



SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON SEPTEMBER 4, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.



NOTE: ALL SECTIONS VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

