Reach S-B26 (Temporary Access Road) Intermittent Spread D Nicholas County, West Virginia

Data	Included
Photos	\checkmark
SWVM Form	\checkmark
FCI Calculator and HGM Form	NA – slope <4%
RBP Physical Characteristics Form	\checkmark
Water Quality Data	\checkmark
RBP Habitat Form	\checkmark
RBP Benthic Form	\checkmark
Benthic Identification Sheet	N/A – Lack of habitat
Wolman Pebble Count	\checkmark
Reference Reach Software Pebble Count Data	\checkmark
Longitudinal Profile and Cross Sections	\checkmark

Spread D Stream S-B26 (Pipeline ROW) Webster County



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, CNJ Lat: 38.339012 Long: -80.659609



Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, CNJ Lat: 38.339012 Long: -80.659609

Spread D Stream S-B26 (Pipeline ROW) Webster County



Photo Type: US View at Center Location, Orientation, Photographer Initials: Center ROW, Upstream View, CNJ Lat: 38.339012 Long: -80.659609



Photo Type: DS View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, CNJ Lat: 38.339012 Long: -80.659609

Spread D Stream S-B26 (Pipeline ROW) Webster County



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, CNJ Lat: 38.339012 Long: -80.659609



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, CNJ Lat: 38.339012 Long: -80.659609

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

USACE FILE NO./ Project Name: (v2.1, Sept 2015)	Mountain Valley Pipeline			IMPACT COORDINATES (in Decimal Degrees)	: Lat.	38.339012	Lon.	-80.659609	WEATHER:	Sunny	DATE:	9/10/2021
IMPACT STREAM/SITE ID AN (watershed size (acreage), una		ON:	S-1	B26		MITIGATION STREAM CLASS (watershed size (acreag				<u>-</u>	Comments:	
STREAM IMPACT LENGTH:		ORM OF TIGATION:	RESTORATION (Levels I-III)			PRECIPITATION PAST 48 HRS:		Mitigation Length:				
Column No. 1- Impact Existing Co	ondition (Debit)		Column No. 2- Mitigation Existing Co	ondition - Baseline (Credit)		Column No. 3- Mitigation P Post Completion	rojected at Five on (Credit)	Years	Column No. 4- Mitigation Proj Post Completion (ected at Ten Years Credit)	Column No. 5- Mitigation Project	ed at Maturity (Credit)
Stream Classification:	Intermittent		Stream Classification:			Stream Classification:		0	Stream Classification:	0	Stream Classification:	0
Percent Stream Channel Slope	e 1.4	L .	Percent Stream Channel Slo	ope		Percent Stream Channel S	Slope	0	Percent Stream Channel SI	ope 0	Percent Stream Channel S	ope 0
HGM Score (attach data	a forms):		HGM Score (attach o	data forms):		HGM Score (attacl	h data forms):		HGM Score (attach da	ata forms):	HGM Score (attach d	ata forms):
Hydrology Biogeochemical Cycling Habitat	Averag		Hydrology Biogeochemical Cycling Habitat	0		Hydrology Biogeochemical Cycling Habitat		Average 0	Hydrology Biogeochemical Cycling Habitat	Average 0	Hydrology Biogeochemical Cycling Habitat	0
PART I - Physical, Chemical and Bio	ological Indicators	570 ST	PART I - Physical, Chemical and	Pains Scale Range Site Score		PART I - Physical, Chemical a	-		PART I - Physical, Chemical and	Points Scale Range Site Score	PART I - Physical, Chemical and	Points Scale Range Sile Scale
PHYSICAL INDICATOR (Applies to all streams class	assifications)		PHYSICAL INDICATOR (Applies to all streams of	classifications)		PHYSICAL INDICATOR (Applies to all stream	is classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)	PHYSICAL INDICATOR (Applies to all streams	classifications)
USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover	0-20 3		USEPA RBP (Low Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover	0-20	_	USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover	0-20		USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover	0-20	USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover	0-20
4. Sediment Deposition 5. Channel Flow Status 6. Channel Alteration 7. Frequency of Kiffles (or bends) 9. Bank Stability (LB & RB) 10. Repair Vegetative Zone Wich (LB & RB) 10. Repair Vegetative Zone Wich (LB & RB) CHEMICAL INDICATOR (Applies to intermitter an WDCFP Water Quality Indicators (General) Specific Conductivity 100-199 - 85 points pH	0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	5	3. Pool Vantability 4. Sediment Deposition 4. Channel Flow Status 6. Channel Flow Status 6. Channel Flow Status 6. Channel Stratation 7. Channel Sanuaghy 8. Anst Stability (LB & RB) 9. Vegetimities Protection (LB & RB) 10. Requirely regetative Znm Widher (LB & RB) Total RBP Score Sup-Total CHEMICAL INDICATOR (Applies to Interminent WDEP Water Quality Indicators (General) Specific Conductivity pH	0-20 0-20 0-20 0-20 0-20 0-20 0-20 0-20		3. Velocity Depth Regime 4. Sediment Departion 5. Channel Flow Status 6. Channel Flow Status 6. Channel Remarks 7. Frequency of Riffles (or bends) 7. Frequency of Riffles (or bends) 9. Vegetative Protection (LB & RB) 10. Repairs Vegetative Zore Widh (LB & RB) 20. Vegetat		0 0 reams)	3. Velocity' Depth Regime 4. Sediment Depatition 5. Channel Flow Status 6. Channel Karation 7. Fraesency of Riffel (or bends) 8. Brain's Statibility (LB & RB) 10. Rigation Vegatietic Zore Width (LB & RB) CHEIMICAL INDICATOR (Applies to Internite WDDEP Water Quality Indicators (General Specific Conductivity pH	Poor 0 0 tt and Perennial Streams)	Subiolity Depth Regime Settimer Deposition Channel Flow Status Channel Flow Status Channel Alteration Trequency of Riffies for binds) Sark Status (LB & RB) Superstative Protection (LB & RB) Total RBP Score Sub-Total Refunct INDICATOR (Applies to Intermitter WDEP Water Quality Indicators (General Specific Conductivity pH	
DO >5.0 = 30 points 1 Sub-Total BIOLOGICAL INDICATOR (Applies to intermittent WY Stream Condition Index (WVSCI)	10-30 5.1 0.625 t and Perennial Streams)	25	DO Sub-Total BIOLOGICAL INDICATOR (Applies to Intermitte WV Stream Condition Index (WVSCI)	10-30 0 ent and Perennial Streams)		DO Sub-Total BIOLOGICAL INDICATOR (Applies to Inter WV Stream Condition Index (WVSCI)	10-30	0 nial Streams)	DO Sub-Total BIOLOGICAL INDICATOR (Applies to Intern WV Stream Condition Index (WVSCI)	10-30 0 ittent and Perennial Streams)	DO Sub-Total BIOLOGICAL INDICATOR (Applies to Intern WV Stream Condition Index (WVSCI)	10-30 0 ittent and Perennial Streams)
0 C	0-100 0-1		Sub-Total	0-100 0-1 0	_	Sub-Total	0-100 0-1	0	Sub-Total	0-100 0-1 0	Sub-Total	0-100 0-1 0
PART II - Index and Unit	: Score		PART II - Index and I	Unit Score		PART II - Index an	d Unit Score		PART II - Index and U	nit Score	PART II - Index and U	nit Score
Index L	Linear Feet Unit Sc	core	Index	Linear Feet Unit Score		Index	Linear Feet	Unit Score	Index	Linear Feet Unit Score	Index	Linear Feet Unit Score
0.495	43 21.28	85	0	0 0		0	0	0	0	0 0	0	0 0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME	LOCATION						
STATION # RIVERMILE	STREAM CLASS						
LAT LONG	RIVER BASIN						
STORET #	AGENCY						
INVESTIGATORS							
FORM COMPLETED BY	DATE TIME	REASON FOR SURVEY					

WEATHER CONDITIONS	Now Past 24 hours Has there been a heavy rain in the last 7 days? storm (heavy rain) rain (steady rain) showers (intermittent) Yes No % %cloud cover clear/sunny Air Temperature0 C
SITE LOCATION/MAP	Draw a map of the site and indicate the areas sampled (or attach a photograph)
STREAM CHARACTERIZATION	Stream Subsystem Stream Type Perennial Intermittent Tidal Stream Origin Spring-fed Glacial Spring-fed Non-glacial montane Spring-fed Swamp and bog Other

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer)	Predominant Surrounding Landuse Local Watershed NPS Pollution Forest Commercial Field/Pasture Industrial Agricultural Other Residential Other Indicate the dominant type and record the dominant species present Herbaceous Trees Shrubs Grasses Dominant species present Herbaceous
INSTREAM FEATURES	Dominant species present
LARGE WOODY	LWDm ²
DEBRIS	Density of LWDm ² /km ² (LWD/ reach area)
AQUATIC	Indicate the dominant type and record the dominant species present
VEGETATION	Rooted emergent Rooted submergent Rooted floating Free floating Floating Algae Attached Algae Booted floating Free floating Free floating Dominant species present
WATER QUALITY (DS, US)	Temperature0 C Water Odors Normal/None Sewage Specific Conductance Petroleum Fishy Chemical Other Dissolved Oxygen Water Surface Oils Slick Sheen None Globs Flecks pH Turbidity (if not measured) Clear Slightly turbid Turbid Turbid Turbid Opaque Turbid
SEDIMENT/	Odors
SUBSTRATE	Normal Sewage Petroleum Deposits Chemical Anaerobic None Sludge Sawdust Paper fiber Sand Other Other Epoking at stones which are not deeply embedded are the undersides black in color? How are the undersides black in color?

INC	ORGANIC SUBSTRATE (should add up to			ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)							
Substrate Type	Diameter	% Composition in Sampling Reach	Substrate Type	Characteristic % Compositio Sampling Ar							
Bedrock			Detritus	sticks, wood, coarse plant							
Boulder	> 256 mm (10")			materials (CPOM)							
Cobble	64-256 mm (2.5"-10")		Muck-Mud	black, very fine organic							
Gravel	2-64 mm (0.1"-2.5")			(FPOM)							
Sand	0.06-2mm (gritty)		Marl	grey, shell fragments							
Silt	0.004-0.06 mm										
Clay	< 0.004 mm (slick)										

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION						
STATION # RIVERMILE	STREAM CLASS						
LAT LONG	RIVER BASIN	RIVER BASIN					
STORET #	AGENCY						
INVESTIGATORS							
FORM COMPLETED BY	DATE TIME AM PM	REASON FOR SURVEY					

	Habitat		Condition	ı Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.				
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.				
ted i	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).				
uram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
Pa	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.				
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.				
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

Habitat		Condition	ı Category				
Parameter	Optimal	Suboptimal	Marginal	Poor			
6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.			
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.			
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
 SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) 	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.			
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.			
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.			
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
SCORE(RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME		LOCATION							
STATION #	_ RIVERMILE	STREAM CLASS							
LAT	LONG	RIVER BASIN							
STORET #		AGENCY							
INVESTIGATORS			LOT NUMBER						
FORM COMPLETED	BY	DATE TIME	REASON FOR SURVEY						
HABITAT TYPES	Cobble% Sn	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%							
SAMPLE COLLECTION	Indicate the number of jab	lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B	anks Sand						
GENERAL COMMENTS									

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						


WOLMAN PEBBLE COUNT FORM

County: Stream Name: HUC Code: Survey Date: Surveyors: Type: Nicholas UNT to Cherry Run 5050005 9/10/2021 CCC, CNJ Bankfull Channel Stream ID: S-B26

Basin:

Impact: 10.92

PEBBLE COUNT							
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	▲ ▼	25	25.00	25.00
	Very Fine	.062125		▲ ▼	69	69.00	94.00
	Fine	.12525		▲ ▼	0	0.00	94.00
	Medium	.255	S A N D	▲ ▼	0	0.00	94.00
	Coarse	.50-1.0		▲ ▼	0	0.00	94.00
.0408	Very Coarse	1.0-2		▲ ▼	0	0.00	94.00
.0816	Very Fine	2 -4		▲ ▼	0	0.00	94.00
.1622	Fine	4 -5.7		▲ ▼	0	0.00	94.00
.2231	Fine	5.7 - 8		▲ ▼	0	0.00	94.00
.3144	Medium	8 -11.3		▲ ▼	0	0.00	94.00
.4463	Medium	11.3 - 16	GRAVEL	▲ ▼	0	0.00	94.00
.6389	Coarse	16 -22.6		▲ ▼	0	0.00	94.00
.89 - 1.26	Coarse	22.6 - 32		▲ ▼	1	1.00	95.00
1.26 - 1.77	Vry Coarse	32 - 45		▲ ▼	0	0.00	95.00
1.77 -2.5	Vry Coarse	45 - 64		▲ ▼	0	0.00	95.00
2.5 - 3.5	Small	64 - 90		▲ ▼	0	0.00	95.00
3.5 - 5.0	Small	90 - 128		▲ ▼	1	1.00	96.00
5.0 - 7.1	Large	128 - 180	COBBLE	▲ ▼	0	0.00	96.00
7.1 - 10.1	Large	180 - 256		▲ ▼	1	1.00	97.00
10.1 - 14.3	Small	256 - 362	1	▲ ▼	3	3.00	100.00
14.3 - 20	Small	362 - 512	BOULDER	▲ ▼	0	0.00	100.00
20 - 40	Medium	512 - 1024		▲ ▼	0	0.00	100.00
40 - 80	Large	1024 -2048		▲ ▼	0	0.00	100.00
80 - 160	Vry Large	2048 -4096		▲ ▼	0	0.00	100.00
	Bedrock		BDRK	▲ ▼	0	0.00	100.00
				Totals:	100		
	Total Tally:						





cumulative %

-

■# of partickes


Size (mm) 💦 🎽	Size Distribution	Туре
D16 0.062	mean 0.1	silt/clay 25%
D35 0.069	dispersion 1.3	sand 69%
D50 0.08	skewness 0.03	gravel 1%
D65 0.093		cobble 2%
D84 0.11		boulder 3%
D95 32		





SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON SEPTEMBER 10, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.



NOTE: ALL SECTIONS VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

