Reach S-F34 (Timber Mat Crossing) Perennial Spread D Webster County, West Virginia

Data	Included
Photos	\checkmark
SWVM Form	✓Water quality readings from benthic sampling
	date
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable)
RBP Physical Characteristics Form	\checkmark
Water Quality Data	\checkmark
RBP Habitat Form	\checkmark
RBP Benthic Form	\checkmark
Benthic Identification Sheet	✓ Sampling date 9/14/2021
Wolman Pebble Count	\checkmark
Reference Reach Software Pebble Count Data	\checkmark
Longitudinal Profile and Cross Sections	\checkmark

Spread D Stream S-F34 (Timber Mat Crossing) Webster County




Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068



Spread D Stream S-F34 (Timber Mat Crossing) Webster County

Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068



Spread D Stream S-F34 (Timber Mat Crossing) Webster County

Photo Type: DS View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068

Spread D Stream S-F34 (Timber Mat Crossing) Webster County



Photo Type: US View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068



Spread D Stream S-F34 (Timber Mat Crossing) Webster County

Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068



Spread D Stream S-F34 (Timber Mat Crossing) Webster County

Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068





Photo Type: Riffle, DS View Location, Orientation, Photographer Initials: Upstream of Riffle, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068






Photo Type: Riffle, US View Location, Orientation, Photographer Initials: Downstream of Riffle, Upstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068





Photo Type: Pool, DS View Location, Orientation, Photographer Initials: Upstream of Pool, Downstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068

Spread D Stream S-F34 (Timber Mat Crossing) Webster County



Photo Type: Pool, US View Location, Orientation, Photographer Initials: Downstream of Pool, Upstream View, DD/LC/KP Lat: 38.423988 Long: -80.57068

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

	Mountain Valley Pipeline		(in Decimal Degrees)							08/28/21
IMPACT STREAM/SITE ID ANI (watershed size (acreage), unalt		S-F34			MITIGATION STREAM CLASS./SITE ID AN (watershed size (acreage), unalitered or				Comments:	
STREAM IMPACT LENGTH:	22 FORM OF MITIGATION:	RESTORATION (Levels I-III)	MIT COORDINATES: (in Decimal Degrees)	Lat.	Lon.		PRECIPITATION PAST 48 HRS:		Mitigation Length:	
Column No. 1- Impact Existing Col	ondition (Debit)	Column No. 2- Mitigation Existing Cond	ition - Baseline (Credit)		Column No. 3- Mitigation Projected at F Post Completion (Credit)	Five Years	Column No. 4- Mitigation Proje Post Completion (C		Column No. 5- Mitigation Project	ed at Maturity (Credit)
Stream Classification:	Perennial	Stream Classification:			Stream Classification:	0	Stream Classification:	0	Stream Classification:	0
Percent Stream Channel Slope	e 1.9	Percent Stream Channel Slope			Percent Stream Channel Slope	0	Percent Stream Channel Slo	ope O	Percent Stream Channel SI	ope 0
HGM Score (attach data f	forms):	HGM Score (attach data	ı forms):		HGM Score (attach data form:	s):	HGM Score (attach da	ta forms):	HGM Score (attach da	ata forms):
	Average		Average			Average		Average		Average
Hydrology		Hydrology			Hydrology		Hydrology		Hydrology	
Biogeochemical Cycling Habitat	0	Biogeochemical Cycling Habitat	0		Biogeochemical Cycling Habitat	0	Biogeochemical Cycling Habitat	0	Biogeochemical Cycling Habitat	0
PART I - Physical, Chemical and Biol	ological Indicators	PART I - Physical, Chemical and Bi	ological Indicators		PART I - Physical, Chemical and Biologica	al Indicators	PART I - Physical, Chemical and E	Biological Indicators	PART I - Physical, Chemical and	Biological Indicators
Pair	oints Scale Range Site Score	Pair	ta Scale Range Sille Score		Points Scale	Range Site Score		Points Scale Range Site Score		Points Scale Range Site Score
PHYSICAL INDICATOR (Applies to all streams class	ssifications)	PHYSICAL INDICATOR (Applies to all streams class	ifications)		PHYSICAL INDICATOR (Applies to all streams classification	16)	PHYSICAL INDICATOR (Applies to all streams	classifications)	PHYSICAL INDICATOR (Applies to all streams	classifications)
USEPA RBP (High Gradient Data Sheet)		USEPA RBP (Low Gradient Data Sheet)			USEPA RBP (High Gradient Data Sheet)		USEPA RBP (High Gradient Data Sheet)		USEPA RBP (High Gradient Data Sheet)	
	0-20 14 0-20 13		0-20		1. Epifaunal Substrate/Available Cover 0-20 2. Embeddedness 0-20		1. Epifaunal Substrate/Available Cover 2. Embeddedness	0-20	1. Epifaunal Substrate/Available Cover 2. Embeddedness	0-20
	0-20 8		0-20		3. Velocity/ Depth Regime 0-20		3. Velocity/ Depth Regime	0-20	3. Velocity/ Depth Regime	0-20
	0-20 16		0-20		4. Sediment Deposition 0-20		4. Sediment Deposition	0-20	4. Sediment Deposition	0-20
	0-20 0-1 13	5. Channel Flow Status	0-20 0-1		5. Channel Flow Status 0-20	0-1	5. Channel Flow Status	0-20 0-1	5. Channel Flow Status	0-20 0-1
	0-20 19		0-20		6. Channel Alteration 0-20		6. Channel Alteration	0-20	6. Channel Alteration	0-20
	0-20 18		0-20		7. Frequency of Riffles (or bends) 0-20		7. Frequency of Riffles (or bends)	0-20	Frequency of Riffles (or bends)	0-20
	0-20 18		0-20		8. Bank Stability (LB & RB) 0-20		8. Bank Stability (LB & RB)	0-20	8. Bank Stability (LB & RB)	0-20
9. Vegetative Protection (LB & RB) 0	0-20 16		0-20		9. Vegetative Protection (LB & RB) 0-20		9. Vegetative Protection (LB & RB)	0-20	9. Vegetative Protection (LB & RB)	0-20
	0-20 10 Suboptimal 145	10. Riparian Vegetative Zone Width (LB & RB) Total RBP Score	Poor 0		10. Riparian Vegetative Zone Width (LB & RB) 0.20 Total RBP Score Poor	0	10. Riparian Vegetative Zone Width (LB & RB) Total RBP Score	0-20 Poor 0	10. Riparian Vegetative Zone Width (LB & RB) Total RBP Score	0-20 Poor 0
Sub-Total	0.725	Sub-Total	Poor U		Sub-Total	0	Sub-Total	Poor U	Sub-Total	Poor U
CHEMICAL INDICATOR (Applies to Intermittent and		CHEMICAL INDICATOR (Applies to Intermittent and			CHEMICAL INDICATOR (Applies to Intermittent and Perenni	-	CHEMICAL INDICATOR (Applies to Intermittent		CHEMICAL INDICATOR (Applies to Intermitten	
WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (General)			WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (General)	
Specific Conductivity		Specific Conductivity			Specific Conductivity		Specific Conductivity		Specific Conductivity	
<=99 - 90 points	0-90 37.2		0-90		0-90			0-90		0-90
nH		nH			nH		nH		nH	
	0-80 0-1 6.77	,	5-90 0-1		5-90	0-1	p.,	5-90 0-1		5-90 0-1
6.0-8.0 = 80 points	0.77									
DO		DO			DO		DO		DO	
>5.0 = 30 points	10-30 7	1	0-30		10-30			10-30		10-30
Sub-Total	1	Sub-Total	0		Sub-Total	0	Sub-Total	0	Sub-Total	0
BIOLOGICAL INDICATOR (Applies to Intermittent a	and Perennial Streams)	BIOLOGICAL INDICATOR (Applies to Intermittent an	d Perennial Streams)		BIOLOGICAL INDICATOR (Applies to Intermittent and Pe	erennial Streams)	BIOLOGICAL INDICATOR (Applies to Intermi	ttent and Perennial Streams)	BIOLOGICAL INDICATOR (Applies to Interm	ittent and Perennial Streams)
WV Stream Condition Index (WVSCI)		WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)		WV Stream Condition Index (WVSCI)		WV Stream Condition Index (WVSCI)	
•	0-100 0-1 70.2	0	-100 0-1		0-100	0-1		0-100 0-1		0-100 0-1
Good Sub-Total	0.702	Sub-Total	0		Sub-Total	0	Sub-Total	0	Sub-Total	0
PART II - Index and Unit S		PART II - Index and Unit			PART II - Index and Unit Score	•	PART II - Index and Ur		PART II - Index and U	nit Score
Index L	Linear Feet Unit Score	Index L	inear Feet Unit Score		Index Linear F	Feet Unit Score	Index	Linear Feet Unit Score	Index	Linear Feet Unit Sco
0.809	22 17.798	0	0 0		0 0	0	0	0 0	0	0 0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME	LOCATION						
STATION # RIVERMILE	STREAM CLASS	STREAM CLASS					
LAT LONG	RIVER BASIN	RIVER BASIN					
STORET #	AGENCY						
INVESTIGATORS							
FORM COMPLETED BY	DATE TIME	REASON FOR SURVEY					

WEATHER CONDITIONS SITE LOCATION/MAP	Now Past 24 hours Has there been a heavy rain in the last 7 days? Storm (heavy rain) rain (steady rain) showers (intermittent) % %cloud cover clear/sunny Has there been a heavy rain in the last 7 days? Yes No Air Temperature0 C Other Draw a map of the site and indicate the areas sampled (or attach a photograph)
	S-F34 and flow direction Re-seeded grass Access road Timber mat North
STREAM CHARACTERIZATION	Stream Subsystem Stream Type Perennial Intermittent Tidal Stream Origin Coldwater Warmwater Glacial Spring-fed Catchment Areakm² Non-glacial montane Mixture of origins Swamp and bog Other

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer)	Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential	Local Watershed NPS Pollution No evidence Some potential sources Obvious sources Local Watershed Erosion None Moderate Heavy ant species present Grasses Herbaceous
INSTREAM FEATURES	Dominant species present	Canopy Cover Partly open Partly shaded Shaded High Water Mark m Proportion of Reach Represented by Stream Morphology Types Riffle % Riffle % Pool % Channelized Yes No No
LARGE WOODY DEBRIS	LWDm ² Density of LWDm ² /km ² (LWD/ reac	h area)
AQUATIC VEGETATION	Indicate the dominant type and record the dominant record the dominant type and record the domin Rooted submergent Rooted submergent Attached Algae Dominant species present Portion of the reach with aquatic vegetation	Rooted floating Free floating
WATER QUALITY (DS, US)	Temperature0 C Specific Conductance Dissolved Oxygen pH Turbidity WQ Instrument Used	Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Slick Sheen Globs Flecks None Other Turbidity (if not measured) Clear Slightly turbid Clear Slightly turbid Turbid Opaque Stained Other
SEDIMENT/ SUBSTRATE	Odors Petroleum Normal Sewage Petroleum Chemical Anaerobic None Other	Deposits Paper fiber Sand Sludge Sawdust Paper fiber Sand Relict shells Other

INC	DRGANIC SUBSTRATE (should add up to			ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)							
Substrate Type	Diameter	% Composition in Sampling Area									
Bedrock			Detritus	sticks, wood, coarse plant							
Boulder	> 256 mm (10")			materials (CPOM)							
Cobble	64-256 mm (2.5"-10")		Muck-Mud	black, very fine organic							
Gravel	2-64 mm (0.1"-2.5")			(FPOM)							
Sand	0.06-2mm (gritty)		Marl	grey, shell fragments							
Silt	0.004-0.06 mm										
Clay	< 0.004 mm (slick)										

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION						
STATION # RIVERMILE	STREAM CLASS						
LAT LONG	RIVER BASIN	RIVER BASIN					
STORET #	AGENCY						
INVESTIGATORS							
FORM COMPLETED BY	DATE TIME AM PM	REASON FOR SURVEY					

	Habitat		Condition	ı Category						
	Parameter	Optimal	Suboptimal	Marginal	Poor					
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.					
ted iı	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).					
Iram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
P	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

Habitat		Condition	ı Category	
Parameter	Optimal	Suboptimal	Marginal	Poor
6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
 SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) 	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0
9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0
10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE(RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-F	34	LOCATION Webster County								
STATION #	RIVERMILE	STREAM CLASS Perennial								
LAT 38.423988	LONG80.57068	RIVER BASIN None	RIVER BASIN None							
STORET #		AGENCY WVDEP								
INVESTIGATORS P	= SM		LOT NUMBER							
FORM COMPLETED	^{BY} SM	DATE 9/14/2021 TIME 1315	REASON FOR SURVEY Baseline Assessment							
	r									
HABITAT TYPES	Cobble <u>20</u> %	dicate the percentage of each habitat type present Cobble 20 % □Snags% □Vegetated Banks% □Sand% Submerged Macrophytes% □Other ()%								
SAMPLE COLLECTION		lected? ☑ wading ☐ fi s/kicks taken in each habitat ty ags ☐ Vegetated B	rom bank							
GENERAL COMMENTS			DO: 6.14 mg/L, pH: 6.87 , DO: 7.00 mg/L, pH: 6.77							

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)


Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera		1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

Insects	Count T	Tolerance	Z	Insects	Count	Tolerance	≥	Non-Insects	Count	Tolerance	2	SITE ID:	S-F34
Ephemeroptera			18	Odonata			4	Crustacea			0		9/14/2021
Ameletidae		2	0	Aeshnidae	1	3	'n	Asellidae		7	0		
Baetidae		4	0	Calopterygidae	2	9	12	Cambaridae		5	0		
Beatiscidae		4	0	Coenagrionidae		7	0	Gammaridae		5	0		
Caenidae		5	0	Cordulegastridae		3	0	Palaemonidae		5	0		
Ephemerellidae	12	3	36	Gomphidae	1	5	2	Annelida			0		
Ephemeridae		5	0	Lestidae		۲	0	Hirudinea		10	0		
Heptageniidae	9	з	18	Libellulidae		۲	0	Nematoda		10	0		
Isonychiidae		æ	0	Coleoptera			1	Nematomorpha		10	0		
Leptophlebiidae		4	0	Chrysomelidae		7	0	Oligochaeta		10	0		
Potamanthidae		5	0	Dryopidae		5	0	Turbellaria			0		
Siphlonuridae		æ	0	Dytiscidae		9	0	Turbellaria		7	0		
Tricorythidae		5	0	Elmidae		4	0	Bivalvia			0		
Plecoptera			27	Gyrinidae		5	0	Corbiculidae		6	0		
Capniidae		2	0	Haliplidae		7	0	Sphaeriidae		5	0		
Chloroperlidae		2	0	Hydrophilidae	1	7	7	Unionidae		4	0		
Leuctridae	21	2	42	Psephenidae		8	0	Gastropoda			0		
Nemouridae		2	0	Ptilodactylidae		5	0	Ancylidae		7	0		
Peltoperlidae		H	0	Hemiptera			0	Hydrobiidae		4	0		
Perlidae		н,	0	Belostomatidae		8	0	Physidae		7	0		
Perlodidae	9	1	9	Corixidae		8	0	Planorbidae		5	0		
Pteronarcyidae		1	0	Gerridae		10	0	Pleuroceridae		5	0		
Taeniopterygidae		2	0	Hydrometridae		8	0	Viviparidae		5	0		
Trichoptera			14	Nepidae		8	0	Miscellaneous			7		
Brachycentridae		2	0	Notonectidae		8	0	Collembola	7	6	42		
Glossosomatidae		2	0	Megaloptera			1	Lepidoptera		5	0		
Helicopsychidae		3	0	Corydalidae		3	0	Neuroptera		5	0		
Hydropsychidae	8	5	40	Sialidae	1	9	9	Hydrachnidae		6	0		
Hydroptilidae		з	0	Diptera			58	Totals	Total	Total number	130		
Lepidostomatidae		3	0	Athericidae		3	0	618101	Total	Total families	18		
Leptoceridae		3	0	Blephariceridae		2	0			Σ	Metric calculations	lations	
Limnephilidae		4	0	Ceratopogonidae	7	8	56	SIVIN	MNSCI Matric Scores	Scores		Additional metrics	S
Molannidae		З	0	Chironomidae	32	6	288			2000		Ephemeroptera Taxa	2
Philopotamidae	1	4	4	Culicidae		10	0	Total Taxa		18	81.8	Plecoptera Taxa	2
Phryganeidae		4	0	Dixidae	5	6	30	ЕРТ Таха		7	53.8	Trichoptera Taxa	3
Polycentropodidae		5	0	Empididae		7	0	% EPT Abundance	ance	45.4	50.8	Long-lived Taxa	11
Psychomiidae		4	0	Psychodidae		8	0	% Chironomidae	dae	24.6	76.7	Odonata Taxa	3
Rhyacophilidae	5	3	15	Ptychopteridae		8	0	Hilsenhoff Biotic Index (HBI)	idex (HBI)	5.29	63.7	Diptera Taxa	5
Uenoidae		2	0	Simuliidae		7	0	% 2 Dominant Taxa	Таха	40.8	94.5	СОЕТ Таха	6
To	Total Tolerance Value	nce Value	688	Stratiomyidae		10	0					% Sensitive	40.0
West Virginia Stream Condition Index (WVSCI)	Condition	Index (W	/SCI)	Syrphidae		10	0					% Tolerant	33.8
Gerritson, J., J. Burton, and M.T. Barbour. 2000. A stream	A.T. Barbou	r. 2000. A 5 bla <i>ctream</i> c	tream	Tabanidae	4	7	28	WV Stream Condition Index	Condition	Index	70.2	% Clingers	34.6
conduction maex for west virginia wadeable su earns. Tetra Tech, Inc. Owing Mills, MD.	gilila wauca	מווב את במוויאי	-	Tipulidae	10	5	50					% Net-spinners	6.9
Spreadsheet uses updated Best Standard Values [BSV] for each metric per WVSCI Addenda dated March 23, 2010	est Standar	d Values [BS	אן for eac	h metric per WVSCI Add	enda dated	March 23, 201	0.						

WOLMAN PEBBLE COUNT FORM

County:	Webster	Stream ID:	S-F34
Stream Name:	UNT to Birch River		
HUC Code:		Basin:	
Survey Date:	8/28/2021		
Surveyors:	DD LC KP	Impact Reach:	22.6 m
Type:	Bankfull Channel		

			LE COUNT		-		
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	•	12	12.00	12.00
	Very Fine	.062125		•	1	1.00	13.00
	Fine	.12525		•	6	6.00	19.00
	Medium	.255	S A N D	▲ ▼	7	7.00	26.00
	Coarse	.50-1.0		* *	3	3.00	29.00
.0408	Very Coarse	1.0-2		▲ ▼	0	0.00	29.00
.0816	Very Fine	2 -4		▲ ▼	0	0.00	29.00
.1622	Fine	4 -5.7		▲ ▼	4	4.00	33.00
.2231	Fine	5.7 - 8		▲ ▼	4	4.00	37.00
.3144	Medium	8 -11.3		▲ ▼	7	7.00	44.00
.4463	Medium	11.3 - 16	G R A V E L	▲ ▼	17	17.00	61.00
.6389	Coarse	16 -22.6		▲ ▼	12	12.00	73.00
.89 - 1.26	Coarse	22.6 - 32		▲ ▼	4	4.00	77.00
1.26 - 1.77	Vry Coarse	32 - 45		▲ ▼	2	2.00	79.00
1.77 -2.5	Vry Coarse	45 - 64		▲ ▼	2	2.00	81.00
2.5 - 3.5	Small	64 - 90		▲ ▼	6	6.00	87.00
3.5 - 5.0	Small	90 - 128	COBBLE	▲ ▼	3	3.00	90.00
5.0 - 7.1	Large	128 - 180	COBBLE	▲ ▼	5	5.00	95.00
7.1 - 10.1	Large	180 - 256		▲ ▼	2	2.00	97.00
10.1 - 14.3	Small	256 - 362	BOULDER	•	1	1.00	98.00
14.3 - 20	Small	362 - 512		▲ ▼	0	0.00	98.00
20 - 40	Medium	512 - 1024		▲ ▼	0	0.00	98.00
40 - 80	Large	1024 -2048		▲ ▼	0	0.00	98.00
80 - 160	Vry Large	2048 -4096		▲ ▼	0	0.00	98.00
	Bedrock		BDRK	▲ ▼	2	2.00	100.00
				Totals:	100		
	Total Tally:						





cumulative %

of particles

Bankfull Channel Pebble Count, S-F34, UNT to Birch River


percent finer than





SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON AUGUST 28, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.



NOTE: ALL SECTIONS VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

