Reach S-H113 (1) (Pipeline ROW) Perennial Spread C Webster County, West Virginia

| Data | Included |
|--|--|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | N/A – Perennial stream (not shadeable) |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | ✓ Readings from benthic sampling date |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | ✓ Sampling date 9/21/21 |
| Wolman Pebble Count | \checkmark |
| Reference Reach Software Pebble Count Data | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, VM Lat: 38.612982, Long: -80.503647



Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, VM Lat: 38.612982, Long: -80.503647



Photo Type: US View at Center Location, Orientation, Photographer Initials: Center ROW, Upstream View, VM Lat: 38.612982, Long: -80.503647



Photo Type: DS View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, VM Lat: 38.612982, Long: -80.503647




Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, HK Lat: 38.612982, Long: -80.503647



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, HK Lat: 38.612982, Long: -80.503647




Photo Type: Riffle, DS View Location, Orientation, Photographer Initials: Upstream of Riffle, Downstream View, VM Lat: 38.612982, Long: -80.503647



Photo Type: Riffle, US View Location, Orientation, Photographer Initials: Downstream of Riffle, Upstream View, VM Lat: 38.612982, Long: -80.503647




Photo Type: Pool, DS View Location, Orientation, Photographer Initials: Upstream of Pool, Downstream View, VM Lat: 38.612982, Long: -80.503647




Photo Type: Pool, US View Location, Orientation, Photographer Initials: Downstream of Pool, Upstream View, VM Lat: 38.612982, Long: -80.503647

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | Mountain | Valley Pipeline | IMPACT COORDINATES: (in Decimal Degrees) | Lat. | 38.612982 Lon. | -80.503647 | WEATHER: | 50% cloud cover | DATE: | 09/21 | 1/21 |
|--|------------------------------|---|---|------|--|----------------------|--|--------------------------------|--|----------------------------------|--------------|
| IMPACT STREAM/SITE ID AN (watershed size (acreage), una | | S-H113 (1) P | ipeline ROW | | MITIGATION STREAM CLASS./SITE ID (watershed size (acreage), unalter | | | | Comments: | Date used from sampling date. | |
| STREAM IMPACT LENGTH: | 74 FORM OF MITIGATION: | RESTORATION (Levels I-III) | MIT COORDINATES: (in Decimal Degrees) | Lat. | Lon. | | PRECIPITATION PAST 48 HRS: | | Mitigation Length: | | |
| Column No. 1- Impact Existing Co | ondition (Debit) | Column No. 2- Mitigation Existing Co | ondition - Baseline (Credit) | | Column No. 3- Mitigation Projected Post Completion (Cred | at Five Years t) | Column No. 4- Mitigation Proj Post Completion (| ected at Ten Years Credit) | Column No. 5- Mitigation Proje | cted at Maturity (Cr | Credit) |
| Stream Classification: | Perennial | Stream Classification: | | | Stream Classification: | 0 | Stream Classification: | 0 | Stream Classification: | 0 | 0 |
| Percent Stream Channel Slope | e 10.1 | Percent Stream Channel Slo | pe | | Percent Stream Channel Slope | 0 | Percent Stream Channel Si | ope 0 | Percent Stream Channel | Slope | 0 |
| HGM Score (attach data | forms): | HGM Score (attach d | ata forms): | | HGM Score (attach data fo | orms): | HGM Score (attach da | ata forms): | HGM Score (attach | data forms): | |
| | Average | | Average | | | Average | | Average | | | Average |
| Hydrology Biogeochemical Cycling | | Hydrology Biogeochemical Cycling | 0 | | Hydrology Biogeochemical Cycling | 0 | Hydrology Biogeochemical Cycling | | Hydrology Biogeochemical Cycling | | 0 |
| Habitat | | Habitat | | | Habitat | | Habitat | | Habitat | | - ° |
| PART I - Physical, Chemical and Bio | blogical Indicators | PART I - Physical, Chemical and | Biological Indicators | | PART I - Physical, Chemical and Biolo | gical Indicators | PART I - Physical, Chemical and | Biological Indicators | PART I - Physical, Chemical ar | nd Biological Indica | ators |
| P | oints Scale Range Site Score | | Points Scale Range Site Score | | Points Sc | ée Range Site Score | | Points Scale Range Site Score | | Points Scale Range | Site Score |
| PHYSICAL INDICATOR (Applies to all streams clar | ssifications) | PHYSICAL INDICATOR (Applies to all streams cl | assifications) | | PHYSICAL INDICATOR (Applies to all streams classific | ations) | PHYSICAL INDICATOR (Applies to all streams | s classifications) | PHYSICAL INDICATOR (Applies to all stream | ms classifications) | |
| USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (Low Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (High Gradient Data Sheet) | | |
| | 0-20 15 | 1. Epifaunal Substrate/Available Cover | 0-20 | | 1. Epifaunal Substrate/Available Cover 0-20 | | 1. Epifaunal Substrate/Available Cover | 0-20 | 1. Epifaunal Substrate/Available Cover | 0-20 | |
| | 0-20 14 0-20 8 | 2. Pool Substrate Characterization 3. Pool Variability | 0-20 | | 2. Embeddedness 0-20 3. Velocity/ Depth Regime 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | |
| | 0-20 18 | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition 0-20 | | 4. Sediment Deposition | 0-20 | 4. Sediment Deposition | 0-20 | |
| | 0-20 0.4 10 | 5. Channel Flow Status | 0-20 0-1 | | 5. Channel Flow Status 0-20 | | 5. Channel Flow Status | 0-20 0.4 | 5. Channel Flow Status | 0-20 0.1 | |
| | 0-20 0-1 18 | 6. Channel Alteration | 0-20 0-1 | | 6. Channel Alteration 0-20 | | 6. Channel Alteration | 0-20 0-1 | 6. Channel Alteration | 0-20 0-1 | |
| | 0-20 18 | 7. Channel Sinuosity | 0-20 | | 7. Frequency of Riffles (or bends) 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | 7. Frequency of Riffles (or bends) | 0-20 | |
| | 0-20 18 | 8. Bank Stability (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) 0-20 | | 8. Bank Stability (LB & RB) | 0-20 | 8. Bank Stability (LB & RB) | 0-20 | |
| | 0.20 18 | 9. Vegetative Protection (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) 0-20 | | 9. Vegetative Protection (LB & RB) | 0-20 | 9. Vegetative Protection (LB & RB) | 0-20 | |
| | 0-20 18 | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 10. Riparian Vegetative Zone Width (LB & RB) 0.20 | | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | 10. Riparian Vegetative Zone Width (LB & RB) | | |
| | Suboptimal 155 | Total RBP Score | Poor 0 | | | Poor 0 | Total RBP Score | Poor 0 | Total RBP Score | Poor | 0 |
| Sub-Total | 0.775 | Sub-Total | 0 | | Sub-Total | 0 | Sub-Total | 0 | Sub-Total | | 0 |
| CHEMICAL INDICATOR (Applies to Intermittent an | nd Perennial Streams) | CHEMICAL INDICATOR (Applies to Intermittent a | and Perennial Streams) | | CHEMICAL INDICATOR (Applies to Intermittent and Pe | ennial Streams) | CHEMICAL INDICATOR (Applies to Intermitter | nt and Perennial Streams) | CHEMICAL INDICATOR (Applies to Intermit | tent and Perennial Stree | eams) |
| WVDEP Water Quality Indicators (General) | | WVDEP Water Quality Indicators (General) | | | WVDEP Water Quality Indicators (General) | | WVDEP Water Quality Indicators (General | 0 | WVDEP Water Quality Indicators (Gener | ral) | |
| Specific Conductivity | | Specific Conductivity | | | Specific Conductivity | | Specific Conductivity | | Specific Conductivity | | |
| <=99 - 90 points | 0-90 31.3 | | 0-90 | | 0-90 | | | 0-90 | | 0-90 | |
| nH | | nH | | | pH | | nH | | nH | | |
| | 0-80 0-1 7,99 | | 5-90 0-1 | | 5.97 | 0-1 | ··· | 5.90 0-1 | | 5-90 0-1 | |
| 6.0-8.0 = 80 points | 1.55 | | | | | | | | | | |
| DO | | DO | | | DO | | DO | | DO | | |
| >5.0 = 30 points | 10-30 8.17 | | 10-30 | | 10-3 | | | 10-30 | | 10-30 | |
| Sub-Total | 1 | Sub-Total | 0 | | Sub-Total | 0 | Sub-Total | 0 | Sub-Total | | 0 |
| BIOLOGICAL INDICATOR (Applies to Intermittent | and Perennial Streams) | BIOLOGICAL INDICATOR (Applies to Intermitter | nt and Perennial Streams) | | BIOLOGICAL INDICATOR (Applies to Intermittent an | d Perennial Streams) | BIOLOGICAL INDICATOR (Applies to Interm | nittent and Perennial Streams) | BIOLOGICAL INDICATOR (Applies to Inte | rmittent and Perennia | ial Streams) |
| WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | 1 1 | |
| Good | 0-100 0-1 70.79 | | 0-100 0-1 | | 0-10 | 0-1 | | 0-100 0-1 | | 0-100 0-1 | |
| G000 Sub-Total | 0.7079 | Sub-Total | 0 | | Sub-Total | 0 | Sub-Total | 0 | Sub-Total | | 0 |
| | | | | | | | | | | | |
| PART II - Index and Unit | Score | PART II - Index and U | Jnit Score | | PART II - Index and Unit S | core | PART II - Index and U | Init Score | PART II - Index and | I Unit Score | |
| Index | Linear Feet Unit Score | Index | Linear Feet Unit Score | | Index Line | ar Feet Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet | Unit Score |
| 0.828 | 74 61.2448667 | 0 | | | 0 | | | 1 1 | | 0 | 0 |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME | REASON FOR SURVEY |



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Local Watershed NPS Pollution Forest Commercial Field/Pasture Industrial Agricultural Other Residential Other Indicate the dominant type and record the dominant species present Herbaceous Trees Shrubs Grasses Dominant species present Herbaceous |
|--|--|
| INSTREAM FEATURES | Dominant species present |
| LARGE WOODY | LWDm ² |
| DEBRIS | Density of LWDm ² /km ² (LWD/ reach area) |
| AQUATIC | Indicate the dominant type and record the dominant species present |
| VEGETATION | Rooted emergent Rooted submergent Rooted floating Free floating Floating Algae Attached Algae Booted floating Free floating Free floating Dominant species present |
| WATER QUALITY (DS, US) | Temperature0 C Water Odors Normal/None Sewage Specific Conductance Petroleum Fishy Chemical Other Dissolved Oxygen Water Surface Oils Slick Sheen None Globs Flecks pH Turbidity (if not measured) Clear Slightly turbid Turbid Turbid Turbid Opaque Turbid |
| SEDIMENT/ | Odors |
| SUBSTRATE | Normal Sewage Petroleum Deposits Chemical Anaerobic None Sludge Sawdust Paper fiber Sand Other Other Epoking at stones which are not deeply embedded are the undersides black in color? How are the undersides black in color? |

| INC | ORGANIC SUBSTRATE (should add up to | | | ORGANIC SUBSTRATE COMPONEN (does not necessarily add up to 100% | | | | | | | | | |
|-------------------|--|------------------------------------|-------------------|--|-----------------------------------|--|--|--|--|--|--|--|--|
| Substrate Type | Diameter | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | | | | | | | | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | | | | | | | | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | | | | | | | | | |
| Cobble | 64-256 mm (2.5"-10") | | Muck-Mud | black, very fine organic | | | | | | | | | |
| Gravel | 2-64 mm (0.1"-2.5") | | | (FPOM) | | | | | | | | | |
| Sand | 0.06-2mm (gritty) | | Marl | grey, shell fragments | | | | | | | | | |
| Silt | 0.004-0.06 mm | | | | | | | | | | | | |
| Clay | < 0.004 mm (slick) | | | | | | | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME AM PM | REASON FOR SURVEY |

| | Habitat | | Condition | ı Category | |
|--|---|---|---|---|---|
| | Parameter | Optimal | Suboptimal | Marginal | Poor |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. |
| ted i | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). |
| uram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Condition | ı Category | |
|---|--|--|--|---|
| Parameter | Optimal | Suboptimal | Marginal | Poor |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE(RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME S-H | 113 (1) | LOCATION Webster County | , |
|----------------------|--|---|--|
| STATION # | RIVERMILE | STREAM CLASS Perennial | |
| LAT <u>38.612982</u> | LONG80.503647 | RIVER BASIN None | |
| STORET # | | AGENCY WVDEP | |
| INVESTIGATORS R | H VM MB | - | LOT NUMBER |
| FORM COMPLETED | RH | DATE 9/21/21 TIME 0737 | REASON FOR SURVEY Baseline Assessment |
| | | - | |
| HABITAT TYPES | Indicate the percentage of ✓Cobble 40 % Sn Submerged Macrophytes | ags% 🗍 Vegetated B | |
| SAMPLE COLLECTION | | lected? ☑ wading ☐ f ps/kicks taken in each habitat ty lags ☐ Vegetated B | rom bank from boat |
| GENERAL COMMENTS | | | C: 36.6 us/cm ph: 8.59 C: 31.3 us/cm ph: 7.99 |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)


| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | - | |

| | Count - | | signated as such in the Sample Methodolgy column on the Benthic ID forms (Family or Gen WVSCI Metrics and Scores 0R6 ID REIC2 |
|--|--|--------------------------------------|---|
| Bestifise ↓ Cambaridae Chironomidae Chironomidae Elmidae Goeridae Heptagenidae Hydropsychidae Leuctridae | 3 2 16 2 1 3 3 33 2 2 | 4 5 1 4 4 4 5 3 | Wetrics BSV BSV Benthic Density % 2 Dominant Taxa (Family) 58.33 37.3 66.45 # of grids Picked 100 Total # of grids % Chironomidae 19.05 1.7 82.35 Total IBI Individuals 84 % EPT (Family) 67.86 89.3 75.99 # of Organisms per Grid 0.84 |
| Oligochaeta Perildae Philopotamidae Psephenidae Tipulidae | 2 8 3 2 4 | 10 1 3 4 3 | HBI (Family) 4.48 2.61 74.75 # EPT Taxa (Family) 8 13 61.54 # Total Taxa (Family) 14 22 63.64 WVSCI Score w/ BSV 1996-2001 70.79 70.79 WVSCI Category Unimpaired-Good 70.70 |
| | | | WVSCI Thresholds Unimpaired = >60.61 to 58.00 Gray Zone = 60.61 to 58.00 Impaired = <60.61 |

WOLMAN PEBBLE COUNT FORM

| County: | Webster | Stream ID: | S-H113 (1) |
|--------------|----------------------|------------|------------|
| Stream Name: | UNT to Elk River (1) | | |
| HUC Code: | | Basin: | |
| Survey Date: | 9/8/2021 | | |
| Surveyors: | AC, VM, HK | Impact: | 34.4 m |
| Type: | Bankfull Channel | | |

| | | | LE COUNT | | | | |
|-------------|--------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | • • | 0 | 0.00 | 0.00 |
| | Very Fine | .062125 | SAND | * * | 0 | 0.00 | 0.00 |
| | Fine | .12525 | | • • | 0 | 0.00 | 0.00 |
| | Medium | .255 | | • • | 0 | 0.00 | 0.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | 6 | 6.00 | 6.00 |
| .0408 | Very Coarse | 1.0-2 | | • • | 6 | 6.00 | 12.00 |
| .0816 | Very Fine | 2 -4 | GRAVEL | * * | 9 | 9.00 | 21.00 |
| .1622 | Fine | 4 -5.7 | | * * | 3 | 3.00 | 24.00 |
| .2231 | Fine | 5.7 - 8 | | • • | 5 | 5.00 | 29.00 |
| .3144 | Medium | 8 -11.3 | | | 5 | 5.00 | 34.00 |
| .4463 | Medium | 11.3 - 16 | | • • | 4 | 4.00 | 38.00 |
| .6389 | Coarse | 16 -22.6 | | • • • | 4 | 4.00 | 42.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | • • | 3 | 3.00 | 45.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | • • | 5 | 5.00 | 50.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | • • • | 3 | 3.00 | 53.00 |
| 2.5 - 3.5 | Small | 64 - 90 | - COBBLE | • • • | 2 | 2.00 | 55.00 |
| 3.5 - 5.0 | Small | 90 - 128 | | | 4 | 4.00 | 59.00 |
| 5.0 - 7.1 | Large | 128 - 180 | | • • • | 2 | 2.00 | 61.00 |
| 7.1 - 10.1 | Large | 180 - 256 | | | 1 | 1.00 | 62.00 |
| 10.1 - 14.3 | Small | 256 - 362 | BOULDER | • • • | 30 | 30.00 | 92.00 |
| 14.3 - 20 | Small | 362 - 512 | | • • | 0 | 0.00 | 92.00 |
| 20 - 40 | Medium | 512 - 1024 | | • • | 8 | 8.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | | • • | 0 | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 | | • • | 0 | 0.00 | 100.00 |
| | Bedrock | | BDRK | • • | 0 | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |
| | Total Tally: | | | | | | |



Bankfull Channel Pebble Count, S-H113 (1), UNT to Elk River (1)

