Reach S-IJ57 (Pipeline ROW) Perennial Spread D Nicholas County, West Virginia

Data	Included
Photos	\checkmark
SWVM Form	\checkmark
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope <4%)
RBP Physical Characteristics Form	\checkmark
Water Quality Data	N/A – Low flow
RBP Habitat Form	\checkmark
RBP Benthic Form	\checkmark
Benthic Identification Sheet	N/A – Low flow
Wolman Pebble Count	\checkmark
Reference Reach Software Pebble Count Data	\checkmark
Longitudinal Profile and Cross Sections	\checkmark



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County

Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, JR/KP Lat: 38.352362 Long: -80.636401



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County

Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, JR/KP Lat: 38.352362 Long: -80.636401



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County

Photo Type: US View at Center Location, Orientation, Photographer Initials: Center ROW, Upstream View, JR/KP Lat: 38.352362 Long: -80.636401



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County

Photo Type: DS View at Center Location, Orientation, Photographer Initials: Center ROW, Downstream View, JR/KP Lat: 38.352362 Long: -80.636401



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County

Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, JR/KP Lat: 38.352362 Long: -80.636401



Spread D Stream S-IJ57 (Pipeline ROW) Nicholas County


Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, JR/KP Lat: 38.352362 Long: -80.636401

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

	USACE FILE NO./ Project Name: (v2.1, Sept 2015)	Mountain V	/alley Pipeline	IMPACT COORDINATES: (in Decimal Degrees)	Lat.	38.352362	Lon.	-80.636401	WEATHER:	Sunny	DATE:	08/31/21
			S-D	57							Comments:	
<form> Autom Lum Control Interm <th< th=""><th>STREAM IMPACT LENGTH: 82</th><th></th><th>RESTORATION (Levels I-III)</th><th></th><th>Lat.</th><th></th><th>Lon.</th><th></th><th>PRECIPITATION PAST 48 HRS:</th><th></th><th>Mitigation Length:</th><th></th></th<></form>	STREAM IMPACT LENGTH: 82		RESTORATION (Levels I-III)		Lat.		Lon.		PRECIPITATION PAST 48 HRS:		Mitigation Length:	
	Column No. 1- Impact Existing Condition (De	Debit)	Column No. 2- Mitigation Existing Cor	dition - Baseline (Credit)				'ears			Column No. 5- Mitigation Project	ed at Maturity (Credit)
	Stream Classification: Per	rennial	Stream Classification:			Stream Classification:		0	Stream Classification:	0	Stream Classification:	0
	Percent Stream Channel Slope	2.9	Percent Stream Channel Slop	Ð		Percent Stream Channel	I Slope	0	Percent Stream Channel Sic	ope 0	Percent Stream Channel S	lope 0
	HGM Score (attach data forms):		HGM Score (attach da	ta forms):		HGM Score (atta	ach data forms):		HGM Score (attach da	ita forms):	HGM Score (attach o	ata forms):
		Average		Average				Average		Average		Average
		U		0		Habitat		-			Habitat	
	PART I - Physical, Chemical and Biological Indi	dicators	PART I - Physical, Chemical and I	Biological Indicators		PART I - Physical, Chemica	al and Biological Ind	icators	PART I - Physical, Chemical and B	Biological Indicators	PART I - Physical, Chemical and	Biological Indicators
	Point Scale Range	pe Site Score		ointa Scale Range Sille Score			Points Scale Range	Site Score		Points Scale Range Site Score		Points Scale Range Site Score
	PHYSICAL INDICATOR (Applies to all streams classifications)		PHYSICAL INDICATOR (Applies to all streams cla	ssifications)		PHYSICAL INDICATOR (Applies to all stres	ams classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)	PHYSICAL INDICATOR (Applies to all stream	classifications)
2. Dots built of the second	USEPA RBP (High Gradient Data Sheet)		USEPA RBP (Low Gradient Data Sheet)			USEPA RBP (High Gradient Data Sheet			USEPA RBP (High Gradient Data Sheet)		USEPA RBP (High Gradient Data Sheet)	
3. Vector Lipsch Reginer 6. No 7. Vector J. Reginer 6. No		10										
S. Owned Needing S. Owned Needing <td< td=""><td>3. Velocity/ Depth Regime 0-20</td><td>8</td><td>3. Pool Variability</td><td></td><td></td><td></td><td></td><td></td><td>3. Velocity/ Depth Regime</td><td></td><td></td><td></td></td<>	3. Velocity/ Depth Regime 0-20	8	3. Pool Variability						3. Velocity/ Depth Regime			
0. Overed Aluretion 0.00 1 0.00 1 0.00 </td <td></td>												
2. Program of dRBs (a for band) as 3. Band Stability (Ad SRB) as 0. Subjective (MBL RBB) as as 0. Subjective (MBL RBB) as 0. Subjective (MBL RBB) as as as 0. Subjective (MBL RBB) as as <td></td>												
 		-										
10. Registed Scaw With (10.8 RB) 0.0												
Total RBP Socie Total RBP Socie <thtotal rbp="" socie<="" th=""> T</thtotal>		10										
Sub-Total 0.52 Sub-Total 0.5 Sub-Total 0		104						0				
							Pool	0				
$ \frac{\text{Specific Conductivity}}{100,090,850016} \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ $	CHEMICAL INDICATOR (Applies to Intermittent and Perennial S	Streams)	CHEMICAL INDICATOR (Applies to Intermittent an	d Perennial Streams)		CHEMICAL INDICATOR (Applies to Interm	ittent and Perennial Stre	zams)	CHEMICAL INDICATOR (Applies to Intermittent	t and Perennial Streams)	CHEMICAL INDICATOR (Applies to Intermitte	nt and Perennial Streams)
indication index (WSC) index ind	WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (General)				eral)		WVDEP Water Quality Indicators (General)		WVDEP Water Quality Indicators (General)
in 00100-06 sponds in 0010-06 sponds in 00			Specific Conductivity			Specific Conductivity	-		Specific Conductivity		Specific Conductivity	
mode	100-199 - 85 points	133		0-90			0-90			0-90		0-90
mode	pH		pH			pH			pH		pH	
DO DO <th< td=""><td>0-80 0-1</td><td>7.7</td><td></td><td>5-90 0-1</td><td></td><td></td><td>5-90 0-1</td><td></td><td></td><td>5-90 0-1</td><td></td><td>5-90 0-1</td></th<>	0-80 0-1	7.7		5-90 0-1			5-90 0-1			5-90 0-1		5-90 0-1
× 0.0 = 30 portis 0 0.0 = 30 portis 0.0 = 30 portis 0 <th< td=""><td></td><td></td><td>00</td><td></td><td></td><td>DO</td><td></td><td></td><td>DO</td><td></td><td>DO.</td><td></td></th<>			00			DO			DO		DO.	
× 0.0 = 30 portis 0 0.0 = 30 portis 0.0 = 30 portis 0 <th< td=""><td>10.20</td><td>7.9</td><td>50</td><td>10-20</td><td></td><td>50</td><td>10.30</td><td></td><td></td><td>10.30</td><td>50</td><td>10.30</td></th<>	10.20	7.9	50	10-20		50	10.30			10.30	50	10.30
BIOLOGICAL INDICATOR (Applies to Intermittent and Perrure) BIOLOGICAL INDICA	>5.0 = 30 points		0.1.7.1.1						0.1.7.1.1		0.1.7.1	
W Stream Condition Index (WVSC) W Stream Condition Ind				U O			termittent and Berenni	U U		U		ittent and Perennial Streeme)
0 0.00 0.10 0.00 0.10 0.00 0.10 <t< td=""><td></td><td>al Streams)</td><td></td><td>and Perennial Streams)</td><td></td><td></td><td>termittent and Ferenin</td><td>an arreants)</td><td></td><td>intent and Perennial Sceans)</td><td></td><td>intent and Perennal Streams)</td></t<>		al Streams)		and Perennial Streams)			termittent and Ferenin	an arreants)		intent and Perennial Sceans)		intent and Perennal Streams)
0 0 0 Sub-Total 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 <td></td> <td>1</td> <td></td> <td>0.100 0.1</td> <td></td> <td>WV Stream Condition Index (WVSCI)</td> <td>0.100 0.1</td> <td></td> <td></td> <td>0.100 0.1</td> <td>Wy Stream condition muex (WySci)</td> <td>0.100 0.1</td>		1		0.100 0.1		WV Stream Condition Index (WVSCI)	0.100 0.1			0.100 0.1	Wy Stream condition muex (WySci)	0.100 0.1
PART II - Index and Unit Score PART	0	-										
Index Linear Feet Unit Score Index	Sub-Total	0	Sub-Total	0	J	Sub-Total		0	Sub-Total	0	Sub-Total	0
	PART II - Index and Unit Score		PART II - Index and U	nit Score		PART II - Index	and Unit Score		PART II - Index and U	nit Score	PART II - Index and	Jnit Score
0.748 82 61.295 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Index Linear Feet	t Unit Score	Index	Linear Feet Unit Score		Index	Linear Feet	Unit Score	Index	Linear Feet Unit Score	Index	Linear Feet Unit Score
	0.748 82	61.295	0	0 0	1	0	0	0	0	0 0	0	0 0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME	LOCATION					
STATION # RIVERMILE	STREAM CLASS					
LAT LONG	RIVER BASIN	RIVER BASIN				
STORET #	AGENCY					
INVESTIGATORS						
FORM COMPLETED BY	DATE TIME	REASON FOR SURVEY				



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer)	Predominant Surrounding Landuse Local Watershed NPS Pollution Forest Commercial Field/Pasture Industrial Agricultural Other Residential Other Indicate the dominant type and record the dominant species present Herbaceous Trees Shrubs Grasses Dominant species present Herbaceous
INSTREAM FEATURES	Dominant species present
LARGE WOODY	LWDm ²
DEBRIS	Density of LWDm ² /km ² (LWD/ reach area)
AQUATIC	Indicate the dominant type and record the dominant species present
VEGETATION	Rooted emergent Rooted submergent Rooted floating Free floating Floating Algae Attached Algae Booted floating Free floating Free floating Dominant species present
WATER QUALITY (DS, US)	Temperature0 C Water Odors Normal/None Sewage Specific Conductance Petroleum Fishy Chemical Other Dissolved Oxygen Water Surface Oils Slick Sheen None Globs Flecks pH Turbidity (if not measured) Clear Slightly turbid Turbid Turbid Turbid Opaque Turbid
SEDIMENT/	Odors
SUBSTRATE	Normal Sewage Petroleum Deposits Chemical Anaerobic None Sludge Sawdust Paper fiber Sand Other Other Epoking at stones which are not deeply embedded are the undersides black in color? How are the undersides black in color?

INC	ORGANIC SUBSTRATE (should add up to			ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)						
Substrate Type	Diameter	meter % Composition in Sampling Reach		Characteristic	% Composition in Sampling Area					
Bedrock			Detritus	sticks, wood, coarse plant						
Boulder	> 256 mm (10")			materials (CPOM)						
Cobble	64-256 mm (2.5"-10")		Muck-Mud	black, very fine organic						
Gravel	2-64 mm (0.1"-2.5")			(FPOM)						
Sand	0.06-2mm (gritty)		Marl	grey, shell fragments						
Silt	0.004-0.06 mm									
Clay	< 0.004 mm (slick)									

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION				
STATION # RIVERMILE	STREAM CLASS				
LAT LONG	RIVER BASIN				
STORET #	AGENCY				
INVESTIGATORS					
FORM COMPLETED BY	DATE TIME AM PM	REASON FOR SURVEY			

	Habitat		Condition	ı Category			
	Parameter	Optimal	Suboptimal	Marginal	Poor		
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.		
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.		
ted i	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).		
uram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
P	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.		
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		
	5. Channel Flow Status						
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0		

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

Habitat		Condition	ı Category	
Parameter	Optimal	Suboptimal	Marginal	Poor
6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.
SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
 SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) 	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0
9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0
10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.
SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0
SCORE(RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME		LOCATION						
STATION #	_ RIVERMILE	STREAM CLASS						
LAT	LONG	RIVER BASIN						
STORET #		AGENCY						
INVESTIGATORS			LOT NUMBER					
FORM COMPLETED	BY	DATE TIME	REASON FOR SURVEY					
HABITAT TYPES	Cobble% Sn	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%						
SAMPLE COLLECTION	Indicate the number of jab	lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B	anks Sand					
GENERAL COMMENTS								

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

WOLMAN PEBBLE COUNT FORM

County:NicholasStream Name:UNT to Barn RunHUC Code:Survey Date:8/31/2021Surveyors:JR KPType:Bankfull Channel


S-IJ57

Reach:: 23m

Stream ID:

Basin:

			LE COUNT			1	
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	• •	42	42.00	42.00
	Very Fine	.062125		•	2	2.00	44.00
	Fine	.12525		•	6	6.00	50.00
	Medium	.255	S A N D	▲ ▼	13	13.00	63.00
	Coarse	.50-1.0		•	4	4.00	67.00
.0408	Very Coarse	1.0-2		• •	2	2.00	69.00
.0816	Very Fine	2 -4		▲ ▼	1	1.00	70.00
.1622	Fine	4 -5.7]	• •	2	2.00	72.00
.2231	Fine	5.7 - 8	1	• •	2	2.00	74.00
.3144	Medium	8 -11.3		• •	2	2.00	76.00
.4463	Medium	11.3 - 16	GRAVEL	• •	4	4.00	80.00
.6389	Coarse	16 -22.6		▲ ▼	2	2.00	82.00
.89 - 1.26	Coarse	22.6 - 32	1	• •	5	5.00	87.00
1.26 - 1.77	Vry Coarse	32 - 45		▲ ▼	4	4.00	91.00
1.77 -2.5	Vry Coarse	45 - 64	1	• •	4	4.00	95.00
2.5 - 3.5	Small	64 - 90		• •	3	3.00	98.00
3.5 - 5.0	Small	90 - 128	CODDLE	▲ ▼	2	2.00	100.00
5.0 - 7.1	Large	128 - 180	COBBLE	▲ ▼	0	0.00	100.00
7.1 - 10.1	Large	180 - 256	1	▲ ▼	0	0.00	100.00
10.1 - 14.3	Small	256 - 362		• •	0	0.00	100.00
14.3 - 20	Small	362 - 512	1	▲ ▼	0	0.00	100.00
20 - 40	Medium	512 - 1024	BOULDER	▲ ▼	0	0.00	100.00
40 - 80	Large	1024 -2048]	▲ ▼	0	0.00	100.00
80 - 160	Vry Large	2048 -4096]	▲ ▼	0	0.00	100.00
	Bedrock		BDRK	▲ ▼	0	0.00	100.00
				Totals:	100		




cumulative %


-


■# of particles

 Size (mm) 💦 📩		Size Distr	Size Distribution		Туре		
D16	0.062	mean	1.3	silt/clay	42%		
D35	0.062	dispersion	54.0	sand	27%		
D50	0.25	skewness	0.44	gravel	26%		
D65	0.71			cobble	5%		
D84	26			boulder	0%		
D95	64						



AS-BUILT TABLE: S-IJ57 CROSS SECTION A										
	PI	AS-BUILT								
PT. LOC.	NORTHING	EASTING	ELEV	VERT. DIFF.	HORZ. DIFF.					
TS-L	13927058.5300	1744621.6190'	2213.805'							
BS-L	13927059.9200	1744622.32501	2213.385'							
THW	13927060.5600	1744622.6740'	2213.222'							
BS-R	13927061.0700	1744622.7940'	2213.283'							
TS-R	13927061.6600	1744623.23001	2213.739'							





SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON AUGUST 31, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.

3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.

- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.

