Baseline Assessment – Stream Attributes

Reach S-K82 (Pipeline ROW) Perennial Spread B Harrison County, West Virginia

| Data | Included |
|--|--|
| Photos | ✓ |
| SWVM Form | ✓ |
| FCI Calculator and HGM Form | NA- Perennial Stream (not shadeable slope <4%) |
| RBP Physical Characteristics Form | ✓ |
| Water Quality Data | ✓ |
| RBP Habitat Form | ✓ |
| RBP Benthic Form | ✓ |
| Benthic Identification Sheet | NA- lack of habitat |
| Wolman Pebble Count | ✓ |
| Reference Reach Software Pebble Count Data | ✓ |
| Longitudinal Profile and Cross Sections | √ |



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181



Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, DP/VM/HK
Lat: 39.167753 Long: -80.578181



Photo Type: US View at Center
Location, Orientation, Photographer Initials: Center ROW, Upstream View, DP/VM/HK
Lat: 39.167753 Long: -80.578181



Photo Type: DS View at Center Location, Orientation, Photographer Initials: Center ROW, Downstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181



Photo Type: Pool, DS View Location, Orientation, Photographer Initials: Upstream of Pool, Downstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181



Photo Type: Pool, US View Location, Orientation, Photographer Initials: Downstream of Pool, Upstream View, DP/VM/HK Lat: 39.167753 Long: -80.578181

| USACE FILE NO./ Project Name: Mount (v2.1, Sept 2015) | nin Valley Pipeline IMPACT COORDINA (in Decimal Degree | | Lat. | t. 39.167753 Lon80.578181 | | WEATHER: | S | teady Rain | DATE: | 8/31/2021 |
|--|--|--|------|--|--------------------------------------|--|---------------------|--------------|--|-------------------------------|
| IMPACT STREAM/SITE ID AND SITE DESCRIPTION: (watershed size (acreage), unaltered or impairments) | S- | K82 | | MITIGATION STREAM CLASS./SITE ID AI (watershed size (acreage), unaltered o | ID SITE DESCRIPTION: impairments) | | | | Comments: | |
| STREAM IMPACT LENGTH: 110 FORM OF MITIGATION: | RESTORATION (Levels I-III) | MIT COORDINATES: (in Decimal Degrees) | Lat. | Lon. | | PRECIPITATION PAST 48 HRS: | | | Mitigation Length: | |
| Column No. 1- Impact Existing Condition (Debit) | Column No. 2- Mitigation Existing C | ondition - Baseline (Credit) | | Column No. 3- Mitigation Projected at F Post Completion (Credit) | ive Years | Column No. 4- Mitigation Proj Post Completion (| | ars | Column No. 5- Mitigation Projected | at Maturity (Credit) |
| Stream Classification: Perennial | Stream Classification: | | | Stream Classification: | 0 | Stream Classification: | (| D | Stream Classification: | 0 |
| Percent Stream Channel Slope 0.2 | Percent Stream Channel Slo | оре | | Percent Stream Channel Slope | 0 | Percent Stream Channel SI | оре | 0 | Percent Stream Channel Slo | pe 0 |
| HGM Score (attach data forms): | HGM Score (attach | data forms): | | HGM Score (attach data form | 5): | HGM Score (attach da | ata forms): | | HGM Score (attach dat | a forms): |
| Average Hydrology Blogeochemical Cycling Dahottat Habitat Physical, Chemical and Biological Indicators | Hydrology Biogeochemical Cycling Habitat PART I - Physical, Chemical an | Average 0 d Biological Indicators | | Hydrology Biogeochemical Cycling Habitat PART I - Physical, Chemical and Biologic. | Average 0 | Hydrology Biogeochemical Cycling Habitat PART I - Physical, Chemical and | Biological Indic | Average 0 | Hydrology Biogeochemical Cycling Habitat PART I - Physical, Chemical and B | Average 0 |
| Prints Scine Range Site Score | | Points Scale Range Site Score | | Point Scale | Range Site Score | • | Points Scale Range | Site Score | | Points Scale Range Site Score |
| PHYSICAL INDICATOR (Applies to all streams classifications) | PHYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDICATOR (Applies to all streams classification | 3) | PHYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams of | assifications) |
| USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover 0-20 16 | USEPA RBP (Low Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover | | | USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover 0-20 | | USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover | | | USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover | |
| 1. Epifarum Substrate/Available Cover | 2. Pool Substates Chrostorials of Control State of Contro | 0-90 5-90 0-1 | | 1. Epiflurian Substate Mealistable Cover | 0-1 | 2. Embeldischenss 3. Velocity Orgin Regime 4. Sediment Deposition 5. Channel Flow Status 6. Channel Alteration 7. Frequency of Rittles (or bends) 8. Bank Statuliny (18.8 A RB) 10. Reparter Verbeding (18.8 A RB) 10. Part Verbeding (18.8 A RB) 10. Reparter Verbeding (18.8 A RB) 10. Reparter Verbeding (18.8 A RB) 10. Reparter Verbeding (18.8 A RB) 10. Sediment Verbed | 0-90 5-90 0-1 | 0 | 2. Embeddedories 3. Velocity Depth Regime 4. Sediment Deposition 6. Channel Flow Status 6. Channel Flow Status 6. Channel Alteration 7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB) 9. Vegotative Protection (LB & RB) 10. Ripartan Vegetative Zone Width (LB & RB) 17. Ripartan Vegetative Zone Width (LB & RB) 17. Ripartan Vegetative Zone Width (LB & RB) 18. WDDEP Water Quality Indicators (General) 3. Specific Conductivity Ph DO Sub-Total BIOLOGICAL INDICATOR (Applies to Intermit | 5-90 0-1 |
| WV Stream Condition Index (WVSCI) | WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | |
| 0 0-100 0-1 Sub-Total 0 | Sub-Total | 0-100 0-1 | | 0-100 Sub-Total | 0-1 | Sub-Total | 0-100 0-1 | 0 | Sub-Total | 0-100 0-1 |
| PART II - Index and Unit Score | PART II - Index and | Unit Score | | PART II - Index and Unit Score | | PART II - Index and U | Init Score | | PART II - Index and Un | it Score |
| Index Linear Feet Unit Score | Index | Linear Feet Unit Score | | Index Linear I | eet Unit Score | Index | Linear Feet | Unit Score | Index | Linear Feet Unit Score |
| 0.788 110 86.625 | 0 | 0 0 | | 0 0 | 0 | 0 | 0 | 0 | 0 | 0 0 |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | | LOCATION | | | |
|-------------------|-----------|--------------|-------------------|--|--|
| STATION# | RIVERMILE | STREAM CLASS | | | |
| LAT | LONG | RIVER BASIN | | | |
| STORET# | | AGENCY | | | |
| INVESTIGATORS | | | | | |
| FORM COMPLETED BY | | DATE | REASON FOR SURVEY | | |

| | | | | yy |
|----------------------------|---|--|----------------|---|
| WEATHER CONDITIONS | | storm (heavy rain) rain (steady rain) nowers (intermittent) %cloud cover clear/sunny | Past 24 hours | Has there been a heavy rain in the last 7 days? Yes No Air Temperature0 C Other |
| SITE LOCATION/MAP | Draw a map of t | Lob | k 92 | oming In Going Away |
| STREAM CHARACTERIZATION | Stream Subsyste Perennial Stream Origin Glacial Non-glacial m Swamp and bo | Spring-fee | l f origins | Stream Type Coldwater Warmwater Catchment Areakm² |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERS FEATURI | | Fores Field Agric | Pasture Industria | rcial | No evidence Sor Obvious sources Local Watershed Erosi None Moderate | ne potential sources |
|--|----------------------------------|--|--|--|---|-----------------------------------|
| RIPARIA VEGETA (18 meter | ΓION | Trees | e the dominant type and Sl ant species present | hrubs | Grasses He | brbaceous |
| INSTREA FEATURI | | Estimat Samplin Area in Estimat | red Stream Depthm | m m² km² m | Canopy Cover Partly open Part High Water Mark Proportion of Reach R Morphology Types Riffle Pool 9 Channelized Yes Dam Present Yes | epresented by Stream Run% No |
| LARGE V DEBRIS | VOODY | | m² of LWDm | 1 ² /km ² (LWD / 1 | reach area) | |
| AQUATIO VEGETA | | Domina | | | minant species present nt Rooted floating | Ü |
| WATER ((DS, US) | QUALITY | Specific Dissolve pH Turbidi | rature0 C Conductance ed Oxygen ty trument Used | | Water Odors Normal/None Sewage Petroleum Fishy Water Surface Oils Slick Sheen None Other Turbidity (if not measu Clear ☐ Slightly tu Opaque Stained | Chemical Other Globs Flecks |
| SEDIMENT/ SUBSTRATE Odors Normal Sewage Chemical Anaerobic Other Oils Absent Slight Modera | | | | | are the undersides blac | th are not deeply embedded, |
| INC | ORGANIC SUBS (should a | | COMPONENTS 00%) | | ORGANIC SUBSTRATE C (does not necessarily add | |
| Substrate Type | Diamet | er | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area |
| Bedrock | | | | Detritus | sticks, wood, coarse plant materials (CPOM) | |
| Boulder Cobble | > 256 mm (10") 64-256 mm (2.5 | | | Muck-Mud | black, very fine organic | |
| Gravel | 2-64 mm (0.1"-2 | | | IVIUCK-IVIUU | (FPOM) | |

Sand

Silt

Clay

0.06-2mm (gritty)

< 0.004 mm (slick)

0.004-0.06 mm

grey, shell fragments

Marl

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET# | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE AM PM | REASON FOR SURVEY |

| | Habitat | | Condition | ı Category | | | | |
|--|---|---|---|---|---|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | | |
| ted in | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/depth regime (usually slow-deep). | | | |
| ıram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| Pa | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| | Habitat | Condition Category | | | | | | | | |
|--|--|--|--|--|---|--|--|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | | | |
| | 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | | |
| oling reach | 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | | | |
| samp | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | | |
| Parameters to be evaluated broader than sampling reach | 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion | Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | | | |
| e eva | SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |
| to be | SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |
| Parameters | 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | | | |
| | SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |
| | SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |
| | 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | | | | |
| | SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |
| ĺ | SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | |

| Total | Caama | |
|--------|-------|--|
| i otai | Score | |

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME | | LOCATION | | | | |
|---|-------------|-----------------------------|------------|--|--|--|
| STATION # | _ RIVERMILE | STREAM CLASS | | | | |
| LAT | LONG | RIVER BASIN | | | | |
| STORET# | | AGENCY | | | | |
| INVESTIGATORS | | | LOT NUMBER | | | |
| FORM COMPLETED BY | | DATE REASON FOR SURVEY TIME | | | | |
| HABITAT TYPES Indicate the percentage of each habitat type present Cobbbe % Space % Vacceted Ropks % Space % | | | | | | |

| HABITAT TYPES | Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()% |
|----------------------|--|
| SAMPLE COLLECTION | Gear used D-frame kick-net Other |
| | How were the samples collected? wading from bank from boat |
| | Indicate the number of jabs/kicks taken in each habitat type. Cobble Snags Vegetated Banks Sand Submerged Macrophytes Other () |
| GENERAL COMMENTS | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3 = Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

County: Harrison Stream ID: S-K82

Stream Name: UNT to Kincheloe Creek


HUC Code: 05020002 Basin: West Fork


Survey Date: 8/31/2021


Surveyors: DP, VM, HK Impact: 44m


Type: Bankfull Channel


| | | | LE COUNT | | | • | 1 |
|-------------|--------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | A | 5 | 5.00 | 5.00 |
| | Very Fine | .062125 | | ▲ | 0 | 0.00 | 5.00 |
| | Fine | .12525 | 1 | A | 0 | 0.00 | 5.00 |
| | Medium | .255 | SAND | A | 3 | 3.00 | 8.00 |
| | Coarse | .50-1.0 | 1 | ▲ | 8 | 8.00 | 16.00 |
| .0408 | Very Coarse | 1.0-2 | 1 | A | 0 | 0.00 | 16.00 |
| .0816 | Very Fine | 2 -4 | | A | 0 | 0.00 | 16.00 |
| .1622 | Fine | 4 -5.7 | 1 | A | 0 | 0.00 | 16.00 |
| .2231 | Fine | 5.7 - 8 | 1 | ^ | 0 | 0.00 | 16.00 |
| .3144 | Medium | 8 -11.3 | 1 | ^ | 0 | 0.00 | 16.00 |
| .4463 | Medium | 11.3 - 16 | GRAVEL | A | 0 | 0.00 | 16.00 |
| .6389 | Coarse | 16 -22.6 | 1 | A | 0 | 0.00 | 16.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | 1 | A | 0 | 0.00 | 16.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | 1 | A | 2 | 2.00 | 18.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | 1 | A | 14 | 14.00 | 32.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | A | 26 | 26.00 | 58.00 |
| 3.5 - 5.0 | Small | 90 - 128 | | ^ | 29 | 29.00 | 87.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | ^ | 13 | 13.00 | 100.00 |
| 7.1 - 10.1 | Large | 180 - 256 | 1 | ^ | 0 | 0.00 | 100.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | A | 0 | 0.00 | 100.00 |
| 14.3 - 20 | Small | 362 - 512 | 1 | A | 0 | 0.00 | 100.00 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | ^ | 0 | 0.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | 1 | A | 0 | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 | 1 | ^ | 0 | 0.00 | 100.00 |
| | Bedrock | | BDRK | ▲ | 0 | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |
| | Total Tally: | | | | | | |











SURVEY NOTES:

LEGEND


STUDY AREA (EASEMENT)

EXISTING SURVEY-LOCATED THALWEG


EXISTING SURVEYED GROUND SHOT ELEVATION

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON AUGUST 31, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.

S-K82 BASELINE CROSS-SECTION A POOL



S-K82 BASELINE CROSS-SECTION B RIFFLE



CROSS SECTION LEGEND

— EXISTING GRADE

CROSS SECTION

H: 1"=10'
SCALE: V: 1"-5'

NOTE: ALL SECTIONS VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

PRE-CROSSING PHOTOS



PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS



PHOTO TAKEN LOOKING UPSTREAM FROM DOWNSTREAM IMPACT LIMITS

POST-CROSSING PHOTOS

PENDING CROSSING

PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS

PENDING CROSSING

PHOTO TAKEN LOOKING UPSTREAM FROM

PRE-CROSSING

DOWNSTREAM IMPACT LIMITS

HE PROFILE AND CROSS—S BASELINE SURVE CROSSING S—K82 — UNNAM KINCHELOE CREEK (MP

CAD File No.

1

Drawing No