Reach S-VV13b (Temporary Access Road) Perennial Spread B Lewis County, West Virginia

| Data | Included |
|--|--|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | N/A – Perennial stream (not shadeable, slope <4%) |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | \checkmark |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | N/A – lack of habitat |
| Wolman Pebble Count | \checkmark |
| Reference Reach Software Pebble Count Data | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |



Photo Type: DS, US View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Upstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: DS, DS View Location, Orientation, Photographer Initials: Downstream Edge of ROW, Downstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: US View at Center Location, Orientation, Photographer Initials: Center ROW, Upstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: DS View at Center Location, Orientation, Photographer Initials: ROW Center, Downstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: US, US View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Upstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: US, DS View Location, Orientation, Photographer Initials: Upstream Edge of ROW, Downstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: Riffle, DS View Location, Orientation, Photographer Initials: Upstream of Riffle, Downstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: Riffle, US View Location, Orientation, Photographer Initials: Downstream of Riffle, Upstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: Pool, DS View Location, Orientation, Photographer Initials: Upstream of Pool, Downstream View, MAG/DPF Lat: 38.898431 Long: -80.56825



Photo Type: Pool, US View Location, Orientation, Photographer Initials: Downstream of Pool, Upstream View, MAG/DPF Lat: 38.898431 Long: -80.56825

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | | Mountain \ | Valley Pipeline | IMPACT COORDINAT (in Decimal Degrees | | 38.898431 | Lon. | -80.56825 | WEATHER: | 15% cloud cover | DATE: | 9/13/2021 |
|---|-------------------------------|------------------|---|---|---|---|---|---------------|---|----------------------------------|---|------------------------------|
| IMPACT STREAM/SITE ID (watershed size (acreage), | | : | S-VV13b Tempo | arary Access Road | | MITIGATION STREAM CL (watershed size (| ASS./SITE ID AND S (acreage), unaltered or imp | | | | Comments: | |
| STREAM IMPACT LENGTH: | | RM OF BATION: | RESTORATION (Levels I-III) | MIT COORDINATES (in Decimal Degrees) | | | Lon. | | PRECIPITATION PAST 48 HRS: | | Mitigation Length: | |
| Column No. 1- Impact Existing | g Condition (Debit) | | Column No. 2- Mitigation Existing C | Condition - Baseline (Credit) | | Column No. 3- Mitigati Post Com | tion Projected at Five pletion (Credit) | Years | Column No. 4- Mitigation P Post Completion | | Column No. 5- Mitigation Project | ted at Maturity (Credit) |
| Stream Classification: | Perennial | | Stream Classification: | | | Stream Classification: | | 0 | Stream Classification: | 0 | Stream Classification: | 0 |
| Percent Stream Channel SI | ope 2.4 | | Percent Stream Channel SI | оре | | Percent Stream Chan | nel Slope | 0 | Percent Stream Channel | Slope 0 | Percent Stream Channel S | Slope 0 |
| HGM Score (attach d | ata forms): | | HGM Score (attach | data forms): | | HGM Score (a | attach data forms): | | HGM Score (attacl | n data forms): | HGM Score (attach o | lata forms): |
| | Average | | | Average | | | | Average | | Average | | Averag |
| łydrology Biogeochemical Cycling łabitat | 0 | | Hydrology Biogeochemical Cycling Habitat | 0 | | Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling Habitat | 0 | Hydrology Biogeochemical Cycling Habitat | 0 |
| PART I - Physical, Chemical and | Biological Indicators | | PART I - Physical, Chemical an | d Biological Indicators | | Habitat PART I - Physical, Chemi | ical and Biological In | dicators | PART I - Physical, Chemical a | nd Biological Indicators | PART I - Physical, Chemical and | Biological Indicators |
| | Points Scale Range Site Score | | | Points Scale Range Site Score | | | Points Scale Range | Site Score | | Points Scale Range Site Score | | Painta Scale Range Site Scor |
| HYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDICATOR (Applies to all s | streams classifications) | | PHYSICAL INDICATOR (Applies to all stre | ams classifications) | PHYSICAL INDICATOR (Applies to all stream | s classifications) |
| EPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Low Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sh | | | USEPA RBP (High Gradient Data Shee |) | USEPA RBP (High Gradient Data Sheet) | |
| Epifaunal Substrate/Available Cover | 0-20 5 | _ | 1. Epifaunal Substrate/Available Cover | 0-20 | | 1. Epifaunal Substrate/Available Cover | | | 1. Epifaunal Substrate/Available Cover | 0-20 | 1. Epifaunal Substrate/Available Cover | 0-20 |
| Embeddedness Velocity/ Depth Regime | 0-20 17 | _ | 2. Pool Substrate Characterization 3. Pool Variability | 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 | 2. Embeddedness 3. Velocity/ Depth Regime | 0-20 |
| . Sediment Deposition | 0-20 16 | | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition | 0-20 | | 4. Sediment Deposition | 0-20 | 4. Sediment Deposition | 0-20 |
| Channel Flow Status | 0-20 0.4 8 | | 5. Channel Flow Status | 0-20 | | 5. Channel Flow Status | 0-20 | | 5. Channel Flow Status | 0-20 | 5. Channel Flow Status | 0-20 |
| Channel Alteration | 0-20 0-1 17 | | 6. Channel Alteration | 0-20 0-1 | | 6. Channel Alteration | 0-20 0-1 | | 6. Channel Alteration | 0-20 0-1 | 6. Channel Alteration | 0-20 0-1 |
| Frequency of Riffles (or bends) | 0-20 9 | | 7. Channel Sinuosity | 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | 7. Frequency of Riffles (or bends) | 0-20 |
| Bank Stability (LB & RB) | | | | | | | | | | | 8. Bank Stability (LB & RB) | |
| | 0-20 17 | | 8. Bank Stability (LB & RB) 9. Vegetative Protection (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) 9. Vegetative Protection (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) 9. Vegetative Protection (LB & RB) | 0-20 | 9. Vegetative Protection (LB & RB) | 0-20 |
| Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | 0-20 18 | | Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | 0-20 | | Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & I | | | Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | | Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | 0-20 |
| D. Riparian Vegetative Zone Width (LB & RB) otal RBP Score | | | Total RBP Score | 0-20 Poor 0 | | Total RBP Score | RB) 0-20 Poor | | Total RBP Score | Poor 0 | Total RBP Score | 0-20 Poor (|
| ub-Total | Suboptimal 138 0.69 | - | Sub-Total | Poor | | Sub-Total | Poor | 0 | Sub-Total | Poor | Sub-Total | Poor 0 |
| HEMICAL INDICATOR (Applies to Intermitten | | _ | CHEMICAL INDICATOR (Applies to Intermittent | t and Perennial Streams) | | CHEMICAL INDICATOR (Applies to Inte | ermittent and Perennial St | reams) | CHEMICAL INDICATOR (Applies to Interm | ittent and Perennial Streams) | CHEMICAL INDICATOR (Applies to Intermitte | |
| VDEP Water Quality Indicators (General | 0 | | WVDEP Water Quality Indicators (General) |) | | WVDEP Water Quality Indicators (Ge | eneral) | | WVDEP Water Quality Indicators (Gen | eral) | WVDEP Water Quality Indicators (General | 1) |
| pecific Conductivity | | | Specific Conductivity | | | Specific Conductivity | | | Specific Conductivity | | Specific Conductivity | |
| <=99 - 90 points | 0-90 87.8 | | | 0-90 | | | 0-90 | | | 0-90 | | 0-90 |
| <-99 - 90 points | | | nH | | | nH | | | nH | | nH | |
| | 0-1 | | P.1 | 0-1 | | pri | 0-1 | | pri | 0-1 | pri | 0-1 |
| 6.0-8.0 = 80 points | 0-80 6.81 | | | 5-90 | | | 5-90 | | | 5-90 | | 5-90 |
| 0 | | | DO | · · · · · · · · · · · · · · · · · · · | | po | | | DO | | DO | |
| | 10-30 7.37 | | | 10-30 | | | 10-30 | | | 10-30 | | 10-30 |
| >5.0 = 30 points | 10-30 7.37 | | | 10-30 | | | 10-30 | | | 10-30 | | 10-30 |
| ib-Total | 1 | | Sub-Total | 0 | | Sub-Total | | 0 | Sub-Total | 0 | Sub-Total | |
| OLOGICAL INDICATOR (Applies to Intermitt | tent and Derennial Streams) | | BIOLOGICAL INDICATOR (Applies to Intermitte | ent and Perennial Streame) | | BIOLOGICAL INDICATOR (Applies to | Intermittent and Percent | vial Streame) | BIOLOGICAL INDICATOR (Applies to Int | armittent and Perennial Streams) | BIOLOGICAL INDICATOR (Applies to Inter | nittent and Perennial Stress |
| V Stream Condition Index (WVSCI) | , | | WV Stream Condition Index (WVSCI) | , | | WV Stream Condition Index (WVSCI | | | WV Stream Condition Index (WVSCI) | , | WV Stream Condition Index (WVSCI) | |
| 0 | 0-100 0-1 | | | 0-100 0-1 | | | 0-100 0-1 | | | 0-100 0-1 | Site of the second second second second | 0-100 0-1 |
| U ub-Total | 0 | - | Sub-Total | 0 | | Sub-Total | | 0 | Sub-Total | 0 | Sub-Total | 0 |
| PART II - Index and U | Jnit Score | | PART II - Index and | Unit Score | - | PART II - Inde | ex and Unit Score | 1 | PART II - Index an | d Unit Score | PART II - Index and | Unit Score |
| | | | | | | | | | | | | |
| Index | Linear Feet Unit Scor | e | Index | Linear Feet Unit Scor | e | Index | Linear Feet | Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet Unit S |
| | | | | | | | | | | | | |

0.845

35.49

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | LOCATION | | | | | |
|---------------------|--------------|-------------------|--|--|--|--|
| STATION # RIVERMILE | STREAM CLASS | | | | | |
| LAT LONG | RIVER BASIN | | | | | |
| STORET # | AGENCY | | | | | |
| INVESTIGATORS | | | | | | |
| FORM COMPLETED BY | DATE TIME | REASON FOR SURVEY | | | | |

| WEATHER CONDITIONS | Now Past 24 hours Has there been a heavy rain in the last 7 days? Storm (heavy rain) rain (steady rain) showers (intermittent) % Air Temperature0 C % %cloud cover clear/sunny 0 |
|----------------------------|---|
| SITE LOCATION/MAP | Draw a map of the site and indicate the areas sampled (or attach a photograph) |
| STREAM CHARACTERIZATION | Stream Subsystem Perennial Stream Type Coldwater Warmwater Stream Origin Glacial Spring-fed Mixture of origins Swamp and bog Spring-fed |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential Indicate the dominant type and record the dominant Trees Shrubs Dominant species present | Local Watershed NPS Pollution No evidence Some potential sources Obvious sources Local Watershed Erosion None Moderate Heavy ant species present Grasses Herbaceous |
|--|--|---|
| INSTREAM FEATURES | Estimated Reach Length m Estimated Stream Width m Sampling Reach Area m² Area in km² (m²x1000) km² Estimated Stream Depth m Surface Velocity (at thalweg) m/sec | Canopy Cover Partly open Partly shaded Shaded High Water Mark m High Water Mark m Proportion of Reach Represented by Stream Morphology Types Riffle % Pool % Channelized Yes No Dam Present Yes No |
| LARGE WOODY DEBRIS | LWDm ² Density of LWDm ² /km ² (LWD/ reac | h area) |
| AQUATIC VEGETATION | Indicate the dominant type and record the dominant record the dominant type and record the dominant Rooted submergent Rooted submergent Attached Algae Dominant species present Portion of the reach with aquatic vegetation | Rooted floating Free floating |
| WATER QUALITY (DS) | Temperature0 C Specific Conductance Dissolved Oxygen pH Turbidity WQ Instrument Used | Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Slick Sheen Globs Furbidity (if not measured) Clear Slightly turbid Clear Slightly turbid Turbid Opaque Stained Other |
| SEDIMENT/ SUBSTRATE | Odors Petroleum Normal Sewage Petroleum Chemical Anaerobic None Other | Deposits Paper fiber Sand Sludge Sawdust Paper fiber Sand Relict shells Other |

| INC | DRGANIC SUBSTRATE (should add up to | | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | | | | | |
|-------------------|--|--|---|---|-------------------|--|--|--|--|--|--|
| Substrate Type | | | Diameter % Composition in Sampling Reach | | Substrate Type | Characteristic % Composition Sampling Arc | | | | | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | | | | | | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | | | | | | | |
| Cobble | 64-256 mm (2.5"-10") | | Muck-Mud | black, very fine organic | | | | | | | |
| Gravel | 2-64 mm (0.1"-2.5") | | | (FPOM) | | | | | | | |
| Sand | 0.06-2mm (gritty) | | Marl | grey, shell fragments | | | | | | | |
| Silt | 0.004-0.06 mm | | | | | | | | | | |
| Clay | < 0.004 mm (slick) | | | | | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | | | | |
|---------------------|--------------------|-------------------|--|--|--|
| STATION # RIVERMILE | STREAM CLASS | | | | |
| LAT LONG | RIVER BASIN | | | | |
| STORET # | AGENCY | | | | |
| INVESTIGATORS | | | | | |
| FORM COMPLETED BY | DATE TIME AM PM | REASON FOR SURVEY | | | |

| | Habitat | | Condition | ı Category | | | | | |
|--|---|---|---|---|---|--|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | | | |
| ted i | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | | | | |
| uram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | | | | |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Condition | ı Category | | | | |
|---|--|--|--|---|--|--|--|
| Parameter | Optimal | Suboptimal | Marginal | Poor | | | |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | | |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |
| SCORE(RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | |

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME | | LOCATION | | | | | | | |
|----------------------|----------------------------|--|-------------------|--|--|--|--|--|--|
| STATION # | _ RIVERMILE | STREAM CLASS | | | | | | | |
| LAT | LONG | RIVER BASIN | | | | | | | |
| STORET # | | AGENCY | | | | | | | |
| INVESTIGATORS | | | LOT NUMBER | | | | | | |
| FORM COMPLETED | BY | DATE TIME | REASON FOR SURVEY | | | | | | |
| HABITAT TYPES | Cobble% Sn | Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()% | | | | | | | |
| SAMPLE COLLECTION | Indicate the number of jab | lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B | anks Sand | | | | | | |
| GENERAL COMMENTS | | | | | | | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |


WOLMAN PEBBLE COUNT FORM


| County: | Lewis | Stream ID: |
|--------------|------------------|------------|
| Stream Name: | Second Big Run | |
| HUC Code: | | Basin: |
| Survey Date: | 9/13/2021 | |
| Surveyors: | MAG/DPF | Impact Rea |
| Туре: | Bankfull Channel | |
| | | |

S-VV13b


pact Reach: 14.1m

| | | | LE COUNT | | | | |
|-------------|--------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | • | 0 | 0.00 | 0.00 |
| | Very Fine | .062125 | SAND | • | 0 | 0.00 | 0.00 |
| | Fine | .12525 | | • | 3 | 3.00 | 3.00 |
| | Medium | .255 | | ▲ ▼ | 7 | 7.00 | 10.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | 9 | 9.00 | 19.00 |
| .0408 | Very Coarse | 1.0-2 | | ▲ ▼ | 6 | 6.00 | 25.00 |
| .0816 | Very Fine | 2 -4 | GRAVEL | ▲ ▼ | 0 | 0.00 | 25.00 |
| .1622 | Fine | 4 -5.7 | | • | 4 | 4.00 | 29.00 |
| .2231 | Fine | 5.7 - 8 | | • | 2 | 2.00 | 31.00 |
| .3144 | Medium | 8 -11.3 | | • | 1 | 1.00 | 32.00 |
| .4463 | Medium | 11.3 - 16 | | ▲ ▼ | 2 | 2.00 | 34.00 |
| .6389 | Coarse | 16 -22.6 | | ▲ ▼ | 4 | 4.00 | 38.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | ▲ ▼ | 17 | 17.00 | 55.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | ▲ ▼ | 14 | 14.00 | 69.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | ▲ ▼ | 18 | 18.00 | 87.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ▲ ▼ | 11 | 11.00 | 98.00 |
| 3.5 - 5.0 | Small | 90 - 128 | CODDIE | ▲ ▼ | 2 | 2.00 | 100.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | • | 0 | 0.00 | 100.00 |
| 7.1 - 10.1 | Large | 180 - 256 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 10.1 - 14.3 | Small | 256 - 362 | BOULDER | ▲ ▼ | 0 | 0.00 | 100.00 |
| 14.3 - 20 | Small | 362 - 512 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 20 - 40 | Medium | 512 - 1024 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 | | ▲ ▼ | 0 | 0.00 | 100.00 |
| | Bedrock | | BDRK | ▲ ▼ | 0 | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |
| | Total Tally: | | | | | | |





| | PRE-CROSSING | | | AS-BUILT | |
|----------|--------------|------------|--------|----------------|----------------|
| PT. LOC. | NORTHING | EASTING | ELEV | VERT. DIFF. | HORZ. DIFF. |
| TS-L | 14125918.43 | 1763231.45 | 844.61 | | |
| BS-L | 14125922.56 | 1763231.02 | 842.50 | | |
| тнw | 14125927.57 | 1763230.51 | 841.52 | | |
| BS-R | 14125936.14 | 1763229.62 | 841.78 | | |
| TS-R | 14125937.36 | 1763229.49 | 842.87 | | |



 SURVEY NOTES:

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON SEPTEMBER 13, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.
- 5. POST-CROSSING SURVEY INFORMATION SHOWN IN RED. DATA PENDING.
- 6. POST-CROSSING SURVEY POINTS FOR CROSS SECTIONS AND THALWEG ARE PROJECTED ONTO PRE-CROSSING SECTION AND PROFILE VIEWS FOR COMPARISON.

