Reach S-A32 (Pipeline ROW) Perennial Spread G Giles County, Virginia

Data	Included
Photos	\checkmark
SWVM Form	√
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope
	>4%)
RBP Physical Characteristics Form	\checkmark
Water Quality Data	\checkmark
RBP Habitat Form	\checkmark
RBP Benthic Form	\checkmark
Benthic Identification Sheet	\checkmark
Wolman Pebble Count	√
RiverMorph Data Sheet	\checkmark
USM Form (Virginia Only)	\checkmark
Longitudinal Profile and Cross Sections	\checkmark

Giles County



Photo Type: RB DS VIEW Location, Orientation, Photographer Initials: Standing on RB looking downstream along the ROW looking S, TC



Photo Type: LB DS VIEW Location, Orientation, Photographer Initials: Standing on LB looking downstream along the ROW looking SW, TC

Giles County



Photo Type: RB US VIEW Location, Orientation, Photographer Initials: Standing on RB looking upstream along the ROW looking N, TC



Photo Type: LB US VIEW Location, Orientation, Photographer Initials: Standing on LB looking upstream along the ROW looking N, TC



Photo Type: RB CL Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking E, ES



Photo Type: LB CL Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking W, ES



Photo Type: DS COND Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking S, ES

 $L: |22000s| 22800| 22865.06| Admin| 05-ENVR| Field Data| Spread G| Field Forms| S-A32| S-A32_Photo Doc_BKF10 plus. docx and the second state of the second state of$


Page 4

DEQ Permit #21-0416

	USACE FILE NO./ Project Name: (v2.1, Sept 2015)		Mountain	Valley Pipeline	IMPACT COORDINATES: (in Decimal Degrees)	Lat.	37.335094	Lon.	-80.596868	WEATHER:	Intermittent Showers	DATE:	8/20/2021 & 8/31/2021
Autor <th></th> <th></th> <th></th> <th>S-4</th> <th>132</th> <th></th> <th></th> <th></th> <th></th> <th></th> <th></th> <th>Comments:</th> <th></th>				S-4	132							Comments:	
<form>Added and participation of the product of the product</form>	STREAM IMPACT LENGTH:	78		RESTORATION (Levels I-III)		Lat.		Lon.		PRECIPITATION PAST 48 HRS:	1.25",8/18/21&8/19/21; 0.58",8/	80/21 Mitigation Length:	
	Column No. 1- Impact Existing	g Condition (Deb	pit)	Column No. 2- Mitigation Existing Co	ondition - Baseline (Credit)				Years			Column No. 5- Mitigation Project	ed at Maturity (Credit)
	Stream Classification:	Perer	nnial	Stream Classification:			Stream Classification:		0	Stream Classification:	0	Stream Classification:	0
	Percent Stream Channel SI	lope	6.59	Percent Stream Channel Slo	pe		Percent Stream Chann	nel Slope	0	Percent Stream Channel Si	ope 0	Percent Stream Channel S	lope 0
	HGM Score (attach d	lata forms):		HGM Score (attach d	ata forms):		HGM Score (at	tach data forms):		HGM Score (attach da	ata forms):	HGM Score (attach d	ata forms):
<form> martine martine martine <</form>			Average		Average				Average		Average		Average
<form> PATI - Prycial Chancial and Biological Productional Control Productional Control</form>			0						0		0		0
		Biological Indica	ators		Biological Indicators			cal and Biological In	dicators		Biological Indicators		Biological Indicators
		Points Scale Range	Site Score		Points Scale Range Sille Score			Points Scale Range	Site Score		Points Scale Range Site Score		Points Scale Range Site Score
	PHYSICAL INDICATOR (Applies to all streams	s classifications)		PHYSICAL INDICATOR (Applies to all streams of	assifications)		PHYSICAL INDICATOR (Applies to all st	reams classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)	PHYSICAL INDICATOR (Applies to all streams	classifications)
	USEPA RBP (High Gradient Data Sheet)			USEPA RBP (Low Gradient Data Sheet)			USEPA RBP (High Gradient Data She	eet)		USEPA RBP (High Gradient Data Sheet)		USEPA RBP (High Gradient Data Sheet)	
			16										
			16										
			8										
			18		0-20								
	6. Channel Alteration	0-20	19	6. Channel Alteration	0-20		6. Channel Alteration	0-20		6. Channel Alteration	0-20	6. Channel Alteration	0-20
0. Vogetate Production (L8 A BS) 0.00 1.00 0.00	7. Frequency of Riffles (or bends)	0-20		7. Channel Sinuosity	0-20		7. Frequency of Riffles (or bends)	0-20		7. Frequency of Riffles (or bends)	0-20	7. Frequency of Riffles (or bends)	0-20
		0-20			0-20			0-20			0-20		0-20
<form> Sub_chi Sub_chi<td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></form>													
<form> Sub_Tidd 0.726 Sub_Tidd Sub_Tidd<</form>													
		Suboptimai						Poor	0				
Specific Conductivity 33.4 0 33.4 0 0 0		nt and Perennial Stre						mittent and Perennial St	-				
$ \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} \begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \end{array} \\ 0 \\ 0 \\$	WVDEP Water Quality Indicators (General	ŋ		WVDEP Water Quality Indicators (General)			WVDEP Water Quality Indicators (Ge	neral)		WVDEP Water Quality Indicators (General)	WVDEP Water Quality Indicators (General)
$ \begin{array}{c} $	Specific Conductivity	_		Specific Conductivity			Specific Conductivity			Specific Conductivity		Specific Conductivity	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	300-399 - 70 points	0-90	333.4		0-90			0-90			0-90		0-90
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	pH			pH			pH			pH		pH	
BOD	8 1-9 0 = 45 points	0-80	8.61		5-90			5-90			5-90		5-90
2 + 50 = 30 ords $2 + 50 = 30 ords$ $3 + 50 ords$ <td< td=""><td>DO</td><td></td><td></td><td>DO</td><td></td><td></td><td>DO</td><td></td><td></td><td>DO</td><td></td><td>DO</td><td></td></td<>	DO			DO			DO			DO		DO	
Sub-Total O.7 Sub-Total O Sub-Total O Sub-Total O Sub-Total O BloLOGICAL INDICATOR (Applies to Intermiter and Paral and Control (A NDICATOR (Applies to Intermi		10-30	6.77		10-30			10-30			10-30		10-30
B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): WStream Condition Index (WVSC): WStream Condition Index (WVSC): B0LOGICAL INDICATOR (Applies to Intermitted and Particle Definition (MVSC): B0LOGICAL INDICATOR (Applies to Intermitted And Particle Definition (MVSC): WStream Condition Index (WVSC): WStream Condition Index (WVSC): WStream Condition Index (WVSC): WStream Condition Index (WVSC): B0LOGICAL INDICATOR (Applies to Intermitted And Particle Definition (MVSC): WStream Condition Index (WVSC): WStream Condition Index (WVSC): W	>5.0 = 30 points Sub-Total		0.725	Sub-Total	0		Sub-Total		0	Sub-Total		Sub-Total	
W Stream Condition Index (WVSC) W St		ttent and Perennial S			and Perennial Streams)			Intermittent and Peren	aial Streams)		ittent and Perennial Streams)		ittent and Perennial Streame)
And the set of			,		,								
Groy Zone Groy Zone Groy Zone Groy Zone Groy Zone Grow		0-100 0-1	65.3		0-100 0-1		(HVOCI)				0-100 0-1		0-100 0-1
PART II - Index and Unit Score Index				L									
Image: Note of the second se	Sub-Total		0.653	Sub-Total	0		Sub-Total		0	Sub-Total	0	Sub-Total	0
	PART II - Index and U	Unit Score		PART II - Index and L	Jnit Score		PART II - Inde	x and Unit Score		PART II - Index and U	nit Score	PART II - Index and L	Init Score
0.701 78 54.878 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	Index	Linear Feet	Unit Score	Index	Linear Feet Unit Score		Index	Linear Feet	Unit Score	Index	Linear Feet Unit Score	Index	Linear Feet Unit Score
	0.701	78	54.678	0	0 0		0	0	0	0	0 0	0	0 0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME S-A32	LOCATION Giles County	
STATION # 10882+96 RIVERMILE	STREAM CLASS Perennial	
LAT <u>37.335094</u> LONG <u>-80.596868</u>	RIVER BASIN Middle New	
STORET #	AGENCY VADEQ	
INVESTIGATORS ES, EM		
FORM COMPLETED BY ES	DATE 8/20/2021 TIME 2:30 PM	REASON FOR SURVEY Baseline Assessment



Notes: Recent storm event.

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer)	Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential Indicate the dominant type and record the dominant species present Solidago and Impatients capensis	Local Watershed NPS Pollution ☑ No evidence □ Some potential sources □ Obvious sources Local Watershed Erosion ☑ None □ Moderate □ Moderate □ Heavy nant species present □ Herbaceous
INSTREAM FEATURES	Estimated Reach Length 17.88 m Estimated Stream Width 1.5 m Sampling Reach Area 26.52 m² Area in km² (m²x1000) km² Estimated Stream Depth 0.15 m Surface Velocity (at thalweg) m/sec	Canopy Cover □Partly shaded □Shaded Image: Partly open □Partly shaded □Shaded High Water Mark 02 m Proportion of Reach Represented by Stream Morphology Types Riffle 40 % Pool % Run 30 % Channelized Yes Dam Present Yes
LARGE WOODY DEBRIS	LWDm ² Density of LWDm ² /km ² (LWD/ read	ch area)
AQUATIC VEGETATION	Indicate the dominant type and record the domin Rooted emergent Floating Algae Dominant species present Persicaria sp. Portion of the reach with aquatic vegetation 25	nant species present ☐Rooted floating ☐Free floating _%
WATER QUALITY (DS, US)	Temperature 22.4, 23.1 0 C Specific Conductance 333.4, 328.0 uS/cm Dissolved Oxygen 6.77, 7.75 mg/L pH 8.61, 8.62 Turbidity N/A WQ Instrument Used VA-4	Water Odors Petroleum Petroleum Fishy Bitck Slick Other Turbidity (if not measured) Clear Opaque Stained
SEDIMENT/ SUBSTRATE	Odors ✓ ✓ Normal Chemical Anaerobic Other None Oils ✓ ✓ Absent Slight	Deposits □Sludge □Sawdust □Paper fiber □Sand □Relict shells □Other □ Lpoking at stones which are not deeply embedded, are the undersides black in color? □ Yes ☑ No

INC	DRGANIC SUBSTRATE (should add up to 1			ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)						
Substrate Type	Diameter	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area					
Bedrock		25	Detritus	sticks, wood, coarse plant	10					
Boulder	> 256 mm (10")	0		materials (CPOM)	10					
Cobble	64-256 mm (2.5"-10")	10	Muck-Mud	black, very fine organic	0					
Gravel	2-64 mm (0.1"-2.5")	40		(FPOM)	0					
Sand	0.06-2mm (gritty)	15	Marl	grey, shell fragments	0					
Silt	0.004-0.06 mm	5]		0					
Clay	< 0.004 mm (slick)	5]							

Notes: Recent storm event.

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-A32	LOCATION Giles County
STATION #_10882+96 RIVERMILE	STREAM CLASS Perennial
LAT <u>37.335094</u> LONG <u>-80.596868</u>	RIVER BASIN Middle New
STORET #	AGENCY VADEQ
INVESTIGATORS ES, EM	
FORM COMPLETED BY ES	DATE 8/20/2021 REASON FOR SURVEY TIME 2:30 PM AM PM Baseline Assessment

	Habitat		Condition	Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.				
	_{SCORE} 16	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.				
ted iı	score 8	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).				
uram	_{score} 16	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
P	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.				
	_{score} 8	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.				
	_{score} 18	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				

Notes: Recent storm event.

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	n Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.				
	_{score} 19	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
ling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.				
ampl	_{score} 18	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing downstream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.				
e evs	SCORE 9	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
to b	SCORE 10	Right Bank 10 9	8 7 6	5 4 3	2 1 0				
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.				
	SCORE 7	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
	SCORE 6	Right Bank 10 9	8 7 6	5 4 3	2 1 0				
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.				
	SCORE 4	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
L	SCORE 6	Right Bank 10 9	8 7 6	5 4 3	2 1 0				

145 Notes:

Notes: Recent storm event.

Total Score ____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-A	32	LOCATION Giles County	
STATION #_10882+96	RIVERMILE	STREAM CLASS Perennial	
LAT37.335094	LONG80.596868	RIVER BASIN Middle New	
STORET #		AGENCY VADEQ	
INVESTIGATORS ES			LOT NUMBER
FORM COMPLETED	^{BY} ES, DW	DATE 8/31/2021 TIME 10:30 AM	REASON FOR SURVEY Baseline Assessment
HABITAT TYPES	Indicate the percentage of ✓Cobble 25 % Sn Submerged Macrophytes	ags%	
SAMPLE COLLECTION		lected? □wading □fi s/kicks taken in each habitat ty lags □Vegetated B	rom bank
GENERAL COMMENTS	4 kicks collected		,

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

Mountain Valley Pipeline Data are not adjusted for subsampling

ECO ANALYSTS, INC.

	Sample ID Collection Date	S-A32 08-31-2021
ORDER GENUS/SF	PECIES	COUNT
Ephemeroptera Diphetor hageni		9
Ephemeroptera Habrophlebiodes sp.		7
Ephemeroptera Isonychia sp.		4
Plecoptera Leuctra sp.		2
Trichoptera Cheumatopsyche sp.		16
Trichoptera Chimarra sp.		4
Trichoptera Diplectrona sp.		1
Trichoptera Hydropsyche sp.		2
Trichoptera Lepidostoma sp.		2
Coleoptera Optioservus sp.		4
Coleoptera Psephenus sp.		112
Coleoptera Stenelmis sp.		12
Diptera-Chironomidae Paracricotopus sp.		1
Diptera-Chironomidae Polypedilum sp.		2 2
Diptera-Chironomidae Thienemannimyia gr.	sp.	2
Diptera Ceratopogoninae		3
Annelida Lumbricina		1
Annelida tubificoid Naididae w/	cap setae	2
Gastropoda Physa sp.		1
Crustacea Gammarus sp.		12
Other Organisms Turbellaria		6
	TOTAL	205

Mountain Valley Pipeline WV SCI Metrics

ECO ANALYSTS, INC.

Sample ID Collection Date	
WVSCI Metric Values	
Total taxa	16
EPT taxa	7
% EPT	22.9
% Chironomidae	2.4
% 2 Dominant HBI	63.9 4.20
пы	4.20
WVSCI Metric Scores	
Total taxa	76.2
EPT taxa	53.8
% EPT % Chironomidae	24.9 98.5
% 2 Dominant	98.5 56.4
HBI	81.7
WVSCI Metric Scores	70.0
Total taxa EPT taxa	76.2 53.8
% EPT	24.9
% Chironomidae	98.5
% 2 Dominant	56.4
НВІ	81.7
WVSCI Total Score	65.3

WVSCI Thresholds

Unimpaired = > 68.00 Gray Zone = 60.61 to 68.00 Impaired = <60.61

WOLMAN PEBBLE COUNT FORM

County:	Giles County
Stream Name:	UNT to Doe Creek
HUC Code:	02080201
Survey Date:	8/20/2021
Surveyors:	ES, EM
Туре:	Representative

Stream ID:

Basin:

Middle New

S-A32


			LE COUNT				a. (
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	▲ ▼	9	9.00	9.00
	Very Fine	.062125		▲ ▼	0	0.00	9.00
	Fine	.12525		▲ ▼	0	0.00	9.00
	Medium	.255	SAND	▲ ▼	1	1.00	10.00
	Coarse	.50-1.0		▲ ▼	8	8.00	18.00
.0408	Very Coarse	1.0-2		▲ ▼	3	3.00	21.00
.0816	Very Fine	2 -4		▲ ▼	2	2.00	23.00
.1622	Fine	4 -5.7		▲ ▼	5	5.00	28.00
.2231	Fine	5.7 - 8		▲ ▼	9	9.00	37.00
.3144	Medium	8 -11.3	GRAVEL	▲ ▼	11	11.00	48.00
.4463	Medium	11.3 - 16		▲ ▼	5	5.00	53.00
.6389	Coarse	16 -22.6		▲ ▼	2	2.00	55.00
.89 - 1.26	Coarse	22.6 - 32		▲ ▼	3	3.00	58.00
1.26 - 1.77	Vry Coarse	32 - 45		▲ ▼	3	3.00	61.00
1.77 -2.5	Vry Coarse	45 - 64		▲ ▼	4	4.00	65.00
2.5 - 3.5	Small	64 - 90		▲ ▼	0	0.00	65.00
3.5 - 5.0	Small	90 - 128	CODDIE	▲ ▼	3	3.00	68.00
5.0 - 7.1	Large	128 - 180	COBBLE	▲ ▼	3	3.00	71.00
7.1 - 10.1	Large	180 - 256		▲ ▼	1	1.00	72.00
10.1 - 14.3	Small	256 - 362		▲ ▼	0	0.00	72.00
14.3 - 20	Small	362 - 512		▲ ▼	0	0.00	72.00
20 - 40	Medium	512 - 1024	BOULDER	▲ ▼	0	0.00	72.00
40 - 80	Large	1024 -2048	1	▲ ▼	0	0.00	72.00
80 - 160	Vry Large	2048 -4096	1	▲ ▼	0	0.00	72.00
	Bedrock		BDRK	▲ ▼	28	28.00	100.00
	1			Totals	100		

Reach Name: S- Sample Name: Re	T to Doe Cre A32 presentative /20/2021			
Size (mm)	тот #	ITEM %	CUM %	
$\begin{array}{r} 0 & - & 0.062 \\ 0.062 & - & 0.125 \\ 0.125 & - & 0.25 \\ 0.25 & - & 0.50 \\ 0.50 & - & 1.0 \\ 1.0 & - & 2.0 \\ 2.0 & - & 4.0 \\ 4.0 & - & 5.7 \\ 5.7 & - & 8.0 \\ 8.0 & - & 11.3 \\ 11.3 & - & 16.0 \\ 16.0 & - & 22.6 \\ 22.6 & - & 32.0 \\ 32 & - & 45 \\ 45 & - & 64 \\ 64 & - & 90 \\ 90 & - & 128 \\ 128 & - & 180 \\ 180 & - & 256 \\ 256 & - & 362 \\ 362 & - & 512 \\ 512 & - & 1024 \\ 1024 & - & 2048 \\ Bedrock \end{array}$	9 0 1 8 3 2 5 9 11 5 2 3 3 4 0 3 3 1 0 0 0 0 2 8	9.00 0.00 0.00 1.00 8.00 3.00 2.00 5.00 9.00 11.00 5.00 2.00 3.0	9.00 9.00 9.00 10.00 18.00 21.00 23.00 28.00 37.00 48.00 53.00 55.00 58.00 61.00 65.00 65.00 65.00 65.00 65.00 65.00 72.00 72.00 72.00 72.00 72.00 72.00	
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Gravel (%) Boulder (%) Bedrock (%)	0.88 7.49 13.18 Bedrock Bedrock 9 12 44 7 0 28			


Total Particles = 100.


			Strear		essm tream Method			form 1)		
					able channels cla			al			
Project #	Projec	ct Name (App		Locality	Cowardin Class.	HUC	Date	SAR #	Impact Length	Impact Factor	
22865.06		alley Pipeline ey Pipeline, L		Giles County	R3	05050002	8/20/2021	S-A32	78	1	
Nam	e(s) of Evaluat	tor(s)	Stream Name	e and Informa	tion				SAR Length		
	ES, EM		UNT to Doe (Creek					7	8	
. Channel C	condition: Asses	ss the cross-secti	on of the stream a								
	Optimal Subopti				Conditional Catego	ginal	Po	or	Sev	vere	
Channel Condition	Very little incision or 100% stable banks. protection or nature (80-100%). AND/OF bankfull benches ar to their original fi developed wide ban channel bars and tr Transient sediment less than 109	r active erosion; 80- Vegetative surface al rock, prominent 8 Stable point bars / re present. Access loodplain or fully kfull benches. Mid- ansverse bars few.	Slightly incised, fr erosion or unprotect of banks are s Vegetative protec prominent (60 Depositional feat stability. The ban channels are well d has access to ba	ew areas of active ted banks. Majority table (60-80%). tion or natural rock -80%) AND/OR ures contribute to hkfull and low flow fined. Stream likely inkfull benches, or floodplains along	Often incised, but Poor. Banks more or Poor due to Ic Erosion may be pr both banks. Veget 40-60% of banks. S vertical or und 40-60% Sediment transient, contr	less than Severe or stable than Severe	Overwidened/in laterally unstable further. Majority of banks. vegttaal. Erosion pr banks. Vegetative on 20-40% of bank to prevent erosion. the stream is cov Sediment is temp	cised. Vertically / . Likely to widen both banks are near esent on 60-80% of protection present s, and is insufficient AND/OR 60-80% of ered by sediment. orary / transient in buting to instability.	Deeply incised vertical/lateral in incision, flow control Streambed below av majority of banks Vegetative protect than 20% of banks erosion. Obviou present. Erosion/raw	(or excavated), stability. Severe red within the banks. erage rooling depth, vertical/undercut. ion present on less s, is not preventing s bank sloughing	
			portions of the r sediment covers 1 bot	each. Transient 0-40% of the stream tom.	may be forming/pr shaped channels protection on > 40 depositional feature to sta	resent. AND/OR V- s have vegetative % of the banks and es which contribute ability.	AND/OR V-shap vegetative protect 40% of the banks a deposition	bed channels have tion is present on > and stable sediment h is absent.	than 80% of stream deposition, contrib Multiple thread subterran	n bed is covered by buting to instability. channels and/or nean flow.	CI
Scores	3	3	2	.4		2	1	.6	-	1	2.40
	Opti			ditional Cate	gory				NOTES>>		
		imai	Subo	ptimal	Mar	ginal		oor	NOTES		
Riparian Buffers	Tree stratum (dbh > with > 60% tree Wetlands located i are:	 3 inches) present, canopy cover. within the riparian 	Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	Low Suboptimal:	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production,	Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NUTE3		
•	with > 60% tree Wetlands located	 3 inches) present, canopy cover. within the riparian 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30%	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable			
•	with > 60% tree Wetlands located	 3 inches) present, canopy cover. within the riparian as. 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.	High Poor: Lawns, mowed, and maintained areas, nurseries, no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NUTE3		
Buffers Scores Delineate ripa	with > 60% tree Wetlands located area	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegetated non-maintained area; recently seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NUTE3		
Buffers Scores Delineate ripa Determine sq	with > 60% tree Wetlands located area 1. Trian areas along ex uare footage for ea tiparian Area and S	 3 inches) present, e canopy cover. within the riparian as. 5 ach stream bank ach by measuring 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cal	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegret, seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5	NUTE3		
Buffers Scores Delineate ripa Determine sq	with > 60% tree Wetlands located area 1. Trian areas along ex uare footage for ea Riparian Area and S % Riparian Area>	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ach by measuring Score for each ripa 50% 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30%	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cal- be blocks below. 20%	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegret, seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lost, trails, or other comparable conditions.	NUTE3		
Buffers Scores Delineate ripa Determine sq Enter the % R	with > 60% tree Wetlands located area 1. Trian areas along ex uare footage for ea tiparian Area and S	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ich by measuring score for each ripar 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cal-	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegret, seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lost, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100	CI= (Sum % RA * Sc	pores*0.01)/2	
Buffers Scores Delineate ripa Determine sq Enter the % R Right Bank	with > 60% tree Wetlands located are 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ach by measuring Score for each ripa 50% 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30%	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cal- be blocks below. 20%	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegret, seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lost, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100		ores*0.01)/2	ŭ
Buffers Scores Delineate ripa Determine sq Enter the % R	with > 60% tree Wetlands located arei 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ach by measuring Score for each ripa 50% 0.75 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30% 0.6	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call th and width. Call th blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegret, seeded and stabilized, or other comparable condition. High 0.6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lost, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100	CI= (Sum % RA * Sc		CI 0.69
Buffers Scores Delineate ripa Determine sq Enter the % F Right Bank Left Bank . INSTREAN	with > 60% tree Wetlands located are 1 . Arian areas along ea uare footage for ea Riparian Area and S % Riparian Area> Score > % Riparian Area> Score > 1 HABITAT: Var	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ach by measuring Score for each riparian 50% 0.75 45% 0.5 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30% 0.6	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call th and width. C	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provid	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	Cl= (Sum % RA * Sc Rt Bank Cl >	0.66 0.73	
Buffers Scores Delineate ripa Determine sq Enter the % R Right Bank Left Bank	with > 60% tree Wetlands located are 1 . Arian areas along ea uare footage for ea Riparian Area and S % Riparian Area> Score > % Riparian Area> Score > 1 HABITAT: Var	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank ach by measuring Score for each riparian 50% 0.75 45% 0.5 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30% 0.6	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call th and width. Call th and width. Call th blocks below. 20% 0.5 25% 1.1 and depths; wood	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dhh > 3 inches) present, with <30% tree canopy cover. High 0.85 tion Scores using culators are provid	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI >	0.66 0.73	
Buffers Scores Delineate ripa Determine sq Enter the % F Right Bank Left Bank	with > 60% tree Wetlands located are 1 . Arian areas along ea uare footage for ea Riparian Area and S % Riparian Area> Score > % Riparian Area> Score > 1 HABITAT: Var	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank to by measuring Score for each ripa 50% 0.75 45% 0.5 ried substrate size 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30% 0.6 30% 0.75 as, water velocity a	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call th and width. Call th and width. Call th blocks below. 20% 0.5 25% 1.1 and depths; wood	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dhb > 3 inches) present, with <30% tree canopy cover. High 0.85 ttion Scores using culators are provid	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors.	High Poor: Lawns, mowed, and maintained areas, parsely egetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$	0.66 0.73	
Scores Delineate ripa Determine sq Enter the % F Right Bank Left Bank S. INSTREAN omplexes, stabl	with > 60% tree Wetlands located area International and S Version and S Riparian Area and S Riparian Area Score > Riparian Area Score > A HABITAT: Vari le features.	 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank the by measuring Score for each ripa 50% 0.75 45% 0.5 ried substrate size imal re typically present 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating leng arian category in th 30% 0.6 30% 0.75 es, water velocity a Stable habitat ele present in 30-50% adequate for r	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call th and width. Call th and width. Call th blocks below. 20% 0.5 25% 1.1 and depths; woody	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provid	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below.	High Poor: Lawns, mowed, and maintained areas, sparsely vegetated non-maintained sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I Blocks e Blocks e Habitat elements lacking or are u elements are typic	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$	0.66 0.73 SAV; riffle/pool	0.69
Scores Scores Delineate ripa Determine sq Enter the % F Right Bank Left Bank . INSTREAN omplexes, stabl Instream Habitat/ Available	with > 60% tree Wetlands located are in a reas along ea uare footage for ea tiparian Area and S % Riparian Area> Score > % Albaita Area> Score > % Albaita Area> Score >	 3 inches) present, a canopy cover. within the riparian as. 5 ach stream bank ach stream bank ach by measuring Score for each riparian 50% 0.75 45% 0.5 ried substrate size imal re typically present 0% of the reach. 	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Catt or estimating leng arian category in th 30% 0.6 30% 0.75 as, water velocity a Stable habitat ele present in 30-50% adequate for r popul	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Call re blocks below. 20% 0.5 25% 1.1 and depths; wood ptimal ments are typically of the reach and are	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provid culators are provid	Low Marginal: Non-maintained, dense herbaceous vegetation, ripariar areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ted for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries: no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area; recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks 6 Blocks 6 Habitat elements lacking or are u elements are typic than 10% c	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100% 100% 5 cor s listed above are nstable. Habitat ally present in less	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$ NOTES>>	0.66 0.73 SAV; riffle/pool	

	S	tream Ir	npact A	ssessn	nent For	rm Page	2			
Project #	Project Name (Applicant)		Locality	Cowardin Class.	HUC	Date	SAR #	Impact Length	Impact Factor	
22865.06	Mountain Valley Pipeline (Mountain Valley Pipeline, LLC)		Giles County	R3	05050002	8/20/2021	S-A32	78	1	
. CHANNEL	ALTERATION: Stream crossin	ngs, riprap, concret	te, gabions, or con	icrete blocks, stra	ightening of chann	el, channelization			ons, livestock	
			Conditiona	al Category				NOTES>>		
	Negligible	Mir	nor		erate	Sev	/ere			
Channel Alteration	hardening absent. Stream has an unaltered pattern or has naturalized.	the channel	the channel	40 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	60 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	Greater than 80% of by any of the chanr in the parameter g 80% of banks sh riprap, or	nel alterations listed uidelines AND/OR ored with gabion, cement.			CI
Scores	1.5	1.3	1.1	0.9	0.7	0	.5			1.50
	REACH	CONDITION	INDEX and S	STREAM CO	NDITION UN	ITS FOR THI	S REACH			
OTE: The CIs a	nd RCI should be rounded to 2 deci	mal places. The Cl	R should be round	ed to a whole nun	nber.		THE REACH	H CONDITION IN	IDEX (RCI) >>	1.22
						RCI= (Sum of	all CI's)/5, exce	ept if stream is ep	hemeral RCI = (I	Riparian Cl
							COMPENSA	TION REQUIRE	MENT (CR) >>	95
							CR = RC	CI X L _I X IF		




PROVIDED UNDER SEPARATE COVER





CL STAKEOUT POINTS: S-A32 CROSS SECTION B (PIPE CL)									
	PR	POST-CF	ROSSING						
	NODTUNC	FACTING		VERT.	HORZ.				
PT. LOC.	NORTHING	EASTING	ELEV	DIFF.	DIFF.				
TS-L	13556802.66	1757595.87	2139.20						
BS-L	13556803.23	1757593.09	2137.52						
THW	13556803.75	1757589.77	2136.90						
BS-R	13556805.30	1757583.40	2137.91						
TS-R	13556806.63	1757576.89	2142.07						



SURVEY NOTES:

1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on October 17, 2018.


2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.


3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).

4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.

5. All section views shown are left to right facing downstream.

6. Cross-section B shot at location of pipe centerline (based on best professional judgement).





LEGEND

EXISTING SURVEY-LOCATED THALWEG

EXISTING SURVEY-LOCATED EDGE OF WATER (AS NECESSARY)

EXISTING CONTOUR LINE (MAJOR)

EXISTING SURVEYED GROUND SHOT ELEVATION

