Baseline Assessment - Stream Attributes

Reach S-G24 (Pipeline ROW) Intermittent Spread H Franklin County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	✓
RBP Physical Characteristics Form	✓
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	✓
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	√
Longitudinal Profile and Cross Sections	✓



Photo Type: DS VIEW Location, Orientation, Photographer Initials: Downstream view of ROW looking S, AW



Photo Type: US VIEW Location, Orientation, Photographer Initials: Upstream view of ROW looking N, AW



Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking W, AW



Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking NE, AW



Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking S, AW

USACE FILE NO./ Project Name: (v2.1, Sept 2015)		Mountain	Vally Pipeline		COORDINATES: imal Degrees)	Lat.	37.126412	Lon.	-80.121398	WEATHER:	Me	ostly Sunny		DATE:	August 25, 2021	
IMPACT STREAM/SITE ID			S	-G24			MITIGATION STREAM CLASS						Co	mments:		
(watershed size (acreage)	unaltered or impair	ments)					(watershed size (acrea	ge), unaltered or imp	airments)							
STREAM IMPACT LENGTH:	75	FORM OF MITIGATION:	RESTORATION (Levels I-III)		OORDINATES: imal Degrees)	Lat.		Lon.		PRECIPITATION PAST 48 HRS:	N PAST 48 HRS: Non		Mitiga	tion Length:		
Column No. 1- Impact Existing	olumn No. 1- Impact Existing Condition (Debit)		Column No. 2- Mitigation Existing Condition - Baseline (Credit)			Column No. 3- Mitigation I Post Completi	Projected at Five Years on (Credit)		Column No. 4- Mitigation Proj Post Completion	jected at Ten Ye (Credit)	ears	Column No. 5- Mitigation Proje		rojected at Maturity (Credit)		
Stream Classification:	Interm	nittent	Stream Classification:				Stream Classification:	(Stream Classification:	0		Stream Classification:		0	
Percent Stream Channel SI	оре	10.36	Percent Stream Channel S	lope			Percent Stream Channel	Slope	0	Percent Stream Channel S	lope	0	Perc	ent Stream Channel S	lope	0
HGM Score (attach d	ata forms):		HGM Score (attach	data forms):			HGM Score (attac	h data forms):		HGM Score (attach d	ata forms):			HGM Score (attach d	ata forms):	
		Average			Average				Average			Average				Average
Hydrology	0.55		Hydrology				Hydrology			Hydrology			Hydrology			
Biogeochemical Cycling	0.52	0.49	Biogeochemical Cycling		0		Biogeochemical Cycling		0	Biogeochemical Cycling		0	Biogeochemical C	Cycling		0
PART I - Physical, Chemical and	0.4 Biological Indica	ators	PART I - Physical, Chemical ar	nd Biological Indi	icators		PART I - Physical, Chemical	and Biological In	dicators	PART I - Physical, Chemical and	Biological Indic	cators	Habitat PART I -	Physical, Chemical and	Biological Ind	licators
	Points Scale Range	Site Score		Points Scale Range	Site Score			Points Scale Range	Site Score		Points Scale Range	Site Score			Points Scale Ran	nge Site Score
PHYSICAL INDICATOR (Applies to all streams	classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)			PHYSICAL INDICATOR (Applies to all stream	ms classifications)		PHYSICAL INDICATOR (Applies to all stream	s classifications)	1	PHYSICAL INDICA	ATOR (Applies to all streams	classifications)	
USEPA RBP (High Gradient Data Sheet)			USEPA RBP (Low Gradient Data Sheet)				USEPA RBP (High Gradient Data Sheet)			USEPA RBP (High Gradient Data Sheet)			USEPA RBP (High	Gradient Data Sheet)		
Epifaunal Substrate/Available Cover	0-20	18	Epifaunal Substrate/Available Cover	0-20			Epifaunal Substrate/Available Cover	0-20		Epifaunal Substrate/Available Cover	0-20			ate/Available Cover	0-20	
2. Embeddedness	0-20	11	Pool Substrate Characterization	0-20			2. Embeddedness	0-20		2. Embeddedness	0-20		Embeddedness		0-20	
Velocity/ Depth Regime Sediment Deposition	0-20	16	Pool Variability Sediment Deposition	0-20			Velocity/ Depth Regime Sediment Deposition	0-20		Velocity/ Depth Regime Sediment Deposition	0-20		 Velocity/ Depth F Sediment Depos 	Regime	0-20 0-20	
		14 16														
5. Channel Flow Status	0-20 0-1		5. Channel Flow Status	0-20 0-1			5. Channel Flow Status	0-20 0-1		5. Channel Flow Status	0-20 0-1		5. Channel Flow St		0-20	-1
6. Channel Alteration	0-20	19 18	6. Channel Alteration	0-20			6. Channel Alteration	0-20		6. Channel Alteration	0-20		6. Channel Alterati		0-20	
7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB)	0-20	18	7. Channel Sinuosity	0-20			7. Frequency of Riffles (or bends)	0-20		7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB)	0-20		7. Frequency of Rif		0-20	
	0-20	12	8. Bank Stability (LB & RB)	0-20			8. Bank Stability (LB & RB)	0-20			0-20		8. Bank Stability (L		0-20	
Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20	15	Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20			Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20		Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20		Vegetative Prote Ringrigs Vegetative	ction (LB & RB) ive Zone Width (LB & RB)	0-20 0-20	
Total RBP Score	Suboptimal	157	Total RBP Score	Poor	0		Total RBP Score	Poor	0	Total RBP Score	Poor	0	Total RBP Score	ive zone widin (LB & NB)	Poor	n
Sub-Total	•	0.785	Sub-Total		Ö		Sub-Total		Ö	Sub-Total		Ö	Sub-Total			ő
CHEMICAL INDICATOR (Applies to Intermitter		eams)	CHEMICAL INDICATOR (Applies to Intermitter		sams)		CHEMICAL INDICATOR (Applies to Intermit		treams)	CHEMICAL INDICATOR (Applies to Intermitte		treams)		ATOR (Applies to Intermitter		Streams)
WVDEP Water Quality Indicators (General Specific Conductivity)		WVDEP Water Quality Indicators (General Specific Conductivity)			WVDEP Water Quality Indicators (Gener Specific Conductivity	al)		WVDEP Water Quality Indicators (General Specific Conductivity	1)		WVDEP Water Qu Specific Conduct	ality Indicators (General)	
opcome conductivity			opecine conductivity	T			opecine conductivity			opcome conductivity	T		opecine conduct	ivity .		
<=99 - 90 points	0-90	66.6		0-90				0-90			0-90				0-90	
pH			pH				pH			рН			pH			
8.1-9.0 = 45 points	0-80	8.78		5-90				5-90			5-90				5-90	-1
8. 1-9.0 = 45 points	_		no				no			no			DO.			
	10-30	9.77		10-30				10-30			10-30				10-30	
>5.0 = 30 points	.0-30			10-30				10-30		L	10-30				10-30	
Sub-Total BIOLOGICAL INDICATOR (Applies to Intermit	tent and Perennial S	0.825	Sub-Total BIOLOGICAL INDICATOR (Applies to Intermit	tent and Perennial S	(treams)		Sub-Total BIOLOGICAL INDICATOR (Applies to Inte	rmittent and Peren	nial Streams)	Sub-Total BIOLOGICAL INDICATOR (Applies to Intern	mittent and Peren	nial Streams)	Sub-Total BIOLOGICAL IND	CATOR (Applies to Interm	ittent and Pere	nnial Streams)
WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)				WV Stream Condition Index (WVSCI)		, ,	WV Stream Condition Index (WVSCI)		, , , , , , , , , , , , , , , , , , , ,		tion Index (WVSCI)		, , , ,
	0-100 0-1	79.6		0-100 0-1			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	0-100 0-1		,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	0-100 0-1				0-100 0-	4
Very Good Sub-Total	1 1	0.796	Sub-Total		0		Sub-Total		0	Sub-Total		0	Sub-Total		1	0
PART II - Index and U	nit Score		PART II - Index and	Unit Score			PART II - Index a	nd Unit Score		PART II - Index and U	Init Score			PART II - Index and U	nit Score	
															Linear Foot Unit Comm	
Index	Linear Feet	Unit Score	Index	Linear Feet	Unit Score		Index	Linear Feet	Unit Score	Index	Linear Feet	Unit Score		Index	Linear Fee	t Unit Score
0.646	75	48.45	0	0	0		0	0	0	0	0	0		0	0	0

Ver. 10-20-17

FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the <u>UPPERMOST STRATUM</u> of the plant community is determined based on the calculated value for V_{CCANOPY} (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

Project Name: Mountain Valley Pipeline

Location: Franklin County

Sampling Date: 8/25/21 Project Site Before Project

Subclass for this SAR:

Intermittent Stream

Uppermost stratum present at this SAR: SAR number: S-G24

Shrub/Herb Strata

Functional Results Summary: Enter R

Enter Results in Section A of the Mitigation Sufficiency Calculator

Function	Functional Capacity Index
Hydrology	0.55
Biogeochemical Cycling	0.52
Habitat	0.40

Variable Measure and Subindex Summary:

Variable	Name	Average Measure	Subindex
V _{CCANOPY}	Percent canpoy over channel.	Not Used, <20%	Not Used
V _{EMBED}	Average embeddedness of channel.	3.27	0.92
V _{SUBSTRATE}	Median stream channel substrate particle size.	2.80	1.00
V _{BERO}	Total percent of eroded stream channel bank.	26.32	0.93
V_{LWD}	Number of down woody stems per 100 feet of stream.	0.00	0.00
V _{TDBH}	Average dbh of trees.	Not Used	Not Used
V _{SNAG}	Number of snags per 100 feet of stream.	0.00	0.10
V _{SSD}	Number of saplings and shrubs per 100 feet of stream.	421.05	1.00
V _{SRICH}	Riparian vegetation species richness.	0.00	0.00
V _{DETRITUS}	Average percent cover of leaves, sticks, etc.	17.50	0.21
V _{HERB}	Average percent cover of herbaceous vegetation.	73.33	0.98
V _{WLUSE}	Weighted Average of Runoff Score for Catchment.	0.61	0.64

	High-Gradient Headwater Streams in Appalachia Field Data Sheet and Calculator															
	Team: AW JB Latitude/UTM Northing: 37.126412 Project Name: Mountain Valley Pipeline Longitude/UTM Easting: -80.121398 Location: Franklin County Sampling Date: 8/25/21															
Pro	oject Name:	Mountain V	alley Pipelir	ne					_		3					
	Location:	Franklin Co	unty					San	npling Date:	8/25/21						
SA	AR Number:	S-G24	Reach	Length (ft):	19	Stream Ty	/pe: Inter	mittent Strea	m		_					
	Top Strata:	Sh	rub/Herb St	rata	(determine	d from perce	ent calculate	d in V _{CCANO}	PY)							
		Project Site				•	Before Proje	ct			•					
Sample		I-4 in stream channel Average percent cover over channel by tree and sapling canopy. Measure at no fewer than 10 roughly														
	V _{CCANOPY}	equidistant 20%, enter	points along at least one	g the stream value betw	. Measure een 0 and 1	only if tree/s	sapling cove Top Strata o	r is at least :			Not Used, <20%					
	0	cent cover r	neasureme	ns at each p	oint below.											
2	V_{EMBED}						at no fewer noving it, de				3.3					
	surface and area surrounding the particle that is covered by fine sediment, and enter the rating according to the following table. If the bed is an artificial surface, or composed of fine sediments, use a rating score of 1. If the bed is composed of bedrock, use a rating score of 5. [Embeddedness rating for gravel, cobble and boulder particles (rescaled from Platts, Megahan, and															
	Embeddedness rating for gravel, cobble and boulder particles (rescaled from Platts, Megahan, and Minshall 1983)															
	Rating Rating Description 5 <5 percent of surface covered, surrounded, or buried by fine sediment (or bedrock)															
		5 4					buried by fir d, or buried l			()						
		3					ed, or buried									
		2					ed, or buried	•								
	Lint the end	1			covered, su	rrounded, o	r buried by f	ine sedimer	nt (or artificia	al surface)						
	List the rati	ngs at each	point below	3	-											
	3	5	4	5	5 5											
	1	5	1	1	5											
3	Velibethate	Median stre	eam channe	substrate r	particle size.	Measure a	it no fewer t	nan 30 roug	hlv equidista	ant points						
		along the s	tream; use t	he same po	ints and par	ticles as use	ed in V _{EMBED}				2.80 in					
		as 0.0 in, s														
	8.00	0.08	0.08	2.40	10.00											
	3.00 0.08	6.20 5.10	2.80 0.08	1.80 3.00	2.40 7.20											
	0.00	0.10	0.00	0.00	1.20											
4		Tatal				E-ttht-	t-1		al and the section							
4	V_{BERO}		e total perce				otal number otal number otal number				26 %					
		may be ap	Left Bank:	0	ft		Right Bank:	5	ft							
Sample	Variables	5-9 within t	he entire ri	narian/huff	er zone adi	acent to the	stream ch	annel (25 fo	eet from ea	ch hank)						
5	V _{LWD}	Number of	down wood	y stems (at l	east 4 inche	es in diamet	er and 36 in	ches in leng	th) per 100	feet of						
				e number fr will be calcu	lated.		ouffer and w		·	e amount	0.0					
•	1/	A	h - f t /-				oody stems:			-4144						
6	V_{TDBH}	inches (10	cm) in diam	eter. Enter	tree DBHs in	n inches.	g cover is a	•		at least 4	Not Used					
		the stream	below:	ents of Indiv	ridual trees	(at least 4 in) within the		ich side of		•					
			Left Side					Right Side								
							_		_							
7	V	Number of	enage (at la	act A" dbb a	nd 36" toll)	ner 100 foot	of stream.	Enter numb	er of space	on each						
'	V _{SNAG}			ast 4" don a				Linei nuiilD	or or sirags	on eaul	0.0					
			Left Side:		0		Right Side:		0							
8	V_{SSD}	tree cover i	s <20%). E		of saplings		es dbh) per on each sid			asure only if e amount	421.1					
		, o. 100 it t	l off Cido	ouroural			Diabt Cida		-							

9 V _{SI}	SRICH	Group 1 in richness pe	er 100 feet a	na tne subii	HOOK WIII DO	ouloulutou i	ioni incac u	aıa.				
		Grou	ıp 1 = 1.0					Gro	up 2	2 (-1.0)		
Ac	cer rubrur	n		Magnolia ti	ripetala		Ailanthus a	Itissima		J	Lonicera ja	oonica
_ Ac	cer sacch	arum		Nyssa sylv	ratica		Albizia julib	rissin			Lonicera ta	tarica
– Ae	esculus fla	ava		Oxydendrun	n arboreum		Alliaria peti	olata			Lotus corni	culatus
_ As	simina tril	oba		Prunus ser	rotina		Alternanthe	ro			licaria	
_	etula alleg			Quercus a			philoxeroid				n vimineur	
_	_					1 1						
_	etula lenta 			Quercus co			Aster tatari				tomentos	
_	arya alba			Quercus in			Cerastium		7		uspidatur	
Ca	arya glabi	ra		Quercus p	rinus		Coronilla va	aria			Pueraria m	ontana
Ca	arya ovali	is		Quercus ru	ıbra		Elaeagnus u	mbellata		✓ I	Rosa multif	lora
Ca	arya ovata	а		Quercus ve	elutina		Lespedeza	bicolor			Sorghum h	alepense
Co	ornus flor	ida		Sassafras	albidum		Lespedeza	cuneata			Verbena br	asiliensi
Fa	agus gran	difolia		Tilia ameri	cana		Ligustrum ob	otusifolium	1			
Fra	axinus ar	mericana		Tsuga can	adensis		Ligustrum s	sinense				
— ∠ Liri	riodendron	tulipifera		Ulmus ame	ericana							
_		cuminata	_									
IVIC	agriolia a	carriiriata										
		1	Species in	Group 1				2		Species in	Group 2	
										one within	25 feet from	n each
							nch side of to material. Wo			//" diamete	r and <36"	
IU VD	DETRITUS						yer at each s		115 >	4 ulainete	ii aliu \30	17.50
				Side		<u> </u>		t Side			1	
		5	15	10		70	0	5				
11 V _H	HERB						asure only if					
II VH							there may b					73 %
II VH		-		s up inrougr	n 200% are a	accepted. E	Enter the per	cent cove	er o	r grouna ve	egetation at	
II VH		each subbi	OL.									
II VH		each subpl		Side		I	Right	t Side			1 '	
II VH		95		Side 70		5	Right	t Side] '	
mple Va	ariable 1	95 2 within the	Left 90 e entire cate	70 chment of t	the stream.							0.61
mple Va		95 2 within the	Left 90 e entire cate	70 chment of t						D (0.61
mple Va		95 2 within the	Left 90 e entire cate Average of R	70 chment of t		ned:				Runoff	% in Catch- ment	0.61 Runnir Percer (not >10
mple Va	WLUSE	95 2 within the Weighted A	Left 90 e entire cate Average of R	chment of the Runoff Score Use (Choose	e for watersh	ned:			•			Runnir Percer
mple Va	WLUSE	95 2 within the Weighted A	Left 90 e entire cate Average of R Land	chment of the Runoff Score Use (Choose cover)	e for watersh	ned:			• • • • • • • • • • • • • • • • • • •	Score 0.5	ment 26	Runnir Percer (not >10
mple Va	WLUSE	95 2 within the Weighted A	Left 90 e entire cate Average of R	chment of the Runoff Score Use (Choose cover)	e for watersh	ned:			*	0.5 1	ment	Runnir Percer (not >10
mple Va	wLUSE orest and n	95 2 within the Weighted A ative range (a ative ra	Left 90 e entire cate Average of R Land	Chment of the Runoff Score Use (Choose cover)	e for watersh	ned:			*	Score 0.5	ment 26	Runnir Percei (not >10
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate everage of R Land <50% ground >75% ground	chment of the control	e for watersh	ned:		80	* * * * * * * * * * * * * * * * * * *	0.5 1	ment 26 40	Runnir Percei (not >10 26 66
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:		80	ŀ	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:		80	•	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:		80	ŀ	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:		80	•	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and norest and n	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:		80	•	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and no prest and no prest and no prest and no previous appen space	95 2 within the Weighted A ative range (: ative range (: ative range (: pasture, law)	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	chment of the control	e for watersh	ned:	100	80	*	0.5 1 0	26 40 8	Runnir Percei (not >10 26 66 74
mple Va	orest and no prest and no prest and no prest and no previous appen space	95 2 within the Weighted A ative range (ative range (ative range (parking	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	Cover) cover) cover) cover) cover) cover) cover) cover) cover)	e for watersh se From Dro	p List)	100	80 80	▼ ▼ ▼	0.5 1 0 0.3	ment 26 40 8 26	Runnin Perce (not >10 26 66 74 100
mple Va	orest and norest and norest and nopervious appen space	95 2 within the Weighted A ative range (: ative r	Left 90 e entire cate Average of F Land <50% ground >75% ground g lots, roofs, d	Cover)	e for watersh se From Dro c) r > 75%	p List)	No poleted using	tes:	19 [0.5 1 0 0.3	ment 26 40 8 26 and Cover	Runnin Perce (not >10 26 66 74 100
mple Va 12 V _w For For Imp	orest and norest and n	95 2 within the Weighted A ative range (: ative ra	Left 90 e entire cate Average of R Land <50% ground >75% ground g lots, roofs, d ns, parks, etc.	Use (Choose Cover) I cover)	e for watersh se From Dro c) r >75% ver Analysis from Lands	p List) p was compat satellite	No pleted using imagery ar	tes:	▼	Score 0.5 1 0 0.3 National L	ment 26 40 8 26 and Cover	Runnin Perce (not >10 26 66 74 100
mple Va 12 V _w For For Implication Opp Varia	orest and no prest	95 2 within the Weighted A ative range (: ative range (: ative range (: parking (: pasture, law)) 6-G24 Value Not Used, <20%	Left 90 e entire cate Average of F Land <50% ground >75% ground glots, roofs, d ns, parks, etc.; VSI Not Used	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover	Runnii Perce (not >10 26 66 74 100
mple Va 12 Vw Foot Foot Imp Op Varia Vccan Vembe	orest and n orest and n orest and n opervious a pen space	95 2 within the Weighted A ative range (: ative ra	Left 90 e entire cate Average of R Land <50% ground p lots, roofs, d ns, parks, etc.	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
mple Va 12 V _w For For Implication Opp Varia	orest and n orest and n orest and n opervious a pen space	95 2 within the Weighted A ative range (: ative range (: ative range (: parking (: pasture, law)) 6-G24 Value Not Used, <20%	Left 90 e entire cate Average of F Land <50% ground >75% ground glots, roofs, d ns, parks, etc.; VSI Not Used	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
mple Va 12 V _w For For Implementation Varia Vccan Vembe Vsubs	orest and no prest	95 2 within the Weighted A ative range (ative range (ative range, law) (pasture, law) 4-G24 Value Not Used, <20% 3.3	Left 90 e entire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Mple Va 12 Vw Foot Foot Imp Op Varia Vccan Vembe Vsubs Vbero	orest and no prest	95 2 within the Weighted A within the Weighted A within the Weighted A with a with a wind a with a within the	Left 90 e entire cate Average of R Land <50% ground >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
mple Va 12 V _w For For Implementation Varia Vccan Vembe Vsubs	orest and no prest	95 2 within the Weighted A ative range (continue range)	Left 90 Pentire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Mple Va 12 Vw Foot Foot Imp Op Varia Vccan Vembe Vsubs Vbero	orest and n orest and n orest and n opervious a open space Sable NOPY ED STRATE	95 2 within the Weighted A within the Weighted A within the Weighted A with a with a wind a with a within the	Left 90 e entire cate Average of R Land <50% ground >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnin Perce (not >1) 26 66 74 100
Varia Vccan Vsubs Vsubs Vsubs Vsubs	orest and no prest	95 2 within the Weighted A ative range (control of the within the Weighted A ative range (control of the within the with	Left 90 Pentire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Varia Varia Vocan Varia Vsuss Vbero Vtobh Vsnag	orest and no prest	95 2 within the Weighted A within the Weighted A within the Weighted A with a with a wind a with a within the	Left 90 e entire cate Average of F Land <50% ground >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used 0.10	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Varia Vccan Vsubs Vsubs Vsubs Vsubs	orest and no prest	95 2 within the Weighted A ative range (control of the within the Weighted A ative range (control of the within the with	Left 90 Pentire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Varia Varia Vocan Varia Vsuss Vbero Vtobh Vsnag	orest and norest and n	95 2 within the Weighted A within the Weighted A within the Weighted A with a with a wind a with a within the	Left 90 e entire cate Average of F Land <50% ground >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used 0.10	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Waria Varia Vcan Vembe Vsubs Vbero Vtub Vsnag Vssch	orest and no prest	ative range (at	Left 90 e entire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used 0.10 1.00 0.00	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Waria Varia Vcan Varia Vcub Vsub Vsub Vsub Vsub Vsub Vsub Vsub Vs	MLUSE Porest and n porest an	95 2 within the Weighted A ative range (cative range (cative range) (cative rang	Left 90 Pentire cate Average of File Land Section 10 S	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnii Perce (not >10 26 66 74 100
Waria Varia Vcan Vembe Vsubs Vbero Vtub Vsnag Vssch	MLUSE Porest and n porest an	ative range (at	Left 90 e entire cate Average of R Land >75% ground glots, roofs, d ns, parks, etc. VSI Not Used 0.92 1.00 0.93 0.00 Not Used 0.10 1.00 0.00	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runnir Percer (not >10 26 66 74 100
Waria Varia Vcan Varia Vcub Vsub Vsub Vsub Vsub Vsub Vsub Vsub Vs	Sable NOPY ED STRATE D H RITUS	95 2 within the Weighted A ative range (cative range (cative range) (cative rang	Left 90 Pentire cate Average of File Land Section 10 S	To chment of the control of the cont	e for watersh se From Dro c) r > 75% rer Analysis rom Lands d boundari	p List) p was compat satellite es are bas	No pleted using imagery ar sed off of fice	tes: g the 20 d other	▼ ▼ ▼ 19 i supeate	0.5 1 0 0.3 National L pplementa	ment 26 40 8 26 and Cover rry datasets impacts.	Runni Perce (not >1 26 66 74 100

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME S-G24		LOCATION Franklin County	,
STATION # F	RIVERMILE	STREAM CLASS Intermitten	t
LAT <u>37.126412</u> L	ONG80.121398	RIVER BASIN Upper Roan	oke
STORET#		AGENCY VADEQ	
INVESTIGATORS JB, AV	N		
FORM COMPLETED BY	JB, AW	DATE 8/25/21 TIME 10:00 AM	REASON FOR SURVEY Baseline Assessment
WEATHER CONDITIONS	rain (shower 30 % ✓ %c	(steady rain)	Has there been a heavy rain in the last 7 days? Yes No Air Temperature 22 0 C Other
SITE LOCATION/MAP	Access Balbers SULSOC	Vegetation Signature Signature Signature Numbaleous Vegetation Numbaleous Vegetation Numbaleous Vegetation Numbaleous Numbaleou	Rence with AWAY SIL RENCING > 1 ASSESSOD AN UEG- AN UE

Notes: Only downstream water quality measurements were taken due to no assessable reach upstream.

Spring-fed

Mixture of origins
Other

Stream Type Coldwater

Catchment Area 0.11

✓Warmwater

 $\,\mathrm{km}^2$

Stream Subsystem
□ Perennial □ Intermittent □ Tidal

Stream Origin
Glacial
Non-glacial montane
Swamp and bog

STREAM CHARACTERIZATION

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Predom ✓ Fores ✓ Field/ ☐ Agric ☐ Resid	Pasture Industri Ultural Other	rcial al	Local Watershed NPS ☑ No evidence ☐ Son ☐ Obvious sources Local Watershed Erosi ☑ None ☐ Moderate	ne potential sources					
RIPARIA VEGETA (18 meter	TION		e the dominant type and s		ominant species present ☐ Grasses ☐ He outlens capensis	rbaceous					
INSTREA FEATURI		Estimat Samplin Area in Estimat	km² (m²x1000) ed Stream Depth Velocity 0.1 m	m m² km² m	Canopy Cover						
LARGE V DEBRIS	VOODY	LWD Density	of LWD NA n	n ² /km ² (LWD/	reach area)						
AQUATIO VEGETA		✓ Roote Floati Domina	licate the dominant type and record the dominant species present Rooted emergent Rooted submergent Rooted floating Rooted floating Attached Algae minant species present Impatient capensis rtion of the reach with aquatic vegetation 5 96								
WATER ((DS)	QUALITY	Specific Dissolve pH 8.78 Turbidi	cature 16.1 0 C Conductance 66.6 uS/cm ed Oxygen 9.77 mg/L ty N/A ctrument Used YSI	-		Other					
SEDIMEN SUBSTRA		Odors Norm Chem Other Oils Absen		Petroleum None	—————————————————————————————————————	□Paper fiber □Sand □Other □ h are not deeply embedded, k in color?					
INC		STRATE (COMPONENTS 00%)		ORGANIC SUBSTRATE C (does not necessarily add						
Substrate Type	Diamet	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area					
Bedrock			0	Detritus	sticks, wood, coarse plant	10					
Boulder	> 256 mm (10")		0		materials (CPOM)	10					
Cobble	64-256 mm (2.5	5"-10")	25	Muck-Mud							
Gravel	2-64 mm (0.1"-2	2.5")	25		(FPOM)						
Sand	0.06-2mm (gritt	y)	35	Marl	grey, shell fragments						
Silt	0.004-0.06 mm		15			l					
Clay	< 0.004 mm (sli	ok)	n	1							

Notes: Only downstream water quality measurements were taken due to no assessable reach upstream.

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-G24	LOCATION Franklin County						
STATION # RIVERMILE	STREAM CLASS Intermittent						
LAT <u>37.126412</u> LONG <u>-80.121398</u>	RIVER BASIN Upper Roanoke						
STORET#	AGENCY VADEQ						
INVESTIGATORS JB, AW							
FORM COMPLETED BY JB, AW	DATE 8/25/21 REASON FOR SURVEY TIME 10:00 AM PM Baseline Assessment						

	Habitat		Condition	Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.				
	SCORE 18	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.				
ted in	SCORE 11	20 19 18 17 16	10 9 8 7 6	5 4 3 2 1 0					
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).				
ıram	SCORE 16	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
P ₂	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.				
	SCORE 14	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	channel and mostly				
	SCORE 16	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				

Notes:

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)


	Habitat		Condition	n Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.				
	SCORE 19	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
ling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.				
amp	SCORE 18	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing downstream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.				
e ev	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
to b	SCORE 8	Right Bank 10 9	8 7 6	5 4 3	2 1 0				
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.				
	SCORE 6	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
	SCORE 6	Right Bank 10 9	8 7 6	5 4 3	2 1 0				
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.				
	SCORE 7	Left Bank 10 9	8 7 6	5 4 3	2 1 0				
	SCORE 8	Right Bank 10 9	8 7 6	5 4 3	2 1 0				

Total Score 157 Notes:

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-0	STREAM NAME S-G24								√ Fran	ıklin	Co	ount	У	_					_		
STATION #	R	IVE	RMI	LE_			STR	EAM C	CLASS	Int	ern	nitte	nt								
LAT 37.126412	_ L	ONO	}80.	12139	3		RIV	ER BAS	SIN U	Jpp	er F	Roar	noke	:							
STORET#							AGI	ENCY V	/ADEC	2											
INVESTIGATORS J	B, AV	N											I	LOT	NUMBER						
FORM COMPLETED	ЭBY	J	3,	Α	W		DAT TIM		5/21 00 AM				I	REASON FOR SURVEY Baseline Assessme							
HABITAT TYPES	✓	Cob	ble 2	5	%	tage of Sna	ags	habitat %	type p	Veg	geta	ted :	Ban (Silt a	KS 95	%	%					
SAMPLE	G	ear	used		D-fr	ame v	kick	-net			Ot	her									
COLLECTION											_					_					
	Н	ow v	vere	tne	samp	oles coll	ectea	<i>:</i> Ŀ	√wadi	ng		Ш	Iror	n bar	nk from boa	ıτ					
	✓	Cob	ble 4			r of jab Snaphytes_	ags	s taken	in ead	Veg	geta	itat ted her	Ban	ks	Sand)	_					
GENERAL	4	kic	:ks	in	riff	le ha	bita	t 5 (crav	fis	h	dis	SCS	ırde	ed.						
COMMENTS	Ι΄		,,,,			10 110	0110		Jiay		• •		,,,,								
Indicate estimated Dominant Periphyton	d abı	und	anco	e: (Not 2 3	Obser 4	rved,			are	2, 2	= C	ommon, 3= Abun		1	2	3	4	
Filamentous Algae					-	1 2		4					nve	rtebr	rates	-	1			4	
Macrophytes						1 2					ish						1		3		
FIELD OBSERV. Indicate estimated					0 =	Absen	t/Not	Obse							rganisms), 2 = Coo , 4 = Dominant (>				ıs)		
Porifera					4		opter						3						3		
Hydrozoa	0	1	2	3	4	Zygo	_		0		1	2	3	4	Ephemeroptera	0	1	2	3	4	
Platyhelminthes	0	1	2	3	4	Hem	_		0		1	2	3	4	Trichoptera	0	1	2	3	4	
Turbellaria	0	1	2	3	4	Cole	_		0		1	2	3	4	Other	0	1	2	3	4	
Hirudinea	0	1	2	3	4	Lepio	_	ra	0		1	2	3	4							
Oligochaeta	0	1	2	3	4	Sialio			0		1	2	3	4							
Isopoda Amphipoda	0	1	2	3	4 4	Cory Tipu		ae	0		1 1	2	3	4							
Decapoda	0	1	2	3	4	Emp		Δ.	0		1 1	2	3	4							
Gastropoda	0	1	2	3	4	Simu			0		1	2	3	4							
Bivalvia	0	1	2	3	4	Tabii			0		1	2	3	4							
21,41,14	Ü	1	_	J	•	Culc			0		1	2	3	4							

Mountain Valley Pipeline Data are not adjusted for subsampling



S-G24 Sample ID Collection Date 08-25-2021 ORDER GENUS/SPECIES COUNT Ephemeroptera Baetis sp. 4 2 29 Ephemeroptera Diphetor hageni Ephemeroptera Ephemerella sp. Ephemeroptera Eurylophella sp. Ephemeroptera Maccaffertium sp. Ephemeroptera Paraleptophlebia sp. 3 6 Ephemeroptera Stenacron sp. Plecoptera Leuctra sp. 23 Plecoptera Peltoperla sp. Plecoptera Perlodidae 3 4 Plecoptera Soyedina sp. Plecoptera Tallaperla sp. Trichoptera Diplectrona sp. 13 Trichoptera Polycentropodidae Trichoptera Rhyacophila sp. 10 Odonata Cordulegaster sp. Coleoptera Oulimnius sp. 10 Diptera-Chironomidae Larsia sp. Diptera-Chironomidae Micropsectra sp. Diptera-Chironomidae Neostempellina sp. 11 3 2 12 7 Diptera-Chironomidae Parachaetocladius sp. Diptera-Chironomidae Parametriocnemus sp. Diptera-Chironomidae Polypedilum sp. Diptera-Chironomidae Psilometriocnemus triannulatus 2 4 8 Diptera-Chironomidae Rheotanytarsus sp. Diptera-Chironomidae Stempellinella sp. 3 Diptera-Chironomidae Tanytarsus sp. Diptera-Chironomidae Thienemanniella sp. Diptera-Chironomidae Thienemannimyia gr. sp. 2 Diptera-Chironomidae Zavrelimyia sp. 15 Diptera Ceratopogoninae Diptera Dixa sp. 4 1 Diptera Limnophila sp. Diptera Pseudolimnophila sp. 2 Diptera Tabanidae Annelida tubificoid Naididae w/o cap setae 3 Bivalvia Sphaeriidae Gastropoda Lymnaeidae Other Organisms Nematoda

TOTAL

205

Mountain Valley Pipeline WV SCI Metrics



Sample II Collection Date	
WVSCI Metric Values	
Total taxa	22
EPT taxa	11
% EPT	50.2
% Chironomidae	28.8
% 2 Dominant	43.9
HBI	4.41
WVSCI Metric Scores	
Total taxa	104.8
EPT taxa	84.6
% EPT	54.7
% Chironomidae	71.9
% 2 Dominant	87.7
HBI	78.7
WVSCI Metric Scores	
Total taxa	100.0
EPT taxa	84.6
% EPT	54.7
% Chironomidae	71.9
% 2 Dominant	87.7
HBI	78.7
WVSCI Total Score	79.6

WVSCI Thresholds

Unimpaired = > 68.00 Gray Zone = 60.61 to 68.00 Impaired = <60.61

WOLMAN PEBBLE COUNT FORM

County: Franklin County Stream ID: S-G24

Stream Name: UNT to Green Creek

HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 8/25/2021 Surveyors: AW, JB Type: Representative

т 1	DADTICLE		LE COUNT	D (1.1	7F 4 3 "	T/ 0/	0/ 5
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cur
	Silt/Clay	< .062	S/C	A	11	10.89	10.89
	Very Fine	.062125		^	0	0.00	10.89
	Fine	.12525		•	0	0.00	10.89
	Medium	.255	SAND	•	0	0.00	10.89
	Coarse	.50-1.0		•	6	5.94	16.83
.0408	Very Coarse	1.0-2	7	•	12	11.88	28.71
.0816	Very Fine	2 -4		^	0	0.00	28.71
.1622	Fine	4 -5.7	1	•	1	0.99	29.70
.2231	Fine	5.7 - 8	1	•	2	1.98	31.68
.3144	Medium	8 -11.3	1	•	7	6.93	38.61
.4463	Medium	11.3 - 16	GRAVEL	•	1	0.99	39.60
.6389	Coarse	16 -22.6	1	•	1	0.99	40.59
.89 - 1.26	Coarse	22.6 - 32	1	•	0	0.00	40.59
1.26 - 1.77	Vry Coarse	32 - 45	1	•	2	1.98	42.57
1.77 -2.5	Vry Coarse	45 - 64	7	•	15	14.85	57.43
2.5 - 3.5	Small	64 - 90		•	19	18.81	76.24
3.5 - 5.0	Small	90 - 128	COBBLE	^	15	14.85	91.09
5.0 - 7.1	Large	128 - 180		•	6	5.94	97.03
7.1 - 10.1	Large	180 - 256		•	3	2.97	100.0
10.1 - 14.3	Small	256 - 362		•	0	0.00	100.0
14.3 - 20	Small	362 - 512		•	0	0.00	100.0
20 - 40	Medium	512 - 1024	BOULDER	^	0	0.00	100.0
40 - 80	Large	1024 -2048		^	0	0.00	100.0
80 - 160	Vry Large	2048 -4096	1	^	0	0.00	100.0
	Bedrock		BDRK	^	0	0.00	100.0
				Totals	101		

RIVERMORPH PARTICLE SUMMARY

River Name: UNT to Green Creek Reach Name: S-G24 Representative Survey Date: 08/25/2021

Size (mm)	тот #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	11 0 0 0 6 12 0 1 2 7 1 1 0 2 15 19 15 6 3 0 0 0	10.89 0.00 0.00 0.00 5.94 11.88 0.00 0.99 1.98 6.93 0.99 0.00 1.98 14.85 18.81 14.85 5.94 2.97 0.00 0.00 0.00 0.00 0.00 0.00	10.89 10.89 10.89 16.83 28.71 29.70 31.68 38.61 39.60 40.59 40.59 42.57 57.43 76.24 91.09 97.03 100.00 100.00 100.00 100.00 100.00 100.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.93 9.58 54.5 109.86 162.23 256 10.89 17.82 28.72 42.57 0		

Total Particles = 101.

		Strear	Unified S	tream Method	lology for use	in Virginia		' /		
Duelest #	Duciest Name (Any			able channels cla	HUC	ittent or perennia	SAR#	Impact	Impact	
Project #	Project Name (App Mountain Valley Pipelin		Locality Franklin	Class.				Length	Factor	
22865.06	Valley Pipeline,	LLC)	County	R4	03010101	8/25/2021	S-G24	75	1	
Name	ne(s) of Evaluator(s) Stream Name			ition				SAR Length	-	
	JB, AW	n Creek					7	5		
. Channel C	ondition: Assess the cross-sec	tion of the stream a	and prevailing con-	dition (erosion, ag Conditional Catego						
	Optimal	Subo	ptimal	Mar	ginal	Po	oor	Sev	rere	
Channel Condition	Very little incision or active erosion; 80 100% stable banks. Vegetative surfac protection or natural rock, prominent (80-100%). AND/OR Stable point bars bankfull benches are present. Access to their original floodplain or fully developed wide bankfull benches. Mic channel bars and transverse bars few Transient sediment deposition covers less than 10% of bottom.	e erosion or unproted of banks are s / Vegetative protec prominent (60 Depositional feat stability. The ban channels are well dinas access to be newly developed portions of the r sediment covers 1	ew areas of active cted banks. Majority table (60-80%), tion or natural rock -80%) AND/OR tures contribute to hkfull and low flow efined. Stream likely inkfull benches, or [floodplains along reach. Transient 0-40% of the stream tom.	Poor. Banks more or Poor due to Ic Erosion may be pr both banks. Vege 40-60% of banks. Survical or vertical or undid 140-60% Sediment transient, control Deposition that comay be forming/p shaped channels	less than Severe or stable than Severe over bank slopes. esent on 40-60% of tative protection on Streambanks may be ercut. AND/OR may be temporary / ibute instability, resent. AND/OR V-s have vegetative % of the banks and	laterally unstable further. Majority of vertical. Erosion pr banks. Vegetative on 20-40% of bank to prevent erosion. the stream is cow. Sediment is temp nature, and contri AND/OR V-shap vegetative protect	sised. Vertically / e. Likely to widen both banks are near esent on 60-80% of protection present s, and is insufficient AND/OR 60-80% of ored by sediment. orary / transient in buting to instability. sed channels have ion is present on > and stable sediment.	Streambed below av majority of banks Vegetative protecti than 20% of banks erosion. Obviou: present. Erosion/raw AND/OR Aggradin	stability. Severe eed within the banks. erage rooting depth, vertical/undercut. on present on less bank sloughing banks on 80-100%. g channel. Greater ib del is covered by uting to instability.	
Scores	3		.4	depositional featur to sta	res which contribute ability.	deposition is absent.		subterran	ean flow.	CI 2.40
	-			1		<u>'</u>		<u>'</u>		
. RIPARIAN	BUFFERS: Assess both bank	Cor	nditional Cate	gory		-	,	NOTES>>		
. RIPARIAN Riparian Buffers	Optimal Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas.	Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh >	nditional Cate	gory Mar	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrun, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.	-	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NOTES>>		
Riparian Buffers	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas.	Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	Low Suboptimal Riparian areas with tree stratum (dbh > 3 inches) present, with 30% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, hay production, sopen water. If present, tree stratum (dbh >3 inches) present, with 430% tree canopy cover with maintained understory. Low	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NOTES>>		
Riparian	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	nditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NOTES>>		
Riparian Buffers Scores Delineate ripa Determine squ	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5	Righ Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat	Low Suboptimal: Riparian areas with tree stratum (dbn > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Conduct th and width. Cal	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5	NOTES>>		
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas.	Righ Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat	Low Suboptimal: Riparian areas with tree stratum (dbn > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Conduct th and width. Cal	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trais, or other comparable conditions. Low 0.5	NOTES>>		
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank ware footage for each by measuring iparian Area and Score for each rip	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Condetth and width. Caline blocks below.	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5			
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank ware footage for each by measuring iparian Area and Score for each right % Riparian Area	Righ Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat yor estimating lenguarian category in the 20% 1.5	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Condetth and width. Caline blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5	CI= (Sum % RA * Sc		
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank ware footage for each by measuring iparian Area and Score for each right % Riparian Area > 50% Score > 0.85	Righ Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat yor estimating lenguarian category in the 20% 1.5	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Condetth and width. Caline blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5	CI= (Sum % RA * Sc Rt Bank CI >	0.90	CI 0.91
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank uare footage for each by measuring iparian Area and Score for each right % Riparian Area > 50% Score > 0.85 1 HABITAT: Varied substrate size	High Suboptimal: Riparian areas with tree stratum (dbb > 3 inches) present, with 30% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cate or estimating lenguarian category in the 20% 1.5	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Conduct and width. Calculate blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks 6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI >	0.90 0.91	CI 0.91
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank uare footage for each by measuring iparian Area and Score for each right % Riparian Area > 50% Score > 0.85 1 HABITAT: Varied substrate size	High Suboptimal: Riparian areas with tree stratum (dbb > 3 inches) present, with 30% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cate or estimating lenguarian category in the 20% 1.5	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calculate blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 Ition Scores using culators are provided to the control of the	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks 6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$	0.90 0.91	
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank uare footage for each by measuring iparian Area and Score for each right % Riparian Area > 50% Score > 0.85 1 HABITAT: Varied substrate size	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cate 3 or estimating lenguarian category in the 20% 1.5 20% 1.5	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calculate blocks below. 20% 0.5	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provided to the control of the	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks 6	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI >	0.90 0.91	
Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank . INSTREAN	Optimal Tree stratum (dbh > 3 inches) present with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Trian areas along each stream bank ware footage for each by measuring iparian Area and Score for each rip % Riparian Area > 50% Score > 0.85 4 Riparian Area > 60% Score > 0.85 1 HABITAT: Varied substrate size features.	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cate or estimating lenguarian category in the condition Cate or estimation category in the category in the category in the category in the category in	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Conductory (dense vegetation). 20% 0.5 20% 0.5 and depths; woody	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 High 0.85 ition Scores using culators are provided to the control of the cont	ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$	0.90 0.91 SAV; riffle/pool	

Stream Impact Assessment Form Page 2								
Project #	Project Name (Applicant)	Locality	Cowardin Class.	HUC	Date	SAR#	Impact Length	Impact Factor
22865.06 Mountain Valley Pipeline (Mountain Valley Pipeline, LLC) R4 03010101 8/25/2021 S-G24 75 1								

4. CHANNEL ALTERATION: Stream crossings, riprap, concrete, gabions, or concrete blocks, straightening of channel, channelization, embankments, spoil piles, constrictions, livestock

		NOTES>>					
	Negligible	Mir	nor	Mod	erate	Severe	
Channel Alteration	Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized.	Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	disrupted by any of the channel	is disrupted by any of the channel alterations listed in the parameter guidelines. If	60 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.		
Scores	1.5	1.3	1.1	0.9	0.7	0.5	

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

THE REACH CONDITION INDEX (RCI) >> 1.26

RCI= (Sum of all Cl's)/5, except if stream is ephemeral RCI = (Riparian Cl/2)

COMPENSATION REQUIREMENT (CR) >> 95

CI 1.50

CR = RCI X L_I X IF

INSERT PHOTOS:


(WSSI Photo Location "L:\22000s\22800\22865.06\Admin\05-ENVR\Field Data\Spread H\Field Forms\S-G24\Photos\2021-08-25_09-45-31.jpg")



 $Reach S-G24\ looking\ downstream\ within\ ROW.\ Assessment\ is\ limited\ to\ areas\ within\ the\ temporary\ ROW.$


DESCRIBE PROPOSED IMPACT	Γ:	
--------------------------	----	--

PROVIDED UNDER SEPARATE COVER



SURVEY NOTES:

- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on January 3, 2019.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).
- 4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.
- 5. All section views shown are left to right facing downstream.
- 6. Cross section B shot at location of pipe centerline (based on field stakes).





S-G24 PRE-CROSSING PHOTOS

01/03/2019



PHOTO TAKEN LOOKING DOWNSTREAM ON 01/03/2019





S-G25 PRE-CROSSING PHOTOS


01/03/2019





PHOTO TAKEN LOOKING DOWNSTREAM ON














NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

CROSS SECTION LEGEND

CROSS SECTION

EXISTING GRADE

H: 1"=10' V: 1"=5'

		tior		Prepared For: N	F	, in the contract of the contr	unty, nd Stud
POST-CROSSING PHOTOS		-Sec		pared		Eronblin Country	III CO I Wetla
PENDING CROSSING PHOTO TAKEN LOOKING		Profile and Cross-Section			Crossing S-G24 & S-G25 - UNT	ΓΙΔΙΙΚΙΙΙΙ ⊂ΟυΠΙΥ), Copyright © 2021 Wetland Stud	
THOTO TAKEN EOOKING							
PENDING CROSSING							
PHOTO TAKEN LOOKING		App. By					
		Rev. By					
PENDING CROSSING	REVISIONS						SCALE: AS NOTED
PHOTO TAKEN LOOKING	REVI	Description					ember, 2021
		No. Date					DATE: September, 2021
PENDING CROSSING	Hori	zontal D	atur	n: NA	.D 1983 I	JTM ZC)NE 17N
		ical Dati			AVD		
	MVP	ndary an I 2' C.I. T			Sourc	e:	

PHOTO TAKEN LOOKING

Draft

Sheet #

1 of 1

NAS

PFS MGE

Computer File Name:

2865_03 S-H MP 245-253 Sheets.dwg

:\Survey\22000s\22800\22865.03\Spread H Work Dwgs