## **Baseline Assessment – Stream Attributes**

# Reach S-G38 (Timber Mat Crossing) Ephemeral Spread H Montgomery County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	✓
RBP Physical Characteristics Form	✓
Water Quality Data	N/A – No water present
RBP Habitat Form	✓
RBP Benthic Form	N/A – No water present
Benthic Identification Sheet	N/A – No water present
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	✓
Longitudinal Profile and Cross Sections	✓



Photo Type: US VIEW
Location, Orientation, Photographer Initials: Upstream view of ROW looking SW, AO



Location, Orientation, Photographer Initials: Downstream view of ROW looking NE, AO

## Spread H Stream S-G38 (Timber Mat) Montgomery County



Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking SE, AO




Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking NW, AO

# Spread H Stream S-G38 (Timber Mat) Montgomery County



Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking NE, AO



Ver. 10-20-17

#### FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the <u>UPPERMOST STRATUM</u> of the plant community is determined based on the calculated value for V<sub>CCANOPY</sub> (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

**Project Name:** Mountain Valley Pipeline

Location: Montgomery County; Spread H

Sampling Date: 8/23/2021 Project Site Before Project

**Subclass for this SAR:** 

**Ephemeral Stream** 

Uppermost stratum present at this SAR: SAR number: S-G38

Shrub/Herb Strata

Functional Results Summary:

**Enter Results in Section A of the Mitigation Sufficiency Calculator** 

Function	Functional Capacity Index			
Hydrology	0.64			
Biogeochemical Cycling	0.59			
Habitat	0.36			

#### Variable Measure and Subindex Summary:

Variable	Name	Average Measure	Subindex
V <sub>CCANOPY</sub>	Percent canpoy over channel.	Not Used, <20%	Not Used
$V_{\sf EMBED}$	Average embeddedness of channel.	3.60	1.00
V <sub>SUBSTRATE</sub>	Median stream channel substrate particle size.	1.10	0.55
$V_{BERO}$	Total percent of eroded stream channel bank.	15.38	0.99
$V_{LWD}$	Number of down woody stems per 100 feet of stream.	0.00	0.00
$V_{TDBH}$	Average dbh of trees.	Not Used	Not Used
$V_{SNAG}$	Number of snags per 100 feet of stream.	0.00	0.10
$V_{SSD}$	Number of saplings and shrubs per 100 feet of stream.	30.77	0.47
V <sub>SRICH</sub>	Riparian vegetation species richness.	0.00	0.00
V <sub>DETRITUS</sub>	Average percent cover of leaves, sticks, etc.	26.25	0.32
V <sub>HERB</sub>	Average percent cover of herbaceous vegetation.	60.00	0.80
V <sub>WLUSE</sub>	Weighted Average of Runoff Score for Catchment.	1.00	1.00

Pr			Hign-C				ms in Apale	-	1		
Pr	Team:	MM, AO						Latitude/UT	M Northing:	37.267002	
• •	oject Name:						L	-	_	-80.312898	3
		Montgomer						San	npling Date:	8/23/2021	
Si	AR Number:			Length (ft):		Stream Ty		emeral Stream			•
	Top Strata:	Shi	rub/Herb Sti	rata	(determined	d from perce	ent calculate	ed in V <sub>CCANO</sub>	<sub>&gt;Y</sub> )		
Site	and Timing:	Project Site				•	Before Proje	ct			•
	e Variables				11 1	1 1				0 11	
1	V <sub>CCANOPY</sub>	V <sub>CCANOPY</sub> Average percent cover over channel by tree and sapling canopy. Measure at no fewer than 10 roughly equidistant points along the stream. Measure only if tree/sapling cover is at least 20%. (If less than 20%, enter at least one value between 0 and 19 to trigger Top Strata choice.)  List the percent cover measurements at each point below:									Not Use <20%
	0										1
2	$V_{EMBED}$	along the si surface and according to rating score	tream. Seled area surro to the following of the following	ect a particle unding the p ing table. If a bed is com	from the becarticle that the bed is a posed of be	ed. Before r is covered l in artificial s drock, use a	at no fewer moving it, de by fine sedir surface, or co a rating scor	etermine the nent, and er composed of re of 5.	percentage iter the ratir fine sedime	of the ng ents, use a	3.6
		Embeddedr Minshall 19	•	for gravel, c	obble and b	oulder parti	cles (rescale	ed from Plat	ts, Megahar	n, and	Measur at leas
		Rating 5	Rating Des		overed, sur	rounded, or	buried by fi	ne sediment	(or bedroc	<b>(</b> )	30 point
		4	5 to 25 per	cent of surfa	ce covered	, surrounde	d, or buried	by fine sedii	ment	-7	
		3 2					ed, or buried ed, or buried				-
		1					or buried by			al surface)	1
	List the rati	ngs at each	•			·			,	,	•
	3	4	5	4	3	5	4	3	5	5	
	3	4	2	3	1						
	asphalt or o	cle size in in concrete as	ches to the 0.0 in, sand	nearest 0.1 or finer par	inch at each	n point below	,	should be co			1.10 in
	3.50	0.50	1.15	0.80	3.10	0.08	0.30	1.30	0.60	1.40	
	1.05	4.80	1.10	1.20	0.10						
	\/	Tatal name	nt of one do	d -4	anal hank	Custom the a to	-1-1	-f ft -f -u		b	
4	V <sub>BERO</sub>	•	e total perce	entage will b		d If both ba	otal number nks are ero Right Bank:	ded, total er			15 %
amnle	V <sub>LWD</sub>		down woody	y stems (at l	er zone adja east 4 inche			-			
5								_			0.0
				e number fr will be calcu	ılated.	e 50'-wide k	er and 36 in ouffer and w oody stems:	ithin the cha		ne amount	0.0
	$V_{TDBH}$	per 100 fee Average db	t of stream oh of trees (r	will be calcumeasure on	lated.  Number of	e 50'-wide k f downed wo <sub>Y</sub> tree/saplir	ouffer and w	ithin the cha	innel, and th		
5	V <sub>ТДВН</sub>	Average db	oh of trees (i cm) in diam	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and woody stems:	thin the cha	onnel, and the		0.0
5	V <sub>товн</sub>	Average db inches (10 db)	oh of trees (i cm) in diam	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	thin the cha	onnel, and the		
5	V <sub>ТДВН</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
5	V <sub>TDBH</sub>	Average db inches (10 db)	ot of stream on of trees (r cm) in diam n measurem below:	will be calcu measure on eter. Enter	Number of V <sub>CCANOP</sub> tree DBHs in	e 50'-wide to the following of the follo	ouffer and w body stems: ng cover is a	t least 20%)	onnel, and the		
6		Average db inches (10 db the stream	t of stream th of trees (it cm) in diam measurem below: Left Side	measure onleter. Enter	Ilated. Number of ly if V <sub>CCANOP</sub> tree DBHs in vidual trees	e 50'-wide he for downed work tree/sapling inches. (at least 4 in	pouffer and woody stems: ag cover is any within the	t least 20%) buffer on ea	Trees are	e at least 4	
5	V <sub>TDBH</sub>	Average db inches (10 List the dbh the stream	t of stream th of trees (incom) in diam in measurem below: Left Side	measure onleter. Enter	Ilated. Number of ly if V <sub>CCANOP</sub> tree DBHs in vidual trees	e 50'-wide he for downed work tree/saplir in inches. (at least 4 in the formal inches)	pouffer and woody stems:  ng cover is and  n) within the  t of stream.	t least 20%) buffer on ea	Trees are	e at least 4	
6		Average db inches (10 List the dbh the stream	t of stream th of trees (incom) in diam in measurem below: Left Side  snags (at lestream, and	measure onleter. Enter ents of indiverse of indiverse as 4" dbh a the amount	Ilated. Number of ly if V <sub>CCANOP</sub> tree DBHs in vidual trees and a second and a second a secon	e 50'-wide he for downed work tree/saplir in inches. (at least 4 in the formal inches)	pouffer and woody stems: ag cover is and in) within the tof stream. culated.	t least 20%) buffer on ea	Trees are ach side of ach side	e at least 4	Not Use
5	V <sub>SNAG</sub>	Average db inches (10 db inches) (10	t of stream th of trees (rom) in diam measurem below: Left Side snags (at lestream, and	measure onleter. Enter ents of indiv	Ilated. Number of ly if V <sub>CCANOP</sub> tree DBHs in ridual trees  and 36" tall) t per 100 fee	e 50'-wide before downed we want to the following the foll	pouffer and woody stems: ag cover is and an) within the tof stream. culated. Right Side:	t least 20%) buffer on ea	annel, and the	e at least 4	Not Use
5		Average db inches (10 db List the dbh the stream  Number of side of the stream	ch of trees (norm) in diam measurem below:  Left Side  snags (at lestream, and Left Side: saplings and	measure onleter. Enter ents of indiv	Ilated. Number of y if V <sub>CCANOP</sub> tree DBHs in yidual trees and 36" tall) t per 100 fee	e 50'-wide before downed we way tree/saplir in inches. (at least 4 in the same of the same	pouffer and woody stems: ag cover is and in) within the tof stream. culated.	t least 20%) buffer on ea Right Side  Enter numb	annel, and the control of the contro	on each	Not Use

9	V <sub>SRICH</sub>	Group 1 in	the tallest s	tratum. Chec	k all exotic	) feet of stream reach. Check all species present from tic and invasive species present in all strata. Species e calculated from these data.							
		•	er 100 feet a p 1 = 1.0	ind the subinc	dex will be	calculateo							
	Acer rubrui		<u>ρι- 1.5</u>	Magnolia trip	petala		Ailanthus a	-	2 (-1.0)	Lonicera ja	ponica		
	Acer sacch	narum		Nyssa sylvat			Albizia julibi	rissin		Lonicera ta			
	Aesculus fl	'ava		Oxydendrum a	arboreum		Alliaria petio	olata		Lotus corni	culatus		
	Asimina tril	oba		Prunus serot	tina		Alternanthe	ra		Lythrum sa	licaria		
	Betula alleg	haniensis		Quercus alba	а		philoxeroide	es		Microstegium	vimineum		
	Betula lent	а		Quercus coc	ccinea		Aster tatari	cus		tomentosa			
	Carya alba			Quercus imb	oricaria		Cerastium t	fontanum		uspidatum			
	Carya glab			Quercus prin	านร		Coronilla va	aria		ontana			
	Carya oval					V	Elaeagnus ur		<u> </u>	Rosa multif			
	Carya ovata			Quercus velu			Lespedeza 			Sorghum h			
	<del>-</del>			Sassafras alı Tilia america			Lespedeza		Ш	Verbena br	asiliensis		
	Fraxinus ai			Tsuga canac			Ligustrum ob						
	Liriodendron			Ulmus ameri		Ш	Ligustrum	aniense					
	Magnolia a			ominao amon	Journa								
		0	Species in	Group 1				2	Species in	Group 2			
Sample	e Variables	10-11 withi	n at least 8	subplots (40	0" x 40". o	r 1m x 1m)	) in the ripar	ian/buffer z	one within	25 feet from	n each		
_				ed roughly ed		-	-		one within	20 1001 1101	ii cacii		
10	$V_{\text{DETRITUS}}$	• .		of leaves, sti r the percent		•		•	<4" diamete	er and <36"	26.25 %		
		long are inc		Side	cover or un	e detrital la		Side		1			
		30	Lon	Oluc		35	ragin	Olde					
		15				25							
11	$V_{HERB}$			over of herbact t least 4" dbh									
		vegetation	percentage:	s up through 2							60 %		
		each subpl		Cido			Diabt	Cido		1			
	-		Left Side Right Side										
	65					40							
		85				40 50							
Sample	e Variable 1	85	e entire cate	chment of the	e stream.								
Sample 12		85 2 within the		chment of the		50				<u> </u>			
	e Variable 1	85 2 within the				50					1.00		
		85 2 within the	Average of F	Runoff Score f	for watersh	50 ned:			Runoff	% in Catch	Running		
		85 2 within the	Average of F		for watersh	50 ned:			Runoff Score	% in Catch- ment			
	V <sub>WLUSE</sub>	85 2 within the	Average of F Land	Runoff Score f	for watersh	50 ned:					Running Percent		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		<b>*</b>	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:			Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		<b>*</b>	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		<b>*</b>	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		• •	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		<b>*</b>	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		• •	Score	ment	Running Percent (not >100)		
	V <sub>WLUSE</sub>	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:		• • • • • • • • • • • • • • • • • • •	Score	ment	Running Percent (not >100)		
	Forest and n	85  2 within the Weighted A	Average of F Land	Runoff Score f	for watersh	50 ned:	Not	* * * * * * * * * * * * * * * * * * *	Score	ment	Running Percent (not >100)		
12	Forest and n	85  2 within the Weighted A	Average of F Land	Cover)  Land Cover	From Drop	bed:  Description:	pleted using	• • • • • • • • • • • • • • • • • • •	Score 1 National La	ment 100 and Cover [	Running Percent (not >100)  100  Database		
12 V	Forest and n	85  2 within the Weighted Anative range (> S-G38  Value Not Used,	Land 75% ground	Cover)  Land Cover (NLCD), fro	From Drop	was compat satellite	pleted using imagery an	tes:	Score  1  National Lapplemental	ment  100  and Cover Iny datasets.	Running Percent (not >100)  100  Database		
12 V <sub>C</sub>	Forest and n	85  2 within the Weighted A sative range (> S-G38  Value  Not Used, <20%	Land 75% ground VSI Not Used	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
12 V <sub>C</sub> V <sub>E</sub>	Forest and n	85  2 within the Weighted A weighted A water range (> S-G38  Value  Not Used, <20%  3.6	VSI Not Used 1.00	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
12 V V V V V V V V V V V V V V V V V V V	Forest and n  Forest and n  S  ariable  CANOPY  MBED  UBSTRATE	85  2 within the Weighted A sative range (> S-G38  Value  Not Used, <20%	Land 75% ground VSI Not Used	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
12 V V V V V V V V V V V V V V V V V V V	Forest and n	85  2 within the Weighted A with a stive range (> S-G38  Value  Not Used, <20%  3.6	VSI Not Used 1.00	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
12 V V V V V V V V V V V V V V V V V V V	Forest and n  Forest and n  S  ariable  CANOPY  MBED  UBSTRATE  ERO	85  2 within the Weighted A  ative range (>  S-G38  Value  Not Used, <20%  3.6  1.10 in	VSI Not Used 1.00 0.55	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n  Forest and n  S  ariable  CANOPY  MBED  UBSTRATE  ERO	85  2 within the Weighted A weighted A wative range (> S-G38  Value  Not Used, <20%  3.6  1.10 in  15 %	VSI Not Used 1.00 0.55 0.99	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n S ariable CANOPY MBED UBSTRATE ERO WD	85  2 within the Weighted A within the Weighted A within the Meighted A with a with a wind a	VSI Not Used 1.00 0.55 0.99 0.00	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n S ariable CANOPY MBED UBSTRATE ERO WD DBH NAG	85  2 within the Weighted A within the Weighted A within the Weighted A with a wind a	VSI Not Used 1.00 0.55 0.99 0.00 Not Used 0.10	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n Sariable CANOPY MBED UBSTRATE ERO WD DBH NAG SD	85  2 within the Weighted A with a wind a wind with a wind with a wind a	VSI Not Used 1.00 0.55 0.99 0.00 Not Used 0.10 0.47	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n Sariable CANOPY MBED UBSTRATE ERO WD DBH NAG SD RICH	85  2 within the Weighted A within the Weighted A within the A with a wind a wi	VSI Not Used 1.00 0.55 0.99 0.00 Not Used 0.10 0.47 0.00	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n Forest and n S S S S S S S S S S S S S S S S S S S	85  2 within the Weighted A  ative range (>  3-G38  Value  Not Used, <20%  3.6  1.10 in  15 %  0.0  Not Used  0.0  30.8  0.00  26.3 %	VSI Not Used 1.00 0.55 0.99 0.00 Not Used 0.10 0.47 0.00 0.32	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		
V	Forest and n Forest and n Sariable CANOPY MBED UBSTRATE ERO WD DBH NAG SD RICH	85  2 within the Weighted A within the Weighted A within the A with a wind a wi	VSI Not Used 1.00 0.55 0.99 0.00 Not Used 0.10 0.47 0.00	Cover)  Land Cover (NLCD), fro Watershed	From Drop r Analysis om Landsa	was compat satellite es are bas	pleted using imagery and sed off of fie	tes: the 2019 d other sup	Score  1  National Lapplemental ed stream	ment  100  and Cover Iny datasets. impacts.	Running Percent (not >100) 100		

# PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

	I				
STREAM NAME S-G38	LOCATION Montgomery County				
STATION # RIVERMILE	STREAM CLASS Ephemeral				
LAT <u>37.267002</u> LONG <u>-80.312898</u>	RIVER BASIN Upper Roand	oke			
STORET#	AGENCY VADEQ				
INVESTIGATORS AO, MM					
FORM COMPLETED BY AO, MM	DATE 8/23/2021 TIME 1:30PM	REASON FOR SURVEY Baseline Assessment			
	T	Los thous been a bearry pain in the last 7 days?			

WEATHER CONDITIONS	Now  Past 24 hours  Yes No  Air Temperature 29 ° C  Other  Other
SITE LOCATION/MAP	Draw a map of the site and indicate the areas sampled (or attach a photograph)  Roman School
STREAM CHARACTERIZATION	Stream Subsystem

Notes: No water present.

# PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Predom  ✓ Fores  ✓ Field/  ☐ Agric  ☐ Resid	Pasture Industri	ercial ial	Local Watershed NPS  ☑ No evidence ☐ Son ☐ Obvious sources  Local Watershed Erosi ☑ None ☐ Moderate	ne potential sources				
RIPARIA VEGETA (18 meter	TION		e the dominant type and S		ominant species present ☐ Grasses	rbaceous				
INSTREA FEATURI		Estimat Samplin Area in Estimat	ted Stream Depth NA	m m² km²	High Water Mark	✓ Partly open       ☐ Partly shaded       ☐ Shaded         High Water Mark       ○.10       m         Proportion of Reach Represented by Stream Morphology Types       Riffle MA       % Run MA         Pool NA       %       Run NA       %         Channelized       ☐ Yes       ☑ No				
LARGE V DEBRIS	VOODY	LWD Density	of LWD NA n	n²/km² ( <b>LWD</b> /	reach area)					
AQUATIO VEGETA		Roote Floati	e the dominant type and demergent RA A A A A A A The species present MA of the reach with aquations and the reach with aquatical RA A A A A A A A A A A A A A A A A A A	ooted submerge ttached Algae		□Free floating				
WATER (	QUALITY	Specific  Dissolve  pH N/A  Turbidi	cature NA 0 C c Conductance NA ed Oxygen NA city NA ctrument Used NA	-		Other				
SEDIMEN SUBSTRA		Odors Norm Chem Other Oils Absen		Petroleum None	— Lρoking at stones whic are the undersides blace	☐Paper fiber ☐Sand ]Other				
INC		STRATE of	COMPONENTS		ORGANIC SUBSTRATE C (does not necessarily add					
Substrate Type	Diamet	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area				
Bedrock			10	Detritus	sticks, wood, coarse plant	0				
Boulder	> 256 mm (10")	)	0		materials (CPOM)	, and the second				
Cobble	64-256 mm (2.5	5"-10")	20	Muck-Mud	black, very fine organic (FPOM)	0				
Gravel	2-64 mm (0.1"-2	2.5")	60		(11 Olv1)	<u> </u>				
Sand	0.06-2mm (gritt	y)	0	Marl	grey, shell fragments	0				
Silt	0.004-0.06 mm		5							
Clov	< 0.004 mm (cli	ok)	5							

Notes: No water present.

#### HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-G38	LOCATION Montgomery County					
STATION # RIVERMILE	STREAM CLASS Ephemeral					
LAT <u>37.267002</u> LONG <u>-80.312898</u>	RIVER BASIN Upper Roanoke					
STORET#	AGENCY VADEQ					
INVESTIGATORS AO, MM						
FORM COMPLETED BY AO, MM	DATE 8/23/2021 TIME 1:30PM AM PM REASON FOR SURVEY Baseline Assessment					

	Habitat		Condition	Category					
	Parameter	Optimal	Suboptimal	Marginal	Poor				
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.				
	SCORE 0	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.				
ted i	SCORE 15	20 19 18 17 16	5 4 3 2 1 0						
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).				
ıram	SCORE 0	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
P <sub>2</sub>	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.				
	SCORE 14	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.				
	score 0	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0				

Notes: No water present.

#### HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	ı Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor			
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.			
	SCORE 20	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
ling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.			
amp	score 0	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank)  Note: determine left or right side by facing deventram.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.			
eva	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
to be	SCORE 9	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.			
	SCORE 7	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
	SCORE 6	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.			
	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
	SCORE 9	Right Bank 10 9	8 7 6	5 4 3	2 1 0			

100 Notes: No water water present. No samples taken.

A-8

#### BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-G38					L	LOCATION Montgomery County													
STATION #	R	IVE	RM	ILE_		S	STREAM CLASS Ephemeral												
LAT 37.267002	_ L	ONO	j -80.	31289	8	R	RIVER BASIN Upper Roanoke												
STORET#						A	AGENCY VADEQ												
INVESTIGATORS A	O, N	IM										I	LOT	NUMBER					
FORM COMPLETED	FORM COMPLETED BY AO, MM					/ D	ATE .	8/23/2021 1:30PM	_			I	REAS	SON FOR SURVEY Ba	aselin	е А	sses	ssm	ent
HABITAT TYPES	∥∟	Cob	ble_		_%	tage of eac Snags phytes	9	itat typ	$\Box$ V	eget	t ated	Bani	ks	%	%				
SAMPLE	G	ear	used	Г	D-fr	ame $\square$ ki	ck-net			По	ther								
COLLECTION	Gear used D-frame kick-net Other																		
	Н	How were the samples collected? ☐ wading ☐ from bank ☐ from boat																	
		Cob	ble			r of jabs/k Snags phytes			$\Box V$	eget		Ban		Sand )	_				
GENERAL	N	O 1/	wat	er	nre	sent. E	Rent	hice	nΩ	t c	مااد	2Ct	ed						
COMMENTS	'`	O V	vai	Ci	pic	SCIII. L	JCIIII	11103	110	·	Onc	,,,	cu.						
Indicate estimated Dominant  Periphyton Filamentous Algae		und	anc	e:	0		3 4			Sliı	nes			ommon, 3= Abuno	0	1 1	2		4 4
Macrophytes							3 4			Fis		1100	rtebi	ales	-	1	_	3	
	l ab	und	anc	e:	0 = org	Absent/N anisms), 3	lot Ob 3= Ab	serve	ıt (>	>10	org	anis	sms)	rganisms), 2 = Coi , 4 = Dominant (>:	50 oı	gar	nism		
Porifera	0	1	2	3	4	_			0	1	2			Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygopte			0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemipte			0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleopt			0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidop			0	1	2	3	4						
Oligochaeta Isopoda	0	1	2	3	4 4	Sialidae Corydal			0	1 1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulida			0	1	2	3	4						
Decapoda Decapoda	0	1	2	3	4	Empidio			0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simulii			0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinid			0	1	2	3	4						
		_		_		Culcida			0	1	2	3	4						

#### WOLMAN PEBBLE COUNT FORM

County: Montgomery County
Stream Name: UNT to North Fork Roanoke River S-G38 Stream ID:

HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 8/23/2021 Surveyors: AO, MM Type: Representa Representative

		PEBBI	LE COUNT				
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	<b>^</b>	4	4.00	4.00
	Very Fine	.062125		<b>4</b>	0	0.00	4.00
	Fine	.12525		<b>4</b>	0	0.00	4.00
	Medium	.255	SAND	<b>4</b>	0	0.00	4.00
	Coarse	.50-1.0		<b>4</b>	0	0.00	4.00
.0408	Very Coarse	1.0-2		<b>4</b>	0	0.00	4.00
.0816	Very Fine	2 -4		<b>4</b>	2	2.00	6.00
.1622	Fine	4 -5.7		•	4	4.00	10.00
.2231	Fine	5.7 - 8		<b>4</b>	4	4.00	14.00
.3144	Medium	8 -11.3	1	<b>4</b>	4	4.00	18.00
.4463	Medium	11.3 - 16	GRAVEL	<b>^</b>	5	5.00	23.00
.6389	Coarse	16 -22.6	1	<b>4</b>	13	13.00	36.00
.89 - 1.26	Coarse	22.6 - 32	1	•	16	16.00	52.00
1.26 - 1.77	Vry Coarse	32 - 45	1	<b>^</b>	13	13.00	65.00
1.77 -2.5	Vry Coarse	45 - 64		<b>A</b>	8	8.00	73.00
2.5 - 3.5	Small	64 - 90		<b>4</b>	10	10.00	83.00
3.5 - 5.0	Small	90 - 128	COBBLE	<b>4</b>	3	3.00	86.00
5.0 - 7.1	Large	128 - 180	CORRLE	<b>A</b>	5	5.00	91.00
7.1 - 10.1	Large	180 - 256	1	<b>4</b>	0	0.00	91.00
10.1 - 14.3	Small	256 - 362		<b>^</b>	0	0.00	91.00
14.3 - 20	Small	362 - 512		<b>^</b>	0	0.00	91.00
20 - 40	Medium	512 - 1024	BOULDER	<b>^</b>	0	0.00	91.00
40 - 80	Large	1024 -2048		<b>^</b>	0	0.00	91.00
80 - 160	Vry Large	2048 -4096		<b>^</b>	0	0.00	91.00
	Bedrock		BDRK	<b>^</b>	9	9.00	100.00
				Totals	100		
	Total Tally:						

#### RIVERMORPH PARTICLE SUMMARY

UNT to North Fork Roanoke River

S-G38

River Name: Reach Name: Sample Name: Sample Name: Representative Survey Date: 08/23/2021

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	4 0 0 0 0 0 0 2 4 4 4 4 5 13 16 13 8 10 3 5 0 0 0	4.00 0.00 0.00 0.00 0.00 0.00 2.00 4.00 4.00 4.00 4.00 5.00 13.00 16.00 13.00 8.00 10.00 3.00 5.00 0.00 0.00 0.00 0.00	4.00 4.00 4.00 4.00 4.00 6.00 10.00 14.00 18.00 23.00 36.00 52.00 65.00 73.00 83.00 86.00 91.00 91.00 91.00 91.00 91.00 91.00 91.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	9.65 22.09 30.83 102.67 Bedrock Bedrock 4 0 69 18 0		

Total Particles = 100.


#### **Ephemeral Stream Assessment Form (Form 1a)** Unified Stream Methodology for use in Virginia For use in ephemeral streams Project # **Project Name** Locality Class. Length Factor Mountain Valley Pipeline (Mountain Montgomery 22865.06 03010101 8/23/2021 S-G38 Valley Pipeline, LLC) Name(s) of Evaluator(s) Stream Name and Information SAR Length 33 Unnamed Tributary to North Fork Roanoke River 2. RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) Conditional Category Optimal Low Marginal: Non-maintained, dense herbaceou /egetation, riparia areas lacking shrub and tree stratum, hay High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% Low Suboptima Riparian areas with tree stratun (dbh > 3 inches High Marginal: Non-maintained lense herbaceou no-till cropland; actively grazed pasture, sparsely vegetated nonlense herbaceou vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. present, with Free stratum (dbh > 3 inches) present with > 60% tree canopy cover and ar non-maintained understory. Wetlands spoil lands Riparian to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. stratum, hay oroduction, ponds open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained cover and a maintained understory. Recent cutove (dense vegetation). nuded surface Buffers High Low High Low High Low Condition 1.5 1.2 0.85 0.75 0.6 0.5 an areas along each stream bank into Condition Categories and Condition Scores using the 2. Determine square footage for each by measuring or estimating length and width. Calculators are provided for of % Riparian Blocks equal 100 8. Enter the % Riparian Area and Score for each riparian category in the blocks below % Riparian Area> 70% 10% 10% 5% Right Bank 0.5 0.6 0.75 1.5 0.85 CI= (Sum % RA \* Sc % Riparian Area> 75% 25% Rt Bank CI > Left Bank 0.74 0.85 0.5 Lt Bank CI > REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH THE REACH CONDITION INDEX (RCI) >> COMPENSATION REQUIREMENT (CR) >>

#### INSERT PHOTOS:


(WSSI Photo Location L:\(\text{L2000s}\)\(\text{L22000s}\)\(\text{L2800}\)\(\text{L2800}\)\(\text{L2800}\)\(\text{L2800}\)\(\text{L2800}\)\(\text{L380}\)\(\text{L41}\)\(

DESCRIBE PROPOSED IMPACT:

<u>S-G38</u>



CL STAKEOUT POINTS: S-G38 CROSS SECTION A (PIPE CL)							
	PF	POST-CROSSING					
PT. LOC.	NORTHING	EASTING	ELEV	VERT.	HORZ.		
	NORTHING	EASTING	ELEV	DIFF.	DIFF.		
TS-L	13532496.930	1840294.531	1620.583				
BS-L	13532496.070	1840295.250	1620.758				
THW	13532495.851	1840295.483	1620.650				
BS-R	13532495.160	1840295.939	1620.678				
TS-R	13532494.020	1840296.891	1621.095				



#### SURVEY NOTES:

LEGEND

EXISTING SURVEY-LOCATED THALWEG


STUDY AREA (EASEMENT)

-1900- EXISTING MAJOR CONTOUR

--1904--- EXISTING MINOR CONTOUR

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON AUGUST 23, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

# S-G38 BASELINE CROSS-SECTION A PIPE CL



DISTANCE ALONG CROSS-SECTION (FT)

CROSS SECTION LEGEND — EXISTING GRADE

CROSS SECTION H: 1"=10' V: 1"=5' SCALE:

NOTE: ALL SECTIONS VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

### PRE-CROSSING PHOTOS



PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS ON 08/23/2021



PHOTO TAKEN LOOKING UPSTREAM FROM DOWNSTREAM IMPACT LIMITS ON 08/23/21

POST-CROSSING PHOTOS

PENDING CROSSING

PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS

PENDING CROSSING

PHOTO TAKEN LOOKING UPSTREAM FROM

DOWNSTREAM IMPACT LIMITS

CAD File No. Checked



TO .38)

Drawing No.