### **Baseline Assessment - Stream Attributes**

# Reach S-IJ83 (Timber Mat Crossing) Intermittent Spread H Roanoke County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	N/A - Headwater stream <4% slope
RBP Physical Characteristics Form	✓
Water Quality Data	N/A – No flow
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A – No flow
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	<b>√</b>
Longitudinal Profile and Cross Sections	✓



Photo Type: DS VIEW
Location, Orientation, Photographer Initials: Downstream view of ROW looking S, TC





Photo Type: US VIEW Location, Orientation, Photographer Initials: Upstream view of ROW looking N, TC



Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking SW, TC



Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking E, TC

## Spread H Stream S-IJ83 (Timber Mat) Roanoke County




Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking SW, TC

USACE FILE NO./ Project Name: (v2.1, Sept 2015)		Mountain \	/alley Pipeline		COORDINATES: imal Degrees)	Lat.	37.169211	Lon.	-80.138258	WEATHER:		Cloudy	DATE:		Septemb	ber 7, 2021
IMPACT STREAM/SITE ID			S	-IJ83			MITIGATION STREAM CLASS						Comments:			
(watershed size (acreage),	unaltered or impairn	ments)					(watershed size (acrea	ge), unaltered or impo	airments)							
STREAM IMPACT LENGTH:	148	FORM OF MITIGATION:	RESTORATION (Levels I-III)		OORDINATES: imal Degrees)	Lat.		Lon.		PRECIPITATION PAST 48 HRS:	0.0	05"; 9/6/2021	Mitigation Leng	th:		
Column No. 1- Impact Existing	Condition (Deb	it)	Column No. 2- Mitigation Existing	Condition - Base	line (Credit)		Column No. 3- Mitigation F Post Completi	Projected at Five on (Credit)	Years	Column No. 4- Mitigation Pro Post Completion	jected at Ten Ye (Credit)	ars	Column No. 5- Mitig	ation Projected a	at Maturity (	Credit)
Stream Classification:	Intermi	ittent	Stream Classification:				Stream Classification:		0	Stream Classification:		0	Stream Classification:			0
Percent Stream Channel SI	оре	1.5	Percent Stream Channel S	lope			Percent Stream Channel	Slope	0	Percent Stream Channel S	lope	0	Percent Stream	Channel Slop	9	0
HGM Score (attach d	ata forms):		HGM Score (attach	data forms):			HGM Score (attac	h data forms):		HGM Score (attach d	ata forms):		HGM Sco	re (attach data	forms):	
		Average			Average				Average			Average				Average
Hydrology			Hydrology				Hydrology			Hydrology			Hydrology			4
Biogeochemical Cycling		0	Biogeochemical Cycling		0		Biogeochemical Cycling		0	Biogeochemical Cycling		0	Biogeochemical Cycling			0
PART I - Physical, Chemical and	Biological Indica	ators	PART I - Physical, Chemical ar	nd Biological Ind	icators		PART I - Physical, Chemical	and Biological In	dicators	PART I - Physical, Chemical and	Biological Indic	cators	PART I - Physical,	hemical and Bio	logical Indic	cators
	Points Scale Range	Site Score		Points Scale Range	Site Score			Points Scale Range	Site Score		Points Scale Range	Site Score		Pi	oints Scale Range	ge Site Score
PHYSICAL INDICATOR (Applies to all streams	classifications)		PHYSICAL INDICATOR (Applies to all streams	classifications)			PHYSICAL INDICATOR (Applies to all stream	ms classifications)		PHYSICAL INDICATOR (Applies to all stream	s classifications)	1	PHYSICAL INDICATOR (Appli	es to all streams clas	ssifications)	
USEPA RBP (High Gradient Data Sheet)			USEPA RBP (Low Gradient Data Sheet)				USEPA RBP (High Gradient Data Sheet)			USEPA RBP (High Gradient Data Sheet)			USEPA RBP (High Gradient	Data Sheet)		
Epifaunal Substrate/Available Cover	0-20	12	Epifaunal Substrate/Available Cover	0-20			Epifaunal Substrate/Available Cover	0-20		Epifaunal Substrate/Available Cover	0-20		<ol> <li>Epifaunal Substrate/Availab</li> </ol>		0-20	
2. Embeddedness	0-20	5	Pool Substrate Characterization	0-20			2. Embeddedness	0-20		2. Embeddedness	0-20		2. Embeddedness		0-20	
3. Velocity/ Depth Regime	0-20	2	3. Pool Variability	0-20			Velocity/ Depth Regime     Sediment Deposition	0-20		Velocity/ Depth Regime     Sediment Deposition	0-20		Velocity/ Depth Regime     Sediment Deposition		0-20	
4. Sediment Deposition	0-20	6	4. Sediment Deposition	0-20							0-20				0-20	
5. Channel Flow Status	0-20 0-1		5. Channel Flow Status	0-20 0-1			5. Channel Flow Status	0-20 0-1		5. Channel Flow Status	0-20 0-1		5. Channel Flow Status		0-20 0-1	
6. Channel Alteration	0-20	18	6. Channel Alteration	0-20			6. Channel Alteration	0-20		6. Channel Alteration	0-20		6. Channel Alteration		0-20	
7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB)	0-20	20	7. Channel Sinuosity	0-20			7. Frequency of Riffles (or bends)	0-20		7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB)	0-20		7. Frequency of Riffles (or ben		0-20	
	0-20	12	8. Bank Stability (LB & RB)	0-20			8. Bank Stability (LB & RB)	0-20			0-20		8. Bank Stability (LB & RB)		0-20	
Vegetative Protection (LB & RB)     Riparian Vegetative Zone Width (LB & RB)	0-20	20	Vegetative Protection (LB & RB)     Riparian Vegetative Zone Width (LB & RB)	0-20			Vegetative Protection (LB & RB)     Riparian Vegetative Zone Width (LB & RB)	0-20		Vegetative Protection (LB & RB)     Reparian Vegetative Zone Width (LB & RB)	0-20		<ol> <li>Vegetative Protection (LB &amp; 10. Riparian Vegetative Zone Wic</li> </ol>	RB)	0-20	
Total RBP Score	Marginal	101	Total RBP Score	Poor	0		Total RBP Score	Poor	0	Total RBP Score	Poor	0	Total RBP Score	III (LD & ND)	Poor	0
Sub-Total	`	0.505	Sub-Total		0		Sub-Total		Ö	Sub-Total		0	Sub-Total			Ö
CHEMICAL INDICATOR (Applies to Intermitter		eams)	CHEMICAL INDICATOR (Applies to Intermitter		sams)		CHEMICAL INDICATOR (Applies to Intermitt		reams)	CHEMICAL INDICATOR (Applies to Intermitte		reams)	CHEMICAL INDICATOR (App.		d Perennial St	reams)
WVDEP Water Quality Indicators (General Specific Conductivity			WVDEP Water Quality Indicators (General Specific Conductivity	)			WVDEP Water Quality Indicators (General Specific Conductivity	al)		WVDEP Water Quality Indicators (General Specific Conductivity	1)		WVDEP Water Quality Indica Specific Conductivity	tors (General)		
-,,	0-90			0-90				0-90			0-90				0-90	
100-199 - 85 points	0-90			0-90				0-90			0-90				0-90	
pH			pH	- 0.4			pH	- 0.4		рН			рН			
5.6-5.9 = 45 points	0-80			5-90				5-90			5-90				5-90	
DO			DO				DO			DO			DO			
	10-30			10-30				10-30			10-30				10-30	
Sub-Total	1 1		Sub-Total	1 1			Sub-Total	1		Sub-Total	1 1		Sub-Total			0
Sub-Total  BIOLOGICAL INDICATOR (Applies to Intermit	ent and Perennial St	treams)	Sub-Total  BIOLOGICAL INDICATOR (Applies to Intermit	tent and Perennial S	itreams)		Sub-Total BIOLOGICAL INDICATOR (Applies to Inter	rmittent and Perens	nial Streams)	Sub-Total  BIOLOGICAL INDICATOR (Applies to Interr	mittent and Peren	nial Streams)	BIOLOGICAL INDICATOR (A	plies to Intermitte	nt and Pereni	
WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)				WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)			WV Stream Condition Index	(WVSCI)		
0	0-100 0-1			0-100 0-1				0-100 0-1			0-100 0-1				0-100 0-1	
Sub-Total	' '	0	Sub-Total		0		Sub-Total		0	Sub-Total		0	Sub-Total	-		0
PART II - Index and U	nit Score		PART II - Index and	I Unit Score		Ī	PART II - Index ar	nd Unit Score		PART II - Index and U	Init Score		PART II	- Index and Unit	Score	
Index	Linear Feet	Unit Score	Index	Linear Feet	Unit Score		Index	Linear Feet	Unit Score	Index	Linear Feet	Unit Score	Index		Linear Feet	Unit Score
0.653	148	96.57	0	0	0		0	0	0	0	0	0	0		0	0
L	1		L	1	1	l	L		1		1					

## PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME S-IJ83	LOCATION Roanoke County					
STATION # RIVERMILE	STREAM CLASS Intermittent					
LAT <u>37.169211</u> LONG <u>-80.138258</u>	RIVER BASIN Upper Roand	oke				
STORET#	AGENCY VADEQ					
INVESTIGATORS KB/TC						
FORM COMPLETED BY KB	DATE 9/7/21 TIME 9:00 AM	REASON FOR SURVEY Baseline Assessment				



# PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Predom  ✓ Fores  ✓ Field/  ☐ Agric  ☐ Resid	Pasture Industri	ercial ial	Local Watershed NPS  ☑ No evidence ☐ Son ☐ Obvious sources  Local Watershed Erosi ☑ None ☐ Moderate	ne potential sources
RIPARIA VEGETA (18 meter	TION		e the dominant type and S VS		ominant species present ☐ Grasses  ☐ He	rbaceous
INSTREA FEATURI		Estimat Samplin Area in Estimat	ng Reach Area 512  km² (m²x1000)  red Stream Depth 0.03	m² km² m	_ , 1	ly shaded Shaded  Shaded  Run 100 %  No
LARGE V DEBRIS	VOODY	LWD Density	8 m² of LWD n	m²/km² (LWD/	reach area)	
AQUATIO VEGETA		✓ Roote Floati	e the dominant type and demergent RA A A A A A A A A A A A A A A A A A A	ooted submerge ttached Algae		□Free floating
WATER (	QUALITY	Specific Dissolve pH N/A Turbidi	cature NA 0 C c Conductance NA ed Oxygen NA city NA ctrument Used NA	-		Other
SEDIMEN SUBSTRA		Oils		Petroleum None	— Lρoking at stones whic are the undersides blace	□Paper fiber □Sand Other muck h are not deeply embedded, k in color?
INC		STRATE of the state of the stat	COMPONENTS		ORGANIC SUBSTRATE C	
Substrate Type	Diamet	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area
Bedrock			0	Detritus	sticks, wood, coarse plant	0
Boulder	> 256 mm (10")		0		materials (CPOM)	
Cobble	64-256 mm (2.5	5"-10")	0	Muck-Mud	black, very fine organic	30
Gravel	2-64 mm (0.1"-2	2.5")	0		(FPOM)	30
Sand	0.06-2mm (gritt	y)	15	Marl	grey, shell fragments	0
Silt	0.004-0.06 mm		85			l
Clay	< 0.004 mm (sli	ok)	0	7		

Notes: No flow. No water quality measurements collected due to no flow.

#### HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-IJ83	LOCATION Roanoke County					
STATION # RIVERMILE	STREAM CLASS Intermittent					
LAT <u>37.169211</u> LONG <u>-80.138258</u>	RIVER BASIN Upper Roanoke					
STORET#	AGENCY VADEQ					
INVESTIGATORS KB/TC						
FORM COMPLETED BY KB	DATE 97/21 REASON FOR SURVEY TIME 9:00 AM PM Baseline Assessment					

	Habitat		Condition	Category	
	Parameter	Optimal	Suboptimal	Marginal	Poor
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.
	SCORE 12	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.
ted in	SCORE 5	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/ depth regime (usually slow-deep).
ıram	SCORE 2	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
P <sub>2</sub>	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.
	score 4	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.
	score 6	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0

Notes: No flow.

#### HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	ı Category	
	Parameter	Optimal	Suboptimal	Marginal	Poor
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.
	SCORE 18	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
ding reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.
samp	score 2	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank)  Note: determine left or right side by facing dewastream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.
eva	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0
to be	SCORE 10	Right Bank 10 9	8 7 6	5 4 3	2 1 0
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.
	SCORE 6	Left Bank 10 9	8 7 6	5 4 3	2 1 0
	SCORE 6	Right Bank 10 9	8 7 6	5 4 3	2 1 0
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.
	SCORE 10	Left Bank 10 9	8 7 6	5 4 3	2 1 0
	SCORE 10	Right Bank 10 9	8 7 6	5 4 3	2 1 0

101 Notes: No flow.

A-8

#### BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-	M NAME S-IJ83								N Roai	noke	Cou	nty								
STATION #	R	IVE	RMI	LE_			STRE	AM (	CLASS	Inte	rmitte	ent								
LAT 37.169211	L	ONC	· -80.	138258	3		RIVE	R BA	SIN U	рре	r Roa	noke	Э							
STORET#							AGE	NCY	VADEC	)										
INVESTIGATORS	KB/TC	;										]	LOT	NUMBER						
FORM COMPLETE	D BY	K	В				DATI TIME		7/21 00 AM			]	REAS	SON FOR SURVE	Y Base	line	As	ses	sme	∍nt
HABITAT TYPES		Cob	ble_		%	tage of ea Snag	gs	_%	t type p	Vege	etated	Ban	ks	%		, D				
SAMPLE COLLECTION	Ho In	ow v dica Cob	vere te th	the s	samp ımbe	r of jabs/ Snag	cted? kicks	take	□wadi n in eac	ng h ha	Other	from	m baı e. ks_	ik	boat					
GENERAL COMMENTS	- 11					thics i								sence of flo	w an	d h	na	bit	at.	
QUALITATIVE Indicate estimate Dominant					0 = A	Absent/I	Not (	Obse	rved,				; = C	ommon, 3= Ab						
Indicate estimate Dominant  Periphyton	ed abu				0 = A	1 2	<b>Not (</b>	Obse 4	rved,	Sl	imes				(	)	1	2	3	4
Indicate estimate Dominant  Periphyton Filamentous Alga	ed abu				0 = A $0$ $0$	1 2 1 2	3 3	4 4	rved,	Sl M	imes acro		ertebi		(	)	1 1	2	3	4
Indicate estimate Dominant  Periphyton	ed abu	ONS	S OI	F M	0 = A $0$ $0$ $0$ $ACI$ $0 = A$	1 2 1 2 1 2 ROBEN Absent/	3 3 3	4 4 4 4 OS	erved,	S1 M Fi	imes acro sh	inve	ertebi	rates	( ( (	on (	1 1 1 1 (3-9)	2 2	3 3	-
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV	ed abu	ONS	S OI	F M	0 = A $0$ $0$ $0$ $ACI$ $0 = A$	1 2 1 2 1 2 ROBEN Absent/	3 3 3 TTHC (Not (), 3= 1	4 4 4 4 OS	erved,	S1 M Fi	imes acro sh	inve	ertebi	rates rganisms), 2 =	( ( (	on (	1 1 1 1 (3-9)	2 2	3 3	4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate	e e  /ATICed abu	ONS	S OI	F M	0 0 0 ACI 0 = org3	1 2 1 2 1 2 ROBEN Absent/sanisms).	3 3 3 TTH( Not (), 3= 2	4 4 4 4 OS	erved,	Sl M Fi 1 = (>10	imes acro sh Rai	inve re (1	-3 o sms)	rganisms), 2 = , 4 = Dominant	() () () () ()	on (org	1 1 1 (3-9)	2 2 2	3 3 8)	4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes	e ATICed abu	ONS und	S OI ance	F M e:	0 $0$ $0$ ACI $0 = org$	1 2 1 2 1 2 1 2 ROBEN Absent/Anisms).  Anisop Zygop Hemip	3 3 3 TTHC (Not 0, 3= 4)	4 4 4 4 Obse	erved, ndant	Sl M Fi 1 = (>10	imes acro sh Ran	re (1 ganis	l-3 o sms)	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm	) on ( org	1 1 1 (3-9)	2 2 2 9 ism 2 2 2	3 3 s)	4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria	e e  /ATICed abu	ONS Ind	2 2 2 2	F M e: 3	0 0 0 0 <b>ACI</b> 0 = orgs	1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	3 3 3 TTHC Not (, 3= 1)	4 4 4 4 ObseAbun	erved, ndant	Sl M Fi 1 = (>10	Ran 2 2 2 2 2	re (1 ganis	4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera	Comm	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2	3 3 8)	4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea	e ATICed abu	DNS 1 1 1 1 1 1 1	2 2 2 2 2 2	3 3 3 3 3	0 0 0 0 ACI 0 = org:	1 2 1 2 1 2 1 2 Absent/II  ROBEN Absent/Anisms).  Anisop Zygop Hemip Coleop Lepido	3 3 3 TTHO (Not (), 3= 1) ptera tera otera otera	4 4 4 4 ObseAbun	0 0 0 0	Sl M Fi 1 = (>10	Ran 2 2 2 2 2 2	3 3 3 3 3	4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta	e e /ATICed abu	ONS 1 1 1 1 1 1 1 1	2 2 2 2 2 2 2	3 3 3 3 3 3	0 0 0 0 0 ACI 0 = org:	1 2 1 2 1 2 ROBEN Absent/I Anisop Zygop Hemip Coleop Lepido Sialida	3 3 3 3 TTHC  Not 9  No	4 4 4 OS Obse	0 0 0 0 0	S1 M Fi 1 = (>10 1 1 1 1 1 1	Ran 2 2 2 2 2 2	3 3 3 3 3 3 3	4 4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda	ed abu	DNS 1 1 1 1 1 1 1	2 2 2 2 2 2 2	3 3 3 3 3 3 3	0 0 0 0 0 <b>ACCI</b> 0 <b>o o o o o o o o o o</b>	1 2 1 2 1 2 1 2 Absent/In Absent/In Anison Zygop Hemip Coleop Lepido Sialida Coryda	3 3 3 3 TTHC Not 1, 3= 1	4 4 4 OS Obse	0 0 0 0 0 0	S1 M Fi  1 = (>10 1	Ran 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	1-3 o ssms) 4 4 4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda	ed abu	DNS and 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3	0 0 0 0 0 0 0 0 0 0 4 4 4 4 4 4 4 4 4	1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	3 3 3 3 TTHO  (Not ( 1, 3= 1)  ptera  tera  ptera  ptera  ae  ae  alidac  dae	4 4 4 OS Obse	0 0 0 0 0 0 0	S1 M Fi  1 = (>10) 1	Ran 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	4 4 4 4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda Decapoda	ed abu	DNS 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3	0 0 0 0 0 0 0 0 0 4 4 4 4 4 4 4 4 4 4 4	Absent/I  1 2 1 2 1 2 1 2  ROBEN Absent/ anisms).  Anisop Zygop Hemip Coleop Lepido Sialida Coryda Tipulio Empido	3 3 3 3 TTHC (Not c) , 3= 1  ptera ptera ptera ptera ptera dae lidae	4 4 4 OS Obse	0 0 0 0 0 0 0	S1 M Fi  1 = (>1(1) 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Rain	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	4 4 4 4 4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4
Periphyton Filamentous Alga Macrophytes  FIELD OBSERV Indicate estimate  Porifera Hydrozoa Platyhelminthes Turbellaria Hirudinea Oligochaeta Isopoda Amphipoda	ed abu	DNS and 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3	0 0 0 0 0 0 0 0 0 0 4 4 4 4 4 4 4 4 4	1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	3 3 3 3 TTHC (Note a potera tera potera dae dae didae	4 4 4 OS Obse	0 0 0 0 0 0 0	S1 M Fi  1 = (>10) 1	Ran 2 2 2 2 2 2 2 2 2 2	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	4 4 4 4 4 4 4	rganisms), 2 = , 4 = Dominant  Chironomidae Ephemeroptera Trichoptera	Comm ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) ( ) (	on (org	1 1 1 (3-9 ani	2 2 2 9 ism 2 2 2	3 3 3 3 3 3	4 4 4 4

#### WOLMAN PEBBLE COUNT FORM

County: Roanoke County Stream ID: S-IJ83

Stream Name: UNT to Bottom Creek

HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 9/7/2021 Surveyors: KB, TC Type: Representative

			LE COUNT			Т	
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cun
	Silt/Clay	< .062	S/C	<b>A</b>	24	24.00	24.00
	Very Fine	.062125		<b>4</b>	0	0.00	24.00
	Fine	.12525		•	19	19.00	43.00
	Medium	.255	SAND	<b>4</b>	14	14.00	57.00
	Coarse	.50-1.0		•	0	0.00	57.00
.0408	Very Coarse	1.0-2	7	<b>4</b>	0	0.00	57.00
.0816	Very Fine	2 -4		<b>A</b>	3	3.00	60.00
.1622	Fine	4 -5.7	1	<b>A</b>	3	3.00	63.00
.2231	Fine	5.7 - 8	1	<b>4</b>	9	9.00	72.00
.3144	Medium	8 -11.3		<b>A</b>	9	9.00	81.00
.4463	Medium	11.3 - 16	GRAVEL	<b>A</b>	8	8.00	89.00
.6389	Coarse	16 -22.6	1	<b>A</b>	5	5.00	94.00
.89 - 1.26	Coarse	22.6 - 32	1	<b>A</b>	4	4.00	98.00
1.26 - 1.77	Vry Coarse	32 - 45	1	<b>A</b>	2	2.00	100.0
1.77 -2.5	Vry Coarse	45 - 64	1	<b>A</b>	0	0.00	100.0
2.5 - 3.5	Small	64 - 90		<b>A</b>	0	0.00	100.0
3.5 - 5.0	Small	90 - 128	1	<b>A</b>	0	0.00	100.0
5.0 - 7.1	Large	128 - 180	COBBLE	<b>A</b>	0	0.00	100.0
7.1 - 10.1	Large	180 - 256	1	<b>A</b>	0	0.00	100.0
10.1 - 14.3	Small	256 - 362		<b>A</b>	0	0.00	100.0
14.3 - 20	Small	362 - 512	1	<b>A</b>	0	0.00	100.0
20 - 40	Medium	512 - 1024	BOULDER	<b>A</b>	0	0.00	100.0
40 - 80	Large	1024 -2048	1	<b>A</b>	0	0.00	100.0
80 - 160	Vry Large	2048 -4096	1	<b>A</b>	0	0.00	100.0
	Bedrock		BDRK	<b>A</b>	0	0.00	100.0
				Totals	100		

#### RIVERMORPH PARTICLE SUMMARY

River Name: UNT to Bottom Creek Reach Name: S-IJ83 Representative 09/07/2021

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	24 0 19 14 0 0 3 3 3 9 9 8 5 4 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	24.00 0.00 19.00 14.00 0.00 0.00 3.00 3.00 9.00 9.00 9.00 4.00 2.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00	24.00 24.00 43.00 57.00 57.00 60.00 63.00 72.00 81.00 89.00 94.00 98.00 100.00 100.00 100.00 100.00 100.00 100.00 100.00 100.00 100.00 100.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.04 0.2 0.38 13.06 24.95 45 24 33 43 0		

Total Particles = 100.

				Unified S	tream Method	lology for use			1)		
Project #	Project N	Nama (Ann			Cowardin	HUC	Date	SAR#	Impact	Impact	
Project #	Mountain Vall	Name (App		Locality	Class.	нос	Date	SAR#	Length	Factor	
22865.06		Pipeline, L		Roanoke County	R4	03010101	9/7/21	S-IJ83	148	1	
Name	e(s) of Evaluator	r(s)	Stream Name		tion				SAR Length		
	KB, TC		UNT to Botto	m Creek					2	10	
Channel C	ondition: Assess t	he cross-section	on of the stream a	ind prevailing con	dition (erosion, ag Conditional Catego						
	Optima	al	Subo	ptimal	_	ginal	Po	or	Sev	/ere	
Channel Condition	Very little incision or act 100% stable banks. Veg protection or natural ro (80-100%). AND/OR Site bankfull benches are pr to their original flood developed wide bankfull channel bars and transt Transient sediment depless than 10% of	getative surface ock, prominent able point bars / resent. Access Iplain or fully I benches. Mid- verse bars few. position covers	erosion or unprotect of banks are st Vegetative protect prominent (60. Depositional feat stability. The bar channels are well de has access to ba newly developed	ew areas of active cted banks. Majority table (60-80%). Ition or natural rock-80%) AND/OR ures contribute to kfull and low flow efined. Stream likely inkfull benchesory infoodplains along	Poor. Banks more or Poor due to Ic Erosion may be pr both banks. Vege 40-60% of banks. S vertical or und 40-60% Sediment transient, contr Deposition that co	less than Severe or stable than Severe over the stable stank slopes: we seem on 40-60% of tative protection on Streambanks may be ercut. AND/OR may be temporary / ibute instability.	laterally unstable further. Majority of vertical. Erosion pr banks. Vegetative on 20-40% of banks to prevent erosion. the stream is cove Sediment is temp nature, and contril	ised. Vertically / e. Likely to widen both banks are near esent on 60-80% of protection present AND/OR 60-80% of ered by sediment, orary / transient in buting to instability.	vertical/lateral in incision, flow contair Streambed below av majority of banks Vegetative protect than 20% of banks erosion. Obviou present. Erosion/ray AND/OR Aggradin	ion present on less s, is not preventing s bank sloughing v banks on 80-100%. ng channel. Greater	
0	2		sediment covers 1 bott		shaped channels protection on > 40 depositional featur to sta	resent. AND/OR V- s have vegetative % of the banks and res which contribute ability.	vegetative protect 40% of the banks a deposition		deposition, contrib Multiple thread subterran	channels and/or nean flow.	CI
Scores	3		2	.4		2	1.	.6	<u> </u>	1	3.00
	Optima	al		ditional Cate	gory	ginal		ay be acceptable)	NOTES>>		
Riparian Buffers	Optima  Tree stratum (dbh > 3 in with > 60% tree can Wetlands located with areas.	nches) present, nopy cover.	Subo  High Suboptimal: Riparian areas with	ptimal Low Suboptimal:	gory Mar	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree		, ,	NOTES>>		
	Tree stratum (dbh > 3 ir with > 60% tree car Wetlands located with	nches) present, nopy cover.	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy conditions and containing both herbaceous and shub layers or a non-maintained	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Receni cutover (dense	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbb - 3 inches) present, with <30%	Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh > 3 inches) present, with <30% tree canopy cover with maintained	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, tralls, or other comparable	NOTES>>		
	Tree stratum (dbh > 3 ir with > 60% tree car Wetlands located with	nches) present, nopy cover.	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recen cutover (dense vegetation).	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NOTES>>		
Scores  Delineate ripar  Determine squ	Tree stratum (dbh > 3 ir with > 60% tree car Wetlands located with areas.	nches) present, nopy cover. in the riparian	High Suboptimal: Riparian areas with tree stratum (dbh 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High  1.2  Into Condition Cate  or estimating lengers	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Condith and width. Cal	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh -3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	NOTES>>		
Scores  Delineate ripar Determine squ Enter the % Ri	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5	nches) present, nopy cover. in the riparian stream bank is tream bank if the present of the pres	High Suboptimal: Riparian areas with tree stratum (dbh 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High 1.2  Into Condition Cate or estimating lenguarian category in the 45%	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Condith and width. Cal	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh -3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5	NOTES>>		
Scores  Delineate ripar Determine squ Enter the % Ri	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  Tian areas along each lare footage for each laparian Area and Scor	nches) present, nopy cover. in the riparian  stream bank i	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High 1.2  Into Condition Cate	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Concetth and width. Caller blocks below.	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh -3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  he sums		10000 10 10 10 10 10 10 10 10 10 10 10 1	
Scores  Delineate ripar Determine squ Enter the % Ri Right Bank	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  Tian areas along each lare footage for each lare footage for each laparian Area and Scor % Riparian Area>	nches) present, nopy cover. in the riparian stream bank is stream bank is by measuring re for each riparian 1.5	Subol  High Suboptimal: Riparian areas with tree stratum (dbh - 30 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High 1.2  Into Condition Cate or estimating lenguarian category in the 45% 1.2	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Condith and width. Calle blocks below.	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh -3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  he sums ciparian qual 100 100%	NOTES>>  CI= (Sum % RA * Sc Rt Bank CI >	cores*0.01//2	CI
Scores  Delineate ripar Determine squ Enter the % Ri Right Bank	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  Tian areas along each lare footage for each lare footage for each laparian Area and Scor % Riparian Area>  Score >	nches) present, nopy cover. in the riparian stream bank is tream bank if the present of the pres	High Suboptimal: Riparian areas with tree stratum (dbh 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High 1.2  Into Condition Cate or estimating lenguarian category in the 45%	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Condith and width. Calle blocks below.	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh. > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh -3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  he sums	CI= (Sum % RA * Sc		CI 1.37
Scores  Delineate ripar Determine squ Enter the % Ri Right Bank  Left Bank	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  Tian areas along each parian Area and Score >  % Riparian Area > Score >  1 HABITAT: Varied	nches) present, nopy cover. in the riparian  stream bank i by measuring re for each ripa  50%  1.5  70%  1.5	Subol  High Suboptimal: Riparian areas with tree stratum (dsh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High 1.2  Into Condition Cate or estimating leng arian category in the 45% 1.2  30% 1.2	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recen cutover (dense vegetation).  Low 1.1  egories and Cond th and width. Cal ne blocks below. 5% 0.75	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.  High 0.85	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75  the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure t  of % F  Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  the sums tiparian qual 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI >	1.33 1.41	
Scores  Delineate ripar Determine squ Enter the % Ri Right Bank  Left Bank . INSTREAM omplexes, stable	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  1.5  Tian areas along each parian Area and Score >  % Riparian Area > Score >  1 HABITAT: Varied = features.	stream bank is by measuring re for each ripa 1.5 70% 1.5	Subol  High Suboptimal: Riparian areas with tree stratum (dbh 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High  1.2  Into Condition Cate or estimating lenguarian category in the 45%  1.2  30%  1.2  as, water velocity as, water velocity as a simular and a sim	Low Suboptimal: Riparian areas with tree stratum (db. 73 inches) present with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).  Low 1.1  egories and Condition and width. Caller blocks below.  5% 0.75	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.  High 0.85  ition Scores using culators are provided to the control of the	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75  the descriptors. ded for you below.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure to fine Blocks e	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  he sums ciparian qual 100 100%	CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI >	1.33 1.41	
Scores  Delineate ripar Determine squ Enter the % Ri Right Bank  Left Bank	Tree stratum (dbh > 3 in with > 60% tree car Wetlands located with areas.  1.5  Tian areas along each parian Area and Score >  % Riparian Area > Score >  1 HABITAT: Varied	nches) present, nopy cover. in the riparian  stream bank is by measuring re for each ripa 50% 1.5  70% 1.5  I substrate size	Subol  High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.  High  1.2  Into Condition Cate or estimating lenguarian category in the desired of the desired of the desired or estimating lenguarian category in the desired of the desired or estimating lenguarian category in the desired or estimation category in the de	Low Suboptimal: Riparian areas with tree stratum (dbh a) inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recenicutover (dense vegetation).  Low 1.1  egories and Condition and width. Call the blocks below. 5% 0.75  and depths; wood  Condition. ptimal  ments are typically of the reach and are maintenance of	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.  High 0.85  ition Scores using culators are provided to the control of the	ginal  Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.  Low  0.75  the descriptors. ded for you below.	High Poor: Laws, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.  High  0.6  Ensure to fine Blocks expenses we will be been seed to fine Blocks expenses when the seed to fine Blocks expenses we will be been seed to fine Blocks expenses when the seed to fine Blocks expenses we will be be been seed to fine Blocks expenses when the seed to fine Blocks expenses we will be be been seed to fine Blocks expenses when the seed to fine Blocks expenses we will be be been seed to fine Blocks expenses to fine Blocks expense	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.  Low 0.5  he sums  Riparian qual 100  100%  1100%	CI= (Sum % RA * So Rt Bank CI > Lt Bank CI > banks; root mats; \$	1.33 1.41	

	Stream II	npact A	ssessn	nent For	rm Page	2		
Project #	Project Name (Applicant)	Locality	Cowardin Class.	HUC	Date	SAR#	Impact Length	Impact Factor
22865.06	Mountain Valley Pipeline (Mountain Valley Pipeline, LLC)	Roanoke County	R4	03010101	9/7/21	S-IJ83	148	1
4. CHANNEL	. ALTERATION: Stream crossings, riprap, concre	te, gabions, or cor	ncrete blocks, stra	ghtening of chann	nel, channelization	, embankments, s	poil piles, constriction	ons, livestock

Conditional Category NOTES>>

			Conditions	ar Category		
	Negligible	Mi	nor	Mod	erate	Severe
Channel Alteration	Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized.	Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	disrupted by any of the channel	of the channel alterations listed in the parameter guidelines. If	60 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	
Scores	1.5	1.3	1.1	0.9	0.7	0.5

CI 1.50

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

THE REA

THE REACH CONDITION INDEX (RCI) >>

1.41

 $\label{eq:RCI} \mbox{RCI= (Sum of all CI's)/5, except if stream is ephemeral RCI = (Riparian CI/2)}$ 

COMPENSATION REQUIREMENT (CR) >>

>> 209

CR = RCI X L<sub>I</sub> X IF

#### INSERT PHOTOS:


(WSSI\_Photo Location L:\22000s\22800\22865.06\Admin\05-ENVR\Field Data\Spread H\Field Forms\S-IJ83\Photos\DS VIEW.JPG)



 $\label{eq:constream} \mbox{Downstream view within the ROW. Assessment is limited to areas within the temporary ROW.}$ 

DESCRIE	
RE PROP	
OSED	
IMPΔ	
СΤ٠	

PROVIDED UNDER SEPARATE COVER





#### SURVEY NOTES:


LEGEND

EXISTING SURVEY-LOCATED THALWEG

- 1. THIS MAP HAS BEEN ORIENTED TO NAD 1983 UTM ZONE 17N, AND VERTICALLY TO THE NORTH AMERICAN VERTICAL DATUM OF 1988 (NAVD 88), USING REAL TIME DGPS. FIELD LOCATIONS WERE COMPLETED ON SEPTEMBER 7, 2021.
- 2. EASEMENT LINES SHOWN ON PLAN VIEW WERE PROVIDED BY MOUNTAIN VALLEY PIPELINE.
- 3. SURVEY POINTS FOR CROSS SECTIONS AND THALWEG PROFILES COLLECTED IN 2021 HAVE BEEN USED IN COMBINATION WITH SURVEY POINTS COLLECTED PREVIOUSLY IN 2020 IN ORDER TO GENERATE THE PRE-CROSSING SURFACE SHOWN IN PLAN. DUE TO NATURAL EROSIONAL STREAM PROCESSES THAT CAN OCCUR OVER TIME, MINOR ADJUSTMENTS TO THE PROFILE ALIGNMENTS MAY HAVE BEEN REQUIRED IN ORDER TO GENERATE A CLEAN PRE-CROSSING SURFACE.
- 4. ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.








TS-R | 13497266.581 | 1891430.300 | 2679.515

PROFILE LEGEND

EXISTING STREAM PROFILE



CROSS SECTION

H: 1"=10'

V: 1"=5'


PRE-CROSSING PHOTOS



PHOTO TAKEN LOOKING DOWNSTREAM FROM UPSTREAM IMPACT LIMITS ON 09/07/21



PHOTO TAKEN LOOKING UPSTREAM FROM DOWNSTREAM IMPACT LIMITS ON 09/07/21



FROM UPSTREAM IMPACT LIMITS




PHOTO TAKEN LOOKING UPSTREAM FROM DOWNSTREAM IMPACT LIMITS

Checked

CAD File No

Drawn

112IC07157 Project No.

Drawing No.