Baseline Assessment - Stream Attributes

Reach S-MM15 (Pipeline ROW) Intermittent Spread H Montgomery County, Virginia

| Data | Included |
|---|------------------------|
| Photos | ✓ |
| SWVM Form | ✓ |
| FCI Calculator and HGM Form | ✓ |
| RBP Physical Characteristics Form | ✓ |
| Water Quality Data | N/A – No water present |
| RBP Habitat Form | ✓ |
| RBP Benthic Form | ✓ |
| Benthic Identification Sheet | N/A - No water present |
| Wolman Pebble Count | ✓ |
| RiverMorph Data Sheet | ✓ |
| USM Form (Virginia Only) | ✓ |
| Longitudinal Profile and Cross Sections | ✓ |



Photo Type: DS VIEW Location, Orientation, Photographer Initials: Downstream view of ROW looking S, AO



Photo Type: US VIEW Location, Orientation, Photographer Initials: Upstream view of ROW looking N, AO



Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking SW, AO



Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking E, AO



Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking S, AO

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | | Mountain | Valley Pipeline | IMPACT COORDINATES (in Decimal Degrees) | E Lat. | 37.258673 Lon. | -80.296446 | WEATHER: | Sunny | DATE: | August 25 | 5, 2021 | |
|--|----------------------|------------------------|---|--|--------|--|--|---|---|---|-----------------------|-------------|--|
| IMPACT STREAM/SITE ID (watershed size (acreage), | | | S-M | M15 | | MITIGATION STREAM CLASS./SITE ID AI (watershed size (acreage), unaltered or | | i: | | Comments: | | | |
| STREAM IMPACT LENGTH: | 82 | FORM OF MITIGATION: | RESTORATION (Levels I-III) | MIT COORDINATES: (in Decimal Degrees) | Lat. | Lon. | | PRECIPITATION PAST 48 HRS: | None | Mitigation Length: | | | |
| Column No. 1- Impact Existing Condition (Debit) | | | Column No. 2- Mitigation Existing Co | ondition - Baseline (Credit) | | Column No. 3- Mitigation Projected at F Post Completion (Credit) | Column No. 4- Mitigation Proj Post Completion (| | Column No. 5- Mitigation Projected at Maturity (Credit) | | | | |
| Stream Classification: Intermittent | | nittent | Stream Classification: | | | Stream Classification: | 0 | Stream Classification: | 0 | Stream Classification: | 0 | 0 | |
| Percent Stream Channel Si | lope | 13.62 | Percent Stream Channel Sig | ре | | Percent Stream Channel Slope | 0 | Percent Stream Channel S | lope 0 | Percent Stream Channel S | lope | 0 | |
| HGM Score (attach d | lata forms): | | HGM Score (attach o | lata forms): | | HGM Score (attach data forms | s): | HGM Score (attach d | ata forms): | HGM Score (attach d | ata forms): | | |
| | | Average | | Average | | | Average | | Average | | | Average | |
| Hydrology Biogeochemical Cycling | 0.47 | 0.32 | Hydrology Biogeochemical Cycling | 0 | | Hydrology Biogeochemical Cycling | 0 | Hydrology Biogeochemical Cycling | • | Hydrology Biogeochemical Cycling | | | |
| Habitat | 0.13 | 0.32 | Habitat | • | | Habitat | • | Habitat | • | Habitat | | ۰ | |
| PART I - Physical, Chemical and | l Biological Indic | ators | PART I - Physical, Chemical and | Biological Indicators | | PART I - Physical, Chemical and Biological | I Indicators | PART I - Physical, Chemical and | Biological Indicators | PART I - Physical, Chemical and | Biological Indicat | tors | |
| | Points Scale Range | Site Score | | Points Scale Range Site Score | | Points Scale | Range Site Score | | Points Scale Range Site Score | | Points Scale Range | Site Score | |
| PHYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams of | dassifications) | | PHYSICAL INDICATOR (Applies to all streams classification | 3) | PHYSICAL INDICATOR (Applies to all stream | s classifications) | PHYSICAL INDICATOR (Applies to all streams | classifications) | | |
| USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Low Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (High Gradient Data Sheet) | | USEPA RBP (High Gradient Data Sheet) | | | |
| Epifaunal Substrate/Available Cover Embeddedness | 0-20 | 11 | Epifaunal Substrate/Available Cover Pool Substrate Characterization | 0-20 | | Epifaunal Substrate/Available Cover 0-20 Embeddedness 0-20 | | Epifaunal Substrate/Available Cover Embeddedness | 0-20 | Epifaunal Substrate/Available Cover Embeddedness | 0-20 | | |
| Velocity/ Depth Regime | 0-20 | 0 | Pool Substrate Characterization Pool Variability | 0-20 | | 3. Velocity/ Depth Regime 0-20 | | Velocity/ Depth Regime | 0-20 | 3. Velocity/ Depth Regime | 0-20 | | |
| 4. Sediment Deposition | 0-20 | 8 | Sediment Deposition | 0-20 | | 4. Sediment Deposition 0-20 | | Velocity/ Depart Regime Sediment Deposition | 0-20 | 4. Sediment Deposition | 0-20 | | |
| 5. Channel Flow Status | 0-20 | 0 | 5. Channel Flow Status | 0-20 | | 5. Channel Flow Status 0-20 | | 5. Channel Flow Status | 0-20 | 5. Channel Flow Status | 0-20 | | |
| 6. Channel Alteration | 0-20 0-1 | 15 | 6. Channel Alteration | 0-20 | | 6. Channel Alteration 0-20 | 0-1 | 6. Channel Alteration | 0-20 | Channel Alteration | 0-20 | | |
| 7. Frequency of Riffles (or bends) | 0-20 | 0 | 7. Channel Sinuosity | 0-20 | | 7. Frequency of Riffles (or bends) 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | 7. Frequency of Riffles (or bends) | 0-20 | | |
| 8. Bank Stability (LB & RB) | 0-20 | 14 | 8. Bank Stability (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) 0-20 | | 8. Bank Stability (LB & RB) | 0-20 | 8. Bank Stability (LB & RB) | 0-20 | | |
| Vegetative Protection (LB & RB) | 0-20 | 14 | Vegetative Protection (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) 0-20 | | Vegetative Protection (LB & RB) | 0-20 | Vegetative Protection (LB & RB) | 0-20 | | |
| 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | 17 | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | 10. Riparian Vegetative Zone Width (LB & RB) 0-20 | • | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | |
| Total RBP Score Sub-Total | Marginal | 79 0.395 | Total RBP Score Sub-Total | Poor 0 | | Total RBP Score Poor Sub-Total | 0 | Total RBP Score Sub-Total | Poor 0 | Total RBP Score Sub-Total | Poor | 0 | |
| CHEMICAL INDICATOR (Applies to Intermitter | nt and Perennial Str | | CHEMICAL INDICATOR (Applies to Intermittent | | | CHEMICAL INDICATOR (Applies to Intermittent and Perenn | | CHEMICAL INDICATOR (Applies to Intermitte | | CHEMICAL INDICATOR (Applies to Intermitter | nt and Perennial Stre | | |
| WVDEP Water Quality Indicators (General | D | | WVDEP Water Quality Indicators (General) | | | WVDEP Water Quality Indicators (General) | | WVDEP Water Quality Indicators (Genera | D | WVDEP Water Quality Indicators (General) | | | |
| Specific Conductivity | | | Specific Conductivity | | | Specific Conductivity | | Specific Conductivity | | Specific Conductivity | | | |
| 100-199 - 85 points pH | 0-90 | | pH | 0-90 | | 0-90 pH | | pH | 0-90 | pH | 0-90 | | |
| 5.6-5.9 = 45 points | 0-80 | | | 5-90 0-1 | | 5-90 | 0-1 | | 5-90 0-1 | | 5-90 0-1 | | |
| DO | 10-30 | | DO | 10-30 | | DO 10-30 | | DO | 10-30 | DO | 10-30 | | |
| Sub-Total | 10-30 | | Sub-Total | 10-30 | | Sub-Total | 0 | Sub-Total | 10-30 | Sub-Total | 10-30 | | |
| BIOLOGICAL INDICATOR (Applies to Intermit | ttent and Perennial | Streams) | BIOLOGICAL INDICATOR (Applies to Intermitte | <u> </u> | | BIOLOGICAL INDICATOR (Applies to Intermittent and Pe | | BIOLOGICAL INDICATOR (Applies to Intern | | BIOLOGICAL INDICATOR (Applies to Interm | littent and Perennia | al Streams) | |
| WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | | WV Stream Condition Index (WVSCI) | | | |
| 0 | 0-100 0-1 | | | 0-100 0-1 | | 0-100 | 0-1 | | 0-100 0-1 | | 0-100 0-1 | | |
| Sub-Total | 1 1 | 0 | Sub-Total | 0 | | Sub-Total | 0 | Sub-Total | 0 | Sub-Total | | 0 | |
| PART II - Index and U | Unit Score | | PART II - Index and I | Unit Score | | PART II - Index and Unit Score | | PART II - Index and U | Init Score | PART II - Index and U | Init Score | | |
| Index | Linear Feet | Unit Score | Index | Linear Feet Unit Score | | Index Linear F | eet Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet | Unit Score | |
| 0.459 | 82 | 37.6175 | 0 | 0 0 | | 0 0 | 0 | 0 | 0 0 | 0 | 0 | 0 | |
| | 1 | | | L | | L | | L | | ╙ | | | |

Ver. 10-20-17

FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the <u>UPPERMOST STRATUM</u> of the plant community is determined based on the calculated value for V_{CCANOPY} (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

Project Name: Mountain Valley Pipeline

Location: Spread H; Montgomery County

Sampling Date: 8/25/2021 Project Site Before Project

Subclass for this SAR:

Intermittent Stream

Uppermost stratum present at this SAR: SAR number: S-MM15

Shrub/Herb Strata

Functional Results Summary: Er

Enter Results in Section A of the Mitigation Sufficiency Calculator

| Function | Functional Capacity Index |
|------------------------|------------------------------|
| Hydrology | 0.47 |
| Biogeochemical Cycling | 0.36 |
| Habitat | 0.13 |

Variable Measure and Subindex Summary:

| Variable | Name | Average Measure | Subindex |
|------------------------|---|--------------------|----------|
| V _{CCANOPY} | Percent canpoy over channel. | Not Used, <20% | Not Used |
| V_{EMBED} | Average embeddedness of channel. | 1.67 | 0.34 |
| V _{SUBSTRATE} | Median stream channel substrate particle size. | 0.08 | 0.04 |
| V_{BERO} | Total percent of eroded stream channel bank. | 53.13 | 0.79 |
| V_{LWD} | Number of down woody stems per 100 feet of stream. | 0.00 | 0.00 |
| V_{TDBH} | Average dbh of trees. | Not Used | Not Used |
| V _{SNAG} | Number of snags per 100 feet of stream. | 0.00 | 0.10 |
| V_{SSD} | Number of saplings and shrubs per 100 feet of stream. | 343.75 | 1.00 |
| V _{SRICH} | Riparian vegetation species richness. | 0.00 | 0.00 |
| V _{DETRITUS} | Average percent cover of leaves, sticks, etc. | 81.25 | 0.99 |
| V_{HERB} | Average percent cover of herbaceous vegetation. | 28.75 | 0.38 |
| V _{WLUSE} | Weighted Average of Runoff Score for Catchment. | 0.88 | 0.93 |

| | | | High-G | | | ter Strea et and C | | | a | Verein | 011 10-20-17 |
|-------------|----------------------------------|--|---|--|---|---|--|--|---|---------------------------------------|----------------------------------|
| | Team: | AO, MM | | | | | | Latitude/UT | M Northing: | 37.258673 | |
| Pr | oject Name: | | | | | | L | - | - | -80.296446 | i |
| | | Spread H; I | | | | | | San | pling Date: | 8/25/2021 | |
| Si | AR Number: | | | Length (ft): | 32 | Stream Ty | | mittent Strea | | | _ |
| Sito | Top Strata: and Timing: | | rub/Herb Str | ata | (determine | d from perce | | | νγ) | | _ |
| | | | | | | | Before Proje | ct | | | • |
| Sample 1 | e Variables V _{CCANOPY} | Average pe equidistant | rcent cover points along at least one | the stream value betw | . Measure een 0 and 1 | nd sapling ca only if tree/s 9 to trigger | apling cove | r is at least : | | 0 , | Not Used, <20% |
| | 0 | CONT. CO VOI 1 | noadaremer | no at oaon p | Joint Bolow. | | | | | | 1 |
| | | | | | | | | | | | |
| 2 | V _{EMBED} | along the si surface and to the follow of 1. If the | tream. Sele I area surro ving table. I bed is comp | ect a particle unding the p f the bed is posed of bed | from the be particle that in an artificial in drock, use a | I. Measure ed. Before n is covered b surface, or o rating score | noving it, de y fine sedim composed o e of 5. | termine the nent, and en f fine sedime | percentage ter the rating ents, use a r | of the g according rating score | 1.7 |
| | | Minshall 19 | 83) | | obble and b | oulder partio | cies (rescale | d from Platt | s, Meganan | , and | Measure at least 30 points |
| | | Rating 5 | Rating Des <5 percent | | overed, sur | rounded, or | buried by fir | ne sediment | (or bedrock | <u> </u> | oo points |
| | | 4 | 5 to 25 per | cent of surfa | ce covered, | surrounded | l, or buried l | y fine sedin | nent | | |
| | | 3 2 | | | | d, surrounde d, surrounde | - | , | | | |
| | | 1 | >75 percen | t of surface | | rrounded, o | | | | al surface) | |
| | | ngs at each | | | 1 | 1 | 1 | 1 | | 1 | 1 |
| | 3 | 3 | 1 | 1 | 3 | 3 | 1 | 1 | 1 | 2 | |
| | 3 | 2 | 1 | ' | 3 | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| 3 | Enter partic | Median stre along the size in inc as 0.0 in, size | tream; use t ches to the i | he same po nearest 0.1 | ints and par inch at each | ticles as use | ed in V _{EMBED} | | | | 0.08 in |
| | 0.08 | 0.70 | 0.08 | 0.08 | 0.08 | 0.60 | 0.08 | 0.08 | 0.08 | 0.50 | 1 |
| | 4.90 | 1.60 | 1.20 | 0.08 | 1.80 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| 4 | V_{BERO} | side and the | e total perce | | | Enter the to | | | | | 53 % |
| | | may be up | to 200%. Left Bank: | 12 | 2 ft | | Right Bank: | 5 | ft | | |
| ampl | e Variables | 5-9 within t | he entire ri | parian/buff | er zone adj | acent to the | stream ch | annel (25 fe | et from ea | ch bank). | |
| 5 | V_{LWD} | stream read | | e number fr | om the entir lated. | es in diamete e 50'-wide b | uffer and w | thin the cha | | | 0.0 |
| 6 | V_{TDBH} | | | | y if V _{CCANOP} tree DBHs i | _Y tree/saplin n inches. | g cover is a | t least 20%) | Trees are | at least 4 | Not Used |
| | | List the dbh the stream | below: | ents of indiv | ridual trees (| (at least 4 in |) within the | | ch side of | | |
| | | | Left Side | | | | | Right Side | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| 7 | V_{SNAG} | | | | | per 100 feet et will be cal | | Enter numb | er of snags | on each | 0.0 |
| | | | Left Side: | | 0 | | Right Side: | |) | | |
| 8 | V _{SSD} | Number of | | | _ | up to 4 inch | | | | asure only if | |
| | - | tree cover i | s <20%). E f stream will | nter number I be calculat | of saplings ed. | and shrubs | on each sid | le of the stre | eam, and the | | 343.8 |
| | | | Left Side: | 6 | 5 | | Right Side: | 4 | 5 | | |

| 9 V _{SRICH} | Group 1 in richness pe | er 100 feet a | nd the subir | | caiculated fi | rom mese da | ala. | | | | | | |
|---|--|---|--|--|-----------------------------|--|--------------------------|--|--|--|--|--|--|
| | | ıp 1 = 1.0 | | | I | | | p 2 (-1.0) | | | | | |
| Acer ru | ıbrum | | Magnolia tr | ripetala | 7 | Ailanthus a | ltissima | | Lonicera ja | ponica | | | |
| Acer s | accharum | | Nyssa sylva | atica | | Albizia julib | rissin | | Lonicera ta | tarica | | | |
| - ∃ Aescui | lus flava | | Oxydendrum | | | Alliaria peti | | | Lotus corni | iculatus | | | |
| _ | a triloba | | Prunus sen | | | | | | Lythrum sali | | | | |
| | alleghaniensis | | Quercus alba | | | Alternanthe philoxeroide | | | Microstegiun | | | | |
| _ | = | | Quercus coccinea | | | | | | | | | | |
| Betula | | | | | | Aster tatari | | | | Paulownia tomentosa | | | |
| Carya | | | Quercus imbricaria Cerastium fontanum | | | | | Polygonum (| | | | | |
| Carya | glabra | | Quercus pr | rinus | | Coronilla va | aria | | Pueraria m | ontana | | | |
| Carya | ovalis | | Quercus rubra Elaeagnus umbellata | | | | | | Rosa multi | flora | | | |
| Carya | ovata | | Quercus velutina Lespedeza bicolor | | | | | | Sorghum h | alepens | | | |
| Cornus | s florida | | Sassafras a | albidum | | Lespedeza | cuneata | | Verbena bi | rasiliensi | | | |
| Fagus | grandifolia | | Tilia americ | cana | | Ligustrum ob | tusifolium | | | | | | |
| Fraxin | Fraxinus americana | | Tsuga cana | adensis | | Ligustrum s | sinense | | | | | | |
| Lirioder | Liriodendron tulipifera | | Ulmus ame | ericana | | | | | | | | | |
| _ | lia acuminata | _ | | | | | | | | | | | |
| _ wagno | ma acammata | | | | | | | | | | | | |
| | 2 | Species in | Group 1 | | | | 2 | Species | in Group 2 | | | | |
| | | uld be place ercent cover | of leaves, s | equidistant ticks, or oth | ly along ea er organic n | ch side of t | he strean | 1. | in 25 feet from eter and <36" | n each 81.25 | | | |
| | long are in | | | t cover or ur | e uetritai iay | | | | _ | | | | |
| | 05 | 100 | Side | | 50 | Right 80 | Side | | | | | | |
| | 95 | 100 | | | 50 | 80 | | | | | | | |
| I1 V _{HERB} | include wo | ody stems a | t least 4" db | h and 36" ta | all. Because | | e several | layers of g | Do <i>not</i> ound cover vegetation at | 29 % | | | |
| · | vegetation each subpl | ot. | | 1 200% are a | accepted. E | Diabi | | | _ | | | | |
| ·· VHERB | each subpl | ot. Left | Side | 1 200% are a | | | Side | | | | | | |
| , i.e. to | each subpl | Left | Side | he stream. | 70 | Right 25 | | | | 0.88 | | | |
| mple Variat | each subpl | ot. Left 10 e entire cate Average of F | Side chment of t | he stream. | 70 | | | Runof | % in Catch | Runni | | | |
| mple Variat | each subple 10 Die 12 within the Weighted / | Left 10 e entire cate Average of F | Side chment of t cunoff Score | he stream. | 70 | | | | | Runni Perce (not >1 | | | |
| mple Variat | each subpl | Left 10 e entire cate Average of F | Side chment of t cunoff Score | he stream. | 70 | | | Runof | | Runni Perce (not >1 | | | |
| mple Variat 2 VwLuse Forest | each subple 10 Die 12 within the Weighted / | Left 10 10 Average of F | Side Chment of t Runoff Score Use (Choose | he stream. | 70 | | | Runof | ment | Runni Perce (not >1 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | | Runof Score | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplication of the second subplication of | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | | Runof Score | ment 22 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | | Runof Score | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variat 12 VwLuse Forest a | each subplement of the subplem | Land | Side Chment of t Runoff Score Use (Choose Cover) | he stream. for watersh | 70 | 25 | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 | | | |
| mple Variation 12 Vw.Luse Forest Copens | each subplement of the subplem | Land <50% ground >75% ground ns, parks, etc. | Side Chment of t Runoff Score Use (Choos cover) cover) | he stream. If for waterships for materials are from Dro | 70 med: p List) | 25 No | Side | Runof Score 0.5 1 0.3 | 22 77 | Runni Perce (not >1 22 99 100 | | | |
| mple Variati 2 Vw.Luse Forest a Open s Variable | each subplement of the subplem | Left 10 a entire cate Average of F Land <50% ground >75% ground ns, parks, etc. | Side chment of t dunoff Score Use (Choose cover) cover) property of the cover | he stream. for watersh se From Dro >75% | p List) | No oleted using imagery an | tes: | Runof Score 0.5 1 0.3 9 Nationa supplement | ment 22 77 1 Land Cover tary datasets | Runni Perce (not >1 22 99 100 | | | |
| mple Variation 12 Vw.Luse Forest Copens | each subplement of the subplem | Land <50% ground >75% ground ns, parks, etc. | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| mple Variati 2 Vw.Luse Forest a Open s Variable | each subplement of the subplem | Left 10 a entire cate Average of F Land <50% ground >75% ground ns, parks, etc. | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets | Runni Perce (not >1 22 99 100 | | | |
| mple Variation 12 Vw.Luse Forest a Open s Variable Vccanopy Vembed | each subplement of the subplem | Land <50% ground >75% ground ns, parks, etc. VSI Not Used 0.34 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Forest and Forest and Popular State of the Popular | each subplement of the subplem | Left 10 Pe entire cate Average of F Land <50% ground >75% ground ns, parks, etc. VSI Not Used 0.34 0.04 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| mple Variation 12 Vw.Luse Forest a Open s Variable Vccanopy Vembed | each subplement of the subplem | Land <50% ground >75% ground ns, parks, etc. VSI Not Used 0.34 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Forest and Forest and Popular State of the Popular | each subplement of the subplem | Left 10 Pe entire cate Average of F Land <50% ground >75% ground ns, parks, etc. VSI Not Used 0.34 0.04 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest and Company Variable | each subplement of the subplem | VSI Not Used 0.34 0.04 0.79 0.00 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest of Company Variable | each subplement of the subplem | Left 10 e entire cate Average of F Land <50% ground >75% ground ns, parks, etc. VSI Not Used 0.34 0.04 0.79 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest and Company Variable | each subplement of the subplem | VSI Not Used 0.34 0.04 0.79 0.00 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Variable Vacanopy Vember Vsubstrat Vbero Vtobh Vsnag | each subplement of the state of | VSI Not Used 0.04 0.79 0.00 Not Used | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Variable | each subplement of the search subplement of th | VSI Not Used 0.10 Not Used 0.10 1.00 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest and Company Variable | each subplement of the search subplement of th | VSI Not Used 0.04 0.79 0.00 Not Used 0.10 1.00 0.00 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest of Company Variable | seach subplement of the seach | VSI Not Used 0.04 0.79 0.00 Not Used 0.10 1.00 0.09 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |
| Porest and Company Variable | each subplement of the search subplement of th | VSI Not Used 0.04 0.79 0.00 Not Used 0.10 1.00 0.00 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Perce (not >10 22 99 100 | | | |
| Porest of Company Variable | seach subplement of the seach | VSI Not Used 0.04 0.79 0.00 Not Used 0.10 1.00 0.09 | Side Chment of t Runoff Score Use (Choose cover) cover) cover) Land Cove (NLCD), fr Watershee | he stream. for watersh se From Dro >75% er Analysis rom Lands, d boundari | p List) | No oleted using imagery an ed off of fie | tes: g the 201 d other s | Runof Score 0.5 1 0.3 9 Nationa supplementated streated streat | ment 22 77 1 Land Cover tary datasets m impacts. | Runni Perce (not >1 22 99 100 | | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME S-MM15 | | LOCATION Montgomery Co | punty | | | | | | | |
|----------------------------|----------------------------|--------------------------------|--|-------------------|--|--|--|--|--|--|
| STATION #_12077+00 R | IVERMILE | STREAM CLASS Intermitten | t | | | | | | | |
| LAT 37.258673 LC | ONG -80.296446 | RIVER BASIN Upper Roan | RIVER BASIN Upper Roanoke | | | | | | | |
| STORET# | | AGENCY VADEQ | | | | | | | | |
| INVESTIGATORS AO, MI | M | | | | | | | | | |
| FORM COMPLETED BY | AO, MM | DATE 8/25/2021 TIME 9:15 AM | REASON FOR SURVEY Bas | seline Assessment | | | | | | |
| WEATHER CONDITIONS | rain (shower | (heavy rain) (steady rain) | Has there been a heavy rain in Yes √No Air Temperature 24 0 C Other | the last 7 days? | | | | | | |
| SITE LOCATION/MAP | (omino) | RB FQUIPME Bridge | ROW ROW LIS | Going | | | | | | |
| STREAM CHARACTERIZATION | Stream Subsystem Perennial | ermittent Tidal | Stream Type | | | | | | | |

Note: No water present.

Spring-fed

Mixture of origins

Other Precipitation

Catchment Area 0.13

 km^2

Stream Origin
Glacial
Non-glacial montane
Swamp and bog

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERS FEATURI | | ✓ Fores | Pasture Industri | rcial | Local Watershed NPS ☑ No evidence ☐ Son ☐ Obvious sources Local Watershed Erosi ☑ None ☐ Moderate | ne potential sources | | | |
|--------------------------------|---------------------------|--|---|---------------------------------|--|--|--|--|--|
| RIPARIA VEGETA (18 meter | TION | | e the dominant type and | | ominant species present Grasses He | rbaceous | | | |
| INSTREA FEATURI | | Estimat Samplin Area in Estimat | ted Stream Depth NA MA | m m² km² | Canopy Cover ☐ Partly open ☐ Part High Water Mark ☐ Proportion of Reach R Morphology Types Riffle | | | | |
| LARGE V DEBRIS | VOODY | LWD Density | of LWD NA m | n²/km² (LWD / | reach area) | | | | |
| AQUATIC VEGETA | | Roote Floati | e the dominant type and ed emergent RA ng Algae At ant species present AA of the reach with aquat | ooted submerge ttached Algae | ent Rooted floating | ☐Free floating | | | |
| WATER ((DS, US) | QUALITY | Specific Dissolve pH NA Turbidi | cature NA 0 C c Conductance NA ed Oxygen NA strument Used NA | | Petroleum Fishy Water Surface Oils Slick Sheen None Other | Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Sheen Globs Flecks None Other Turbidity (if not measured) | | | |
| SEDIMEN SUBSTRA | | Odors Norm Chem Other Oils Absen | ☐Paper fiber ☐Sand]Other h are not deeply embedded, k in color? | | | | | | |
| INC | ORGANIC SUBS (should a | STRATE dd up to 1 | COMPONENTS | | ORGANIC SUBSTRATE C | | | | |
| Substrate Type | Diamet | er | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | | | |
| Bedrock Boulder | > 256 mm (10") |) | 1 0 | Detritus | sticks, wood, coarse plant materials (CPOM) | 75 | | | |
| Cobble | 64-256 mm (2.5 | | 5 | Muck-Mud | black, very fine organic (FPOM) | 0 | | | |
| Gravel 2-64 mm (0.1"-2.5") | | | 54 | | | | | | |
| Sand | 0.06-2mm (gritt | y) | 10 | Marl | grey, shell fragments | | | | |
| Silt | 0.004-0.06 mm | | 15 | | | | | | |
| Clav | < 0.004 mm (sli | ck) | 15 | | I | | | | |

Note: No water present.

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME S-MM15 | LOCATION Montgomery County | | | | | |
|---|---|--|--|--|--|--|
| STATION #_12077+00 RIVERMILE | STREAM CLASS Intermittent | | | | | |
| LAT <u>37.258673</u> LONG <u>-80.296446</u> | RIVER BASIN Upper Roanoke | | | | | |
| STORET# | AGENCY VADEQ | | | | | |
| INVESTIGATORS AO, MM | | | | | | |
| FORM COMPLETED BY AO, MM | DATE 8/25/2021 TIME 9:15 AM PM REASON FOR SURVEY Baseline Assessment | | | | | |

| | Habitat | | Condition | Category | | | | |
|--|---|---|---|---|---|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | | |
| | SCORE 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | | |
| ted in | SCORE 11 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | | | |
| ıram | SCORE 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| P ₂ | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | | |
| | SCORE 8 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | | | |
| | score 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | |

Notes: No water present.

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| | Habitat | Condition Category | | | | | | | | | | | | |
|--|--|--|--|--|---|--|--|--|--|--|--|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | | | | | | | |
| | 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | | | | | | | |
| | SCORE 15 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | | | | | | |
| ling reach | 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | | | | | | | |
| amb | score 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | | | | | | |
| Parameters to be evaluated broader than sampling reach | 8. Bank Stability (score each bank) Note: determine left or right side by facing development. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | | | | | | | |
| e eva | SCORE 7 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |
| to be | SCORE 7 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |
| Parameters | 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | | | | | | | |
| | SCORE 7 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |
| | SCORE 7 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |
| | 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | | | | | | | | |
| | SCORE 8 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |
| | SCORE 9 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | | | | | | |

Notes: No water present. Erosion observed at base of equipment bridge footer on left bank.

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME S-MM15 | | | | | | LOCATION Mon | LOCATION Montgomery County | | | | | | | | | |
|---|----------|--|-------|--------|-------|---|----------------------------|------------|------|-------|-----------------------|-------|------|------|------|-----|
| STATION # 12077+00 | _ R | IVE | RMI | LE_ | | STREAM CLASS | STREAM CLASS Intermittent | | | | | | | | | |
| LAT 37.258673 | L | ONG | -80.2 | 296446 | | RIVER BASIN U | pper | Roar | noke | | | | | | | |
| STORET# | | | | | | AGENCY VADEO |) | | | | | | | | | |
| INVESTIGATORS AC | | | | | | | | | I | TO. | NUMBER | | | | | |
| FORM COMPLETED | BY | Α | Ο, | ١ | //\ | DATE TIME 9:15 AM | | | R | EAS | SON FOR SURVEY Bas | selin | e As | sses | ssme | ent |
| HABITAT TYPES | \sqcup | Cob | ble_ | | % | tage of each habitat type p Snags % | Veget | nt ated | Bank | | % | _% | | | | |
| SAMPLE COLLECTION | G | ear ı | ısed | | D-fr | ame kick-net | | Other | | | | | | | | |
| | Н | ow w | vere | the s | amp | oles collected? wadi | ng | | fron | ı ban | nk from boat | | | | | |
| | \sqcup | Indicate the number of jabs/kicks taken in each habitat type. Cobble Snags Submerged Macrophytes Other (Other (| | | | | | | | | | | | | | |
| GENERAL COMMENTS | N | o w | vate | er | pre | sent. | | | | | | | | | | |
| | | | | | | | | | | | | | | | | |
| QUALITATIVE I Indicate estimated Dominant | | | | | | ATIC BIOTA Absent/Not Observed, | 1 = ì | Rare | e, 2 | = C | ommon, 3= Abunda | ant, | 4 = | : | | |
| Periphyton | | | | | 0 | 1 2 3 4 | Sli | mes | | | | 0 | 1 | 2 | 3 | 4 |
| Filamentous Algae | | | | | 0 | 1 2 3 4 | Ma | croi | nver | tebr | rates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | | | | | 0 | 1 2 3 4 | Fis | h | | | | 0 | 1 | 2 | 3 | 4 |
| FIELD OBSERVA Indicate estimated | | | | : | 0 = . | ROBENTHOS Absent/Not Observed, anisms), 3= Abundant | | | | | | | | | s) | |
| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta Isopoda | 0 | 1 1 | 2 | 3 | 4 | Sialidae 0 Corydalidae 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | - | | · | Culcidae 0 | _ | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

County: Montgomery County Stream Name: UNT to Flatwoods Branch S-MM15 Stream ID:

03010101 HUC Code: Basin: Upper Roanoke

Survey Date: 8/25/2021 Surveyors: Type: AO, MM Representative

| | | | LE COUNT | | | • | |
|-------------|-------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cun |
| | Silt/Clay | < .062 | S/C | A | 27 | 27.00 | 27.00 |
| | Very Fine | .062125 | | 4 | 2 | 2.00 | 29.00 |
| | Fine | .12525 | | • | 2 | 2.00 | 31.00 |
| | Medium | .255 | SAND | 4 | 0 | 0.00 | 31.00 |
| | Coarse | .50-1.0 | 7 | A | 3 | 3.00 | 34.00 |
| .0408 | Very Coarse | 1.0-2 | 1 | 4 | 1 | 1.00 | 35.00 |
| .0816 | Very Fine | 2 -4 | | A | 2 | 2.00 | 37.00 |
| .1622 | Fine | 4 -5.7 | 1 | A | 0 | 0.00 | 37.00 |
| .2231 | Fine | 5.7 - 8 | 1 | 4 | 3 | 3.00 | 40.00 |
| .3144 | Medium | 8 -11.3 | | A | 9 | 9.00 | 49.00 |
| .4463 | Medium | 11.3 - 16 | GRAVEL | A | 15 | 15.00 | 64.00 |
| .6389 | Coarse | 16 -22.6 | 1 | A | 6 | 6.00 | 70.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | - | A | 6 | 6.00 | 76.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | 1 | A | 5 | 5.00 | 81.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | 1 | A | 7 | 7.00 | 88.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | A | 7 | 7.00 | 95.00 |
| 3.5 - 5.0 | Small | 90 - 128 | 1 | A | 1 | 1.00 | 96.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | A | 1 | 1.00 | 97.00 |
| 7.1 - 10.1 | Large | 180 - 256 | 1 | ^ | 1 | 1.00 | 98.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | A | 0 | 0.00 | 98.00 |
| 14.3 - 20 | Small | 362 - 512 | 1 | A | 0 | 0.00 | 98.00 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | A | 0 | 0.00 | 98.00 |
| 40 - 80 | Large | 1024 -2048 | 1 | A | 0 | 0.00 | 98.00 |
| 80 - 160 | Vry Large | 2048 -4096 | 1 | A | 0 | 0.00 | 98.00 |
| | Bedrock | | BDRK | A | 2 | 2.00 | 100.00 |
| | | | | Totals | 100 | | |

RIVERMORPH PARTICLE SUMMARY

UNT to Flatwoods Branch

S-MM15

River Name: Reach Name: Sample Name: Sample Name: Representative Survey Date: 08/25/2021

| Size (mm) | TOT # | ITEM % | CUM % |
|---|---|--|--|
| 0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock | 27 2 2 0 3 1 2 0 3 9 15 6 6 5 7 7 1 1 1 0 0 0 0 | 27.00 2.00 2.00 0.00 3.00 1.00 2.00 0.00 3.00 9.00 15.00 6.00 6.00 7.00 7.00 7.00 1.00 1.00 0.00 0.00 0.00 0.00 0.00 | 27.00 29.00 31.00 31.00 34.00 35.00 37.00 40.00 49.00 64.00 70.00 76.00 81.00 88.00 95.00 96.00 97.00 98.00 98.00 98.00 98.00 98.00 98.00 98.00 |
| D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%) | 0.04 2 11.61 53.14 90 Bedrock 27 8 53 10 0 | | |

Total Particles = 100.

| | | | Strear | Unified St | tream Method | lology for use | in Virginia | | ' / | | |
|--|--|--|--|--|--|---|---|---|---|--|------------|
| | | | | | cowardin | ssified as interm | | | Impact | Impact | |
| Project # | Project Name (Applicant) | | | Locality | Class. | HUC | Date | SAR# | Length | Factor | |
| 22865.06 | | alley Pipeline ey Pipeline, L | | Montgomery County | R4 | 03010101 | 8/25/2021 | S-MM15 | 82 | 1 | |
| Name | e(s) of Evaluate | or(s) | Stream Name | e and Informa | tion | | | | SAR Length | | |
| | AO, MM | | UNT to Flatw | oods Branch | | | | | 82 | | |
| Channel C | ondition: Asses | ss the cross-secti | on of the stream a | | | | | | | | |
| | Optii | mal | Subo | ptimal | Conditional Catego Mar | ginal | Po | or | Sev | ere | |
| Channel | Very little incision or active erosion; 80-100% stable banks. Vegetative surface erosion or unprote | | Slightly incised, few areas of active erosion or unprotected banks. Majority Po | | | n incised, but less than Severe or Banks more stable than Severe | | sised. Vertically / e. Likely to widen both banks are near | Deeply incised vertical/lateral in incision, flow contain | stability. Severe | |
| Condition | (80-100%). AND/OR bankfull benches are to their original fit developed wide bank channel bars and tra Transient sediment less than 10% | Stable point bars / e present. Access codplain or fully (full benches. Mid- ansverse bars few. deposition covers | Vegetative protec prominent (60 Depositional feat stability. The bar channels are well di- has access to be newly developed portions of the r sediment covers 1 | tition or natural rock -80%) AND/OR tures contribute to nkfull and low flow efined. Stream likely ankfull benches, or I floodplains along reach. Transient 0-40% of the stream tom. | Erosion may be pr both banks. Vege 40-60% of banks. S vertical or und 40-60% Sediment transient, control Deposition that cc may be forming/p shaped channel protection on > 40 depositional featur | esent on 40-60% of tative protection on treambanks may be ercut. AND/OR may be temporary / ibute instability, ntribute to stability, resent. AND/OR V- shave vegetative % of the banks and res which contribute ability. | vertical. Erosion pr banks. Vegetative on 20-40% of bank: to prevent erosion. the stream is cove Sediment is temp nature, and contril AND/OR V-shap vegetative protect | esent on 60-80% of protection present s, and is insufficient AND/OR 60-80% of ered by sediment. orary / transient in puting to instability. eed channels have ion is present on > und stable sediment | Streambed below av majority of banks Vegetative protecti than 20% of banks erosion. Obviou present. Erosion/raw AND/OR Aggradin | vertical/undercut. on present on less i, is not preventing s bank sloughing v banks on 80-100%. g channel. Greater i bed is covered by uting to instability. channels and/or | CI |
| Scores | 3 | | 2 | 4 | | 2 | 1. | .6 | 1 | | 2.40 |
| | | | | | | | | | | | |
| RIPARIAN | BUFFERS: As | ssess both bank's | · | • | , , | measurements of | length & width ma | ay be acceptable) | NOTES: | | |
| . RIPARIAN | | | Cor | nditional Cate | gory | | | , , | NOTES>> | | |
| Riparian Buffers | Option Tree stratum (dbh > with > 60% tree Wetlands located warea | mal 3 inches) present, canopy cover. within the riparian | Cor Subo High Suboptimal: Riparian areas with | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover | gory Mar | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shruh and tree stratum, hay production, ponds, open water. If present, tree | | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | NOTES>> | | |
| Riparian Buffers | Optin Tree stratum (dbh > with > 60% tree Wetlands located v area | mal 3 inches) present, canopy cover. within the riparian as. | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. | nditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with 30% tree canopy cover with maintained understory. Low | High Poor: Laws, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | NOTES>> | | |
| Riparian | Optin Tree stratum (dbh > with > 60% tree Wetlands located v | mal 3 inches) present, canopy cover. within the riparian as. | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. | nditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | NOTES>> | | |
| Riparian Buffers Scores Delineate ripa Determine squ | Option Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 | 3 inches) present, canopy cover. within the riparian as. | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 | NOTES>> | | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Option Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 | 3 inches) present, canopy cover. within the riparian as. | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 | NOTES>> | | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 rian areas along eauare footage for eacuparian Area and S | 3 inches) present, canopy cover, within the riparian as. 5 ach stream bank to by measuring core for each ripar | Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cateror estimating lengarian category in the subset of | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 Logories and Cond th and width. Calc | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 | | | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 Trian areas along eacuare footage for eacuiparian Area and St % Riparian Area> Score > | 3 inches) present, canopy cover, within the riparian as. 5 ach stream bank ch by measuring core for each ripa 45% 0.6 | Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat or estimating lenguarian category in the 30% 0.85 | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 Logories and Cond (th and width. Calc | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums ciparian qual 100 100% | CI= (Sum % RA * Sc | | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank | Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 Trian areas along eacher footage for eacher footage f | 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank och by measuring core for each ripa 45% 0.6 | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat or estimating lenguarian category in the 30% 0.85 | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 Low 1.1 Low 1.1 Low 1.1 Low 1.1 Low 1.1 | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia, hay production, ponds, open water. If present, tree stratum, (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 | CI= (Sum % RA * Sc Rt Bank CI > | 0.85 | CI 0.82 |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN | Tree stratum (dbh > with > 60% tree Wetlands located warea areas along each are footage for each iparian Area and St % Riparian Area > Score > 1 HABITAT: Varian Area > Score > 1 HABITAT: | 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank to the by measuring core for each riparian 45% 0.6 75% 0.85 | High Suboptimal: Riparian areas with tree stratum (dbb > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat or estimating lenguarian category in the 30% 0.85 | nditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Cond th and width. Cale the blocks below. 20% 1.5 | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below. | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure t of % F Blocks e | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums tiparian qual 100 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > | 0.85 0.78 | CI 0.82 |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN | Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 rian areas along ea uare footage for eac iparian Area and Sc % Riparian Area > Score > 1 HABITAT: Varie e features. | 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank och by measuring core for each riparian 45% 0.6 75% 0.85 | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cate or estimating lenguarian category in the 30% 0.85 20% 0.6 ess, water velocity and suboptimal category in the su | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutver (dense vegetation). Low 1.1 egories and Cond (th and width. Calche blocks below. 20% 1.5 5% 0.5 and depths; woody | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 tition Scores using culators are provided to the control of the | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below. | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure to fine Blocks e | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums ciparian qual 100 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > | 0.85 0.78 | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank | Tree stratum (dbh > with > 60% tree Wetlands located warea areas along each are footage for each iparian Area and St % Riparian Area > Score > 1 HABITAT: Varian Area > Score > 1 HABITAT: | 3 inches) present, canopy cover. within the riparian as. 5 ach stream bank och by measuring core for each riparian 45% 0.6 75% 0.85 | Cor Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Cat or estimating lenguarian category in the 30% 0.85 20% 0.6 es, water velocity as Subo | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 Low 1.1 Low 1.1 Low 1.1 Low 0.5 And width. Calc 1.5 Low 1.5 Conditions 20% Conditions 20th 20 conditions 20th 20 conditions 20th 20 conditions 20th 20th 20th 20th 20th 20th 20th 20th | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 High 0.85 dulators are provided to the control of the control | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below. | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland, actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure to fine Blocks e | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums Exparian qual 100 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$ | 0.85 0.78 | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN omplexes, stable | Tree stratum (dbh > with > 60% tree Wetlands located warea 1.4 rian areas along ea uare footage for eac iparian Area and Sc % Riparian Area > Score > 1 HABITAT: Varie e features. | 3 inches) present, canopy cover, within the riparian as. 5 ach stream bank in the stream | Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 Into Condition Cat or estimating lenguarian category in the stratum of the st | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutver (dense vegetation). Low 1.1 egories and Cond (th and width. Calche blocks below. 20% 1.5 5% 0.5 and depths; woody | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using culators are provided to the control of the | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparia and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. ded for you below. | High Poor: Laws, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure to fine Blocks expensely we seed and stabilized, or other comparable condition. | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums Riparian qual 100 100% 1100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; \$ | 0.85 0.78 SAV; riffle/pool | |

| Stream Impact Assessment Form Page 2 | | | | | | | | | |
|--------------------------------------|--|----------------------|--------------------|----------|-----------|--------|------------------|------------------|--|
| Project # | Project Name (Applicant) | Locality | Cowardin Class. | HUC | Date | SAR# | Impact Length | Impact Factor | |
| 22865.06 | Mountain Valley Pipeline (Mountain Valley Pipeline, LLC) | Montgomery County | R4 | 03010101 | 8/25/2021 | S-MM15 | 82 | 1 | |
| 4. CHANNEL | 4. CHANNEL ALTERATION: Stream crossings, riprap, concrete, gabions, or concrete blocks, straightening of channel, channelization, embankments, spoil piles, constrictions, livestock | | | | | | | | |

Conditional Category NOTES>>

| | Negligible | Minor | | Moderate | | Severe | |
|-----------------------|--|-------------|-------------|--|---|--------|--|
| Channel Alteration | Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized. | the channel | the channel | of the channel alterations listed in the parameter guidelines. If | 60 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered. | | |
| Scores | 1.5 | 1.3 | 1.1 | 0.9 | 0.7 | 0.5 | |

CI 1.30

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

THE REA

THE REACH CONDITION INDEX (RCI) >>

1.14

RCI= (Sum of all Cl's)/5, except if stream is ephemeral RCI = (Riparian Cl/2)

 COMPENSATION REQUIREMENT (CR) >>
 93

 CR = RCI X L_I X IF

INSERT PHOTOS:


(WSSI Photo Location L:\22000s\22800\22865.06\Admin\05-ENVR\Field Data\Spread H\Field Forms\S-MM15\Photos)




 $\label{eq:constream} \mbox{Downstream view of ROW looking S. Assessment is limited to areas within the temporary ROW.}$


| DESCRIBE PROPOSED IMPACT | 1: |
|--------------------------|----|
|--------------------------|----|

PROVIDED UNDER SEPARATE COVER







| CL STAKEOUT POINTS: S-MM15 CROSS SECTION B (PIPE CL) | | | | | | |
|--|-------------|-------------|---------|--------|---------|--|
| | PF | RE-CROSSING | | POST-C | ROSSING | |
| DT LOC | NODTHING | FASTING | ELEV | VERT. | HORZ. | |
| PT. LOC. | NORTHING | EASTING | ELEV | DIFF. | DIFF. | |
| TS-L | 13529503.49 | 1845114.03 | 1764.00 | | | |
| BS-L | 13529501.46 | 1845104.87 | 1758.16 | | | |
| THW | 16529501.08 | 1845103.65 | 1757.72 | | | |
| BS-R | 13529500.05 | 1845099.60 | 1758.63 | | | |
| TS-R | 13529498.24 | 1845091.73 | 1762.52 | | | |





SURVEY NOTES:


- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on February 14, 2019.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).
- 4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.
- 5. All section views shown are left to right facing downstream.
- 6. Cross section B shot at location of pipe centerline (based on field stakes)











NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

CROSS SECTION LEGEND EXISTING GRADE



Wetland

to



PHOTO TAKEN LOOKING DOWNSTREAM TO THE SOUTH-SOUTHEAST ON 02/14/2019





PHOTO TAKEN LOOKING UPSTREAM TO THE NORTHWEST ON 02/14/2019

| PENDING CROSSING | |
|---------------------|--|
| PHOTO TAKEN LOOKING | |



CROSS SECTION

SCALE:

H: 1"=10'

V: 1"=5'

PHOTO TAKEN LOOKING

Approved NAS PFS JSF Sheet # 1 of 1

Horizontal Datum: NAD 1983 UTM ZONE 1

Computer File Name: :\Survey\22000s\22800\22865.03\Spread H Work Dwgs 2865_03 S-H MP 227-240 Sheets.dwg

POST-CROSSING PHOTOS

Vertical Datum: NAVD 88 Boundary and Topo Source: WSSI 2' C.I. Topo