#### **Baseline Assessment – Stream Attributes**

# Reach S-A38 (Timber Mat Crossing) Intermittent Spread I Franklin County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	✓
RBP Physical Characteristics Form	✓
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A – Lack of habitat
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	✓
Longitudinal Profile and Cross Sections	✓

#### Spread I Stream S-A38 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking NW upstream, RAH



Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking SE downstream, RAH

#### Spread I Stream S-A38 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking W at right streambank, RAH



Photo Type: RB CL

Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking SE at left streambank, RAH


#### Spread I Stream S-A38 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking NE upstream, RAH



Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking SE downstream, RAH



										Versi	on 10-20-17
			High-C	Gradient					a		
				Field [	Data She	et and C	alculato	r			
	Team:	RH, CL						Latitude/UT	M Northing:	37.036271	
Pro	•	Mountain V					Longitude/UTM Easting: -79.799442				
	Location:	Franklin Co	unty; Spread	11				Sar	npling Date:	8/26/2021	
SA	AR Number:	S-A38	Reach	Length (ft):	64	Stream Ty	/pe: Inter	mittent Strea	m		•
	Top Strata:	Sh	rub/Herb Str	ata	(determined	d from perce	nt calculate	d in V <sub>CCANOP</sub>	<sub>Y</sub> )		
Site	and Timing:	Project Site				_	Before Proje	ct			•
Sample	Variables	1-4 in strea	m channel								
1	V <sub>CCANOPY</sub>	equidistant	rcent cover points along st one value	the stream.	Measure o	nly if tree/sa	pling cover	is at least 20			Not Used, <20%
	List the per	cent cover n	neasuremen	ts at each p	oint below:						
	0	0	0	0	0	0	0	10	10	10	
				5.11	<u> </u>						
2	$V_{EMBED}$		nbeddednes tream. Sele								1.6
		-	l area surrou				-		-		
			ving table. If								
			bed is comp						,	9	
			ness rating f					from Platts	, Megahan,	and	
		Minshall 19	83)								
		Rating	Rating Des								1
		5		of surface c							
		4		ent of surfa							
		3		rcent of surf			,	,			
		2		rcent of surf					nent (or artificial	surface)	
	List the rati	ngs at each		t or surface	covered, 3d	rounded, or	buricu by iii	ic scaiment	(or artificial	Juliacc)	J
	1	1	1	1	1	1	1	3	2	3	ì
	2	1	1	4	4	3	1	1	1	1	
	1	4	3	1	1	1	1	1	1	1	
		7	3		'	'	'	'		'	
3	\/	Median stre	om channol	cubatrata n	ortiolo cizo	Monguro ot	no fower the	an 20 roughl	v oguidieten	t points	
J	▼ SUBSTRATE		ream; use th					an so rough	y equidistan	t points	0.25 in
	Enter partie	le size in inc						ould be sour	stad as 00 in	acabalt or	
		s 0.0 in, sand				Joint Delow (	pediock sile	Julu De Coul	ileu as 99 iii	, aspirali ui	
	0.08	0.08	0.08	0.08	0.08	0.20	0.30	0.30	0.40	4.10	ì
	2.20	0.08	0.08	0.40	0.40	0.20	0.08	0.08	3.00	0.80	
	1.60	1.40	1.10	0.40	0.40	0.50	0.30	0.20	1.30	0.00	
	1.00	1.40	1.10	0.00	0.00	0.50	0.50	0.20	1.50	0.00	
4	$V_{BERO}$	Total perce	nt of eroded	stream cha	nnel bank. I	Enter the tot	al number of	feet of eroo	ded bank on	each side	
			al percentage	e will be calc	ulated If bo	th banks are	e eroded, to	tal erosion f	or the strear	m may be	78 %
		up to 200%									
			Left Bank:	30	) ft		Right Bank:	20	) ft		
Sample	<b>Variables</b>	5-9 within t	he entire rip	oarian/buffe	r zone adja	cent to the	stream cha	nnel (25 fe	et from eac	h bank).	
5	$V_{LWD}$	Number of	down woody	stems (at le	east 4 inches	s in diameter	r and 36 incl	nes in lenath	) ner 100 fe	et of	
	* LWD		ch. Enter the								4.7
			stream will b						,		
					Number o	f downed wo	oody stems:		3		
6	$V_{TDBH}$		h of trees (n				cover is at	least 20%).	Trees are a	it least 4	Not Used
		inches (10	cm) in diame	eter. Enter t	ree DBHs in	inches.					Not Osed
		List the dbh	measureme	ents of indivi	dual trees (a	at least 4 in)	within the bi	uffer on eacl	h side of the		
		stream belo									
			Left Side					Right Side			
1											
1											
7	$V_{SNAG}$	Number of	snags (at lea	ast 4" dbh ar	nd 36" tall) p	er 100 feet	of stream. E	nter numbe	r of snags o	n each side	
1			m, and the a						5 -	_	0.0
			Left Side:		)		Right Side:		0		
8	$V_{SSD}$		saplings and								117.0
1			s <20%). Ei eam will be		or saplings	and shrubs (	on each side	or the strea	am, and the	arnount per	117.2
		100 11 01 511	Left Side:		5		Right Side:	2	20		

9	V <sub>SRICH</sub>	1 in the talle	est stratum.	Check all e		asive speci	n reach. Check ies present in all a.				1.56
		Grou	p 1 = 1.0					Group 2	2 (-1.0)		
	Acer rubrui	m		Magnolia tr	ipetala		Ailanthus altiss	sima		Lonicera ja	oonica
	Acer sacch	arum		Nyssa sylv	atica		Albizia julibriss	sin		Lonicera tat	tarica
I -	Aesculus fl	ava	П	Oxydendrum	arboreum		Alliaria petiolat	ta	Ē	Lotus cornic	culatus
l 🗀	Asimina tril			Prunus ser			Alternanthera			Lythrum sai	
Ιä	Betula alleg		_	Quercus ali			philoxeroides			Microstegium	
	Betula lenta			Quercus co			Aster tataricus			Paulownia t	
l	Carya alba			Quercus im			Cerastium fontanum				
1 -	•		_						_	Polygonum c	
□	Carya glab			Quercus pr			Coronilla varia			Pueraria mo	
	Carya oval			Quercus ru			Elaeagnus umbe			Rosa multifi	
	Carya ovat			Quercus ve			Lespedeza bic			Sorghum ha	
	Cornus flor			Sassafras a			Lespedeza cur			Verbena bra	asiliensis
	Fagus gran			Tilia amerio			Ligustrum obtus				
	Fraxinus ar	mericana		Tsuga cana	adensis		Ligustrum sine	ense			
✓	Liriodendron	tulipifera		Ulmus ame	ericana						
	Magnolia a	cuminata									
		1	Species in	Group 1				0	Species in	Group 2	
		'	Opooloo III	огоар т				-	Орсскоз пт	Oroup 2	
		bplots shou Average pe	Id be place rcent cover	d roughly e	quidistantly	<b>/ along eac</b> er organic m	in the riparian/l ch side of the s naterial. Woody er at each subpl	tream. debris <4			11.88 %
				Side			Right Si	ide		1 '	
		0	40	0 25		0			10		
11	$V_{HERB}$	woody stem	ns at least 4'	dbh and 36	6" tall. Becau	se there ma	sure only if tree ay be several lay cent cover of gr	yers of gr ound veg	ound cove	r vegetation	88 %
				Side			Right Si				
		100	60	100	75	100	100	80	90		
Sampl	e Variable 1	2 within the	entire cato	hment of the	he stream.						
Sample 12	e Variable 1				he stream.	ed:				<u> </u>	0.42
			verage of R	unoff Score					Runoff Score	% in Catch- ment	0.42  Running Percent
	V <sub>WLUSE</sub>	Weighted A	verage of R Land	unoff Score Use (Choos	for watershe				Score	ment	Running Percent (not >100)
	V <sub>WLUSE</sub>	Weighted A	verage of R Land 50% ground	Use (Choos	for watershe				Score 0.5	ment	Running Percent (not >100)
	V <sub>WLUSE</sub>	Weighted A	verage of R Land 50% ground	Use (Choos	for watershe			     	Score	ment	Running Percent (not >100)
	V <sub>WLUSE</sub> Forest and n	Weighted A	Land 50% ground 0% to 75% gr	Use (Choos	for watershe			\ \ \ \	Score 0.5	ment	Running Percent (not >100)
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			\ \ \ \	0.5 0.7	ment 11 36	Running Percent (not >100) 11 47
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A ative range (< ative range (5	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			\ \ \ \ \	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A ative range (< ative range (5	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			•	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A ative range (< ative range (5	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A ative range (< ative range (5	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			•	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
	VwLuse  Forest and n  Forest and n  Impervious a	Weighted A ative range (< ative range (5	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe			<b>~</b>	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
	Forest and n Forest and n Impervious a Open space	Weighted A ative range (< ative range (5 areas (parking (pasture, lawn	Land 50% ground 0% to 75% gr	Use (Choos cover) ound cover) iveways, etc)	for watershe		Notes	<b>* * * *</b>	0.5 0.7 0	ment 11 36 16	Running Percent (not >100) 11 47 63
12	Forest and n Forest and n Impervious a Open space	Weighted A ative range (< ative range (5 areas (parking (pasture, lawn	Land 50% ground 0% to 75% gr lots, roofs, dr s, parks, etc.),	Use (Choos cover) ound cover) iveways, etc) grass cover >	for watershe	D List)		<b>* * * *</b>	0.5 0.7 0 0.3	ment  11  36  16  37	Running Percent (not >100) 11 47 63 100
12 V	Forest and n Forest and n Impervious a Open space	Weighted A ative range (< ative range (5 areas (parking (pasture, lawn	Land 50% ground 0% to 75% gr lots, roofs, dr s, parks, etc.),	Use (Choos cover) ound cover) iveways, etc) grass cover 2	for watershe se From Drop -75% er Analysis	vas com	Notes: pleted using the imagery and c	▼	0.5 0.7 0 0.3	ment  11  36  16  37  and Cover I	Running Percent (not >100) 11 47 63 100
12 V	Forest and n Forest and n Impervious a Open space	Weighted A ative range (< ative range (5 areas (parking (pasture, lawn	Land 50% ground 0% to 75% gr lots, roofs, dr s, parks, etc.),	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12 V	Forest and n Forest and n Impervious a Open space	ative range (< ative range (5 areas (parking (pasture, lawn) S-A38 Value Not Used,	Land 50% ground 0% to 75% gr lots, roofs, dr s, parks, etc.),	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using th imagery and c	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12  VC Vc	Forest and n Forest and n Impervious a Open space	ative range (< ative range (5 areas (parking (pasture, lawn S-A38 Value Not Used, <20%	Land 50% ground 0% to 75% gr lots, roofs, dr s, parks, etc.),  VSI  Not Used	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12  Vo Ve Vs	Forest and n Forest and n Impervious a Open space  Zariable CANOPY  MBED  UBSTRATE	ative range (< ative range (5 areas (parking (pasture, lawn S-A38  Value  Not Used, <20%  1.6  0.25 in	VSI Not Used 0.33 0.13	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12  V Vc VE VS	Forest and n Forest and n Impervious a Open space  (ariable CANOPY MBED UBSTRATE	ative range (< ative range (5 areas (parking (pasture, lawn S-A38 Value Not Used, <20% 1.6 0.25 in 78 %	VSI Not Used 0.33 0.66	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12  Vo Ve Vs	Forest and n Forest and n Impervious a Open space  (ariable CANOPY MBED UBSTRATE	ative range (< ative range (5 areas (parking (pasture, lawn S-A38  Value  Not Used, <20%  1.6  0.25 in	VSI Not Used 0.33 0.13	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12 VC VE VS VB VL	Forest and n Forest and n Impervious a Open space  (ariable CANOPY MBED UBSTRATE	ative range (< ative range (5 areas (parking (pasture, lawn S-A38 Value Not Used, <20% 1.6 0.25 in 78 %	VSI Not Used 0.33 0.66	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12 V <sub>C</sub> V <sub>E</sub> V <sub>S</sub> V <sub>B</sub> V <sub>L</sub> V <sub>T</sub>	Forest and n Forest and n Impervious a Open space  Zariable  CANOPY  MBED  UBSTRATE  JERO  WD  DBH	ative range (< ative range (5 areas (parking (pasture, lawn  S-A38  Value  Not Used, <20%  1.6  0.25 in  78 %  4.7  Not Used	VSI Not Used 0.59 Not Used	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100) 11 47 63 100
12 V V <sub>C</sub> V <sub>E</sub> V <sub>S</sub> V <sub>B</sub> V <sub>L</sub> V <sub>T</sub> V <sub>S</sub>	Forest and n Forest and n Impervious a Open space  Cariable  CANOPY  MBED  UBSTRATE  JERO  WD  DBH  NAG	ative range (< ative range (5 areas (parking (pasture, lawn S-A38  Value Not Used, <20% 1.6 0.25 in 78 % 4.7 Not Used 0.0	VSI Not Used 0.59 Not Used 0.10	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100) 11 47 63 100
12 VCVEVS	Forest and n Forest and n Impervious a Open space  Zariable CANOPY MBED UBSTRATE EERO WD DBH NAG SD	ative range (< ative range (5 areas (parking (pasture, lawn  S-A38  Value  Not Used, <20%  1.6  0.25 in  78 %  4.7  Not Used	VSI Not Used 0.59 Not Used	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100) 11 47 63 100
12 VCVEVS	Forest and n Forest and n Impervious a Open space  Cariable  CANOPY  MBED  UBSTRATE  JERO  WD  DBH  NAG	ative range (< ative range (5 areas (parking (pasture, lawn S-A38  Value Not Used, <20% 1.6 0.25 in 78 % 4.7 Not Used 0.0	VSI Not Used 0.59 Not Used 0.10	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12 VC VE VS VS VS VD	Forest and n Forest and n Impervious a Open space  Gariable CANOPY MBED UBSTRATE JERO WD DBH NAG SD RICH JETRITUS	ative range (< ative range (5 areas (parking (pasture, lawn  S-A38  Value  Not Used, <20%  1.6  0.25 in  78 %  4.7  Not Used  0.0  117.2	VSI Not Used 0.59 Not Used 0.10 1.00	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database
12 VC VE VS VS VS VD	Forest and n Forest and n Impervious a Open space  Gariable CANOPY MBED UBSTRATE EERO WD DBH NAG SD RICH	ative range (< ative range (5 areas (parking (pasture, lawn  S-A38  Value  Not Used, <20%  1.6  0.25 in  78 %  4.7  Not Used  0.0  117.2  1.56	VSI Not Used 0.59 Not Used 0.10 1.00 0.74	Use (Choos cover) ound cover) iveways, etc) grass cover s	er Analysis rom Lands:	was compat satellite	pleted using the imagery and co sed off of field	▼ ▼ ▼ ■ E 2019 Nother supdelineate	0.5 0.7 0 0.3  National Lipplementa ed stream	ment  11  36  16  37  and Cover I ry datasets. impacts.	Running Percent (not >100)  11  47  63  100  Database

Ver. 10-20-17

#### FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the <u>UPPERMOST STRATUM</u> of the plant community is determined based on the calculated value for V<sub>CCANOPY</sub> (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

**Project Name:** Mountain Valley Pipeline **Location:** Franklin County; Spread I

Sampling Date: 8/26/2021 Project Site Before Project

Subclass for this SAR:

Intermittent Stream

Uppermost stratum present at this SAR: SAR number: S-A38

Shrub/Herb Strata

Functional Results Summary:

**Enter Results in Section A of the Mitigation Sufficiency Calculator** 

Function	Functional Capacity Index
Hydrology	0.40
Biogeochemical Cycling	0.30
Habitat	0.19

#### Variable Measure and Subindex Summary:

Variable	Name	Average Measure	Subindex
V <sub>CCANOPY</sub>	Percent canpoy over channel.	Not Used, <20%	Not Used
V <sub>EMBED</sub>	Average embeddedness of channel.	1.63	0.33
V <sub>SUBSTRATE</sub>	Median stream channel substrate particle size.	0.25	0.13
$V_{BERO}$	Total percent of eroded stream channel bank.	78.13	0.66
$V_{LWD}$	Number of down woody stems per 100 feet of stream.	4.69	0.59
V <sub>TDBH</sub>	Average dbh of trees.	Not Used	Not Used
V <sub>SNAG</sub>	Number of snags per 100 feet of stream.	0.00	0.10
V <sub>SSD</sub>	Number of saplings and shrubs per 100 feet of stream.	117.19	1.00
V <sub>SRICH</sub>	Riparian vegetation species richness.	1.56	0.74
V <sub>DETRITUS</sub>	Average percent cover of leaves, sticks, etc.	11.88	0.14
V <sub>HERB</sub>	Average percent cover of herbaceous vegetation.	88.13	1.00
V <sub>WLUSE</sub>	Weighted Average of Runoff Score for Catchment.	0.42	0.44

## PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME S-A38		LOCATION Franklin County						
	IVERMILE	_ STREAM CLASS Inter	rmittent					
LAT 37.036271 LO	ONG79.799442	RIVER BASIN Upper F	RIVER BASIN Upper Roanoke					
STORET#		AGENCY VADEQ						
INVESTIGATORSRH, C	L							
FORM COMPLETED BY	RH	DATE 8/26/21 TIME 0824	REASON FOR SURVEY Baseline Assessment					
WEATHER CONDITIONS	rai show	rm (heavy rain) n (steady rain) vers (intermittent) %cloud cover clear/sunny						
SITE LOCATION/MAP	Pipe (	)\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Dense Dense Downstream					

Spring-fed
Mixture of origins
Other

Stream Type Coldwater

Catchment Area 0.16

✓Warmwater

\_km<sup>2</sup>

Stream Subsystem
□ Perennial □ Intermittent □ Tidal

Stream Origin
Glacial
Non-glacial montane
Swamp and bog

STREAM CHARACTERIZATION

## PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Predon  ✓ Fores  ✓ Field  — Agric  — Resid	Pasture Industri	rcial	Local Watershed NPS Pollution  ☑ No evidence ☐ Some potential sources ☐ Obvious sources  — Local Watershed Erosion ☑ None ☐ Moderate ☐ Heavy			
RIPARIA VEGETA (18 meter	TION	Trees	e the dominant type and s	hrubs	ominant species present ☐ Grasses ☐ Ho	erbaceous		
INSTREA FEATURI		Estimat Samplin Area in Estimat		m m² km² m	Canopy Cover  ☐ Partly open ☐ Partly shaded ☐ Shaded  High Water Mark ○2 m  Proportion of Reach Represented by Stream Morphology Types Riffle 55			
LARGE WOODY DEBRIS  LWD 6 m² Density of LWD 3 m²/km² (LWD/ reach area)								
AQUATIC VEGETATION  Indicate the dominant type and record the dominant species present Rooted submergent Rooted floating Free floating Algae  Dominant species present Dicanthellum sp.  Portion of the reach with aquatic vegetation 8 9%								
WATER (	QUALITY	Specific Dissolve pH 5.66 (c) Turbidi				Chemical Other  Globs Flecks		
SEDIMEN SUBSTRA		Odors Norm Chem Other Oils	nical Anaerobic		Paper fiber Sand Other NA ch are not deeply embedded, ck in color?			
INC		STRATE dd up to 1	COMPONENTS (00%)		ORGANIC SUBSTRATE C (does not necessarily add			
Substrate Type	Diamet	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area		
Bedrock Boulder	> 256 mm (10")	1	5	Detritus	sticks, wood, coarse plant materials (CPOM)	15		
Cobble Gravel	64-256 mm (2.5 2-64 mm (0.1"-2		15 20	Muck-Mud	black, very fine organic (FPOM)			
Sand Silt Clay	0.06-2mm (gritt 0.004-0.06 mm < 0.004 mm (sli		40 10 10	Marl	grey, shell fragments			

#### HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME S-A38	LOCATION Franklin County				
STATION # RIVERMILE	STREAM CLASS Intermittent				
LAT <u>37.036271</u> LONG <u>-79.799442</u>	RIVER BASIN Upper Roanoke				
STORET#	AGENCY VADEQ				
INVESTIGATORS RH, CL					
FORM COMPLETED BY RH	DATE 8/26/21 REASON FOR SURVEY TIME 6824 AM PM Baseline Assessment				

	Habitat		Condition	ı Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor			
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.			
	SCORE 10 ▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.			
ed ir	SCORE 8 ▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/depth regime (usually slow-deep).			
ıram	<sub>SCORE</sub> 8 <b>▼</b>	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
P <sub>2</sub>	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.			
	SCORE 14▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.			
	SCORE 2	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			

#### HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat	Condition Category											
	Parameter	Optimal	Suboptimal	Marginal	Poor								
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.								
	SCORE 18▼	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0								
ling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.								
ampl	SCORE 2	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0								
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank)  Note: determine left or right side by facing decumetrs.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion.	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.								
e eva	SCORE 5	Left Bank 10 9	8 7 6	5 4 3	2 1 0								
to b	SCORE 3	Right Bank 10 9	8 7 6	5 4 3	2 1 0								
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.								
	SCORE 9	Left Bank 10 9	8 7 6	5 4 3	2 1 0								
	SCORE 9 ▼	Right Bank 10 9	8 7 6	5 4 3	2 1 0								
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.								
	SCORE 8	Left Bank 10 9	8 7 6	5 4 3	2 1 0								
	SCORE 8 ▼)	Right Bank 10 9	8 7 6	5 4 3	2 1 0								

Total Score 104

#### BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME S-A38						L	LOCATION Franklin County												
STATION#	R	IVE	RMI	LE_		S	TRE	AM CLA	ASS II	nterr	nitte	nt							
LAT 37.036271	_ L	ONC	j -79.	79944	2	R	IVEF	R BASIN	√Upp	er F	Roar	oke							
STORET#						А	GEN	ICY VAI	DEQ										
INVESTIGATORS R	H, C	L					LOT NUMBER												
FORM COMPLETED	BY	R	Η				ATE IME		REASONT OR SORVET						ssm	ent			
HABITAT TYPES		Cob	ble	-	%	tage of eac Snags	;	bitat ty	'nΪν	eget	ated	Banl	ks	%	_%				
SAMPLE	G	Gear used D-frame kick-net Other																	
COLLECTION																			
	н	ow v	vere	tne	samp	oles collect	ea?	П,	vadin	g	_	Iron	n bar	k from boar	Į.				
		Cob	ble			r of jabs/k ☐ Snags phytes	;	taken in —	$\square$ V	hak eget	ated	Banl	ks	Sand	_				
GENERAL COMMENTS	ш	o b ate		thi	CS,	strean	n is	very	na	rro	)W	and	d c	ontains very lit	ttle	flov	wir	ıg	
Periphyton Filamentous Algae					-	1 2 1 2	3	-			nes		rtehr	ates		1 1	_	-	4 4
Macrophytes						1 2				Fish		11 V C	icoi	ates		1		3	
	l abı	und	anc	e:	0 = org	Absent/N anisms),	Not ( 3= A	Observ Abunda	nt (	>10	org	anis	sms)	rganisms), 2 = Con , 4 = Dominant (>5	50 oı	gar	ism		
Porifera	0	1			4	_			0	1				Chironomidae	0	1	2	3	
Hydrozoa	0	1	2	3	4	Zygopto			0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemipt			0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleop			0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepido		Į.	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidad			0	1	2	3	4						
Isopoda	0	1	2	3	4	Coryda		;	0	1	2	3	4						
Amphipoda	0	l	2	3	4	Tipulid			0	1	2	3	4						
Decapoda	0	l	2	3	4	Empidi			0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simulii			0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinid			0	1	2	3	4						
						Culcida	ıc		U			<u>)</u>	4						

#### WOLMAN PEBBLE COUNT FORM

County: Franklin County Stream ID: S-A38

Stream Name: UNT to Foul Ground Creek

HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 8/26/2021
Surveyors: RH, CL
Type: Representative

Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cun
	Silt/Clay	< .062	S/C	<b>A</b>	21	21.00	21.00
	Very Fine	.062125		<b>A</b>	11	11.00	32.00
	Fine	.12525	_	<b>A</b>	13	13.00	45.00
	Medium	.255	SAND	▼	8	8.00	53.00
	Coarse	.50-1.0		▼	2	2.00	55.00
.0408	Very Coarse	1.0-2		<b>*</b>	5	5.00	60.00
.0816	Very Fine	2 -4		▼	2	2.00	62.00
.1622	Fine	4 -5.7		▼	5	5.00	67.00
.2231	Fine	5.7 - 8	_	<b>~</b>	4	4.00	71.00
.3144	Medium	8 -11.3		<b>▲</b>	7	7.00	78.00
.4463	Medium	11.3 - 16	GRAVEL	<b>*</b>	6	6.00	84.00
.6389	Coarse	16 -22.6	]	<b>A</b>	6	6.00	90.00
.89 - 1.26	Coarse	22.6 - 32	]	<b>▲</b>	3	3.00	93.00
1.26 - 1.77	Vry Coarse	32 - 45		<b>*</b>	1	1.00	94.00
1.77 -2.5	Vry Coarse	45 - 64	]	<b>A</b>		0.00	94.00
2.5 - 3.5	Small	64 - 90		<b>A</b>	1	1.00	95.00
3.5 - 5.0	Small	90 - 128	COBBLE	<b>△</b>	3	3.00	98.00
5.0 - 7.1	Large	128 - 180	CORRLE	<b>A</b>	1	1.00	99.00
7.1 - 10.1	Large	180 - 256	]	<b>▲</b>		0.00	99.00
10.1 - 14.3	Small	256 - 362		<b>▲</b>	1	1.00	100.00
14.3 - 20	Small	362 - 512	1	<b>▲</b>		0.00	100.00
20 - 40	Medium	512 - 1024	BOULDER	<b>A</b>		0.00	100.00
40 - 80	Large	1024 -2048	1	<b>▲</b>		0.00	100.0
80 - 160	Vry Large	2048 -4096	1	<b>▲</b>		0.00	100.0
	Bedrock		BDRK	<b>A</b>		0.00	100.0
				Totals:	100		

#### RIVERMORPH PARTICLE SUMMARY

River Name: UNT to Foul Ground Creek Reach Name: S-A38 Sample Name: Representative 08/26/2021

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	21 11 13 8 2 5 2 5 4 7 6 6 6 3 1 0 1 3 1 0 0 0 0	21.00 11.00 13.00 8.00 2.00 5.00 2.00 5.00 4.00 7.00 6.00 6.00 3.00 1.00 0.00 1.00 0.00 1.00 0.00 0.00 0.00 0.00 0.00 0.00	21.00 32.00 45.00 53.00 55.00 60.00 62.00 67.00 71.00 78.00 84.00 90.00 93.00 94.00 94.00 95.00 98.00 99.00 100.00 100.00 100.00 100.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.05 0.15 0.41 16 90 361.99 21 39 34 5		

Total Particles = 100.

#### **Stream Assessment Form (Form 1)** Unified Stream Methodology for use in Virginia For use in wadeable channels classified as intermittent or perennial Cowardin **Impact Impact Project # Project Name (Applicant)** HUC SAR# Locality Date Length **Factor** Class. **Mountain Valley Pipeline (Mountain Franklin** S-A38 8/26/2021 30 22865.06 R4 03010101 **Valley Pipeline, LLC)** County SAR Length Name(s) of Evaluator(s) Stream Name and Information RH, CL S-A38 64 1. Channel Condition: Assess the cross-section of the stream and prevailing condition (erosion, aggradation) **Conditional Category Optimal Suboptimal Poor** Severe Marginal 100% stable banks. Vegetative Slightly incised, few areas of active Often incised, but less than Severe or laterally unstable. Likely to widen vertical/lateral instability. Severe surface protection or natural rock, erosion or unprotected banks. Majority Poor. Banks more stable than Severe further. Majority of both banks are incision, flow contained within the prominent (80-100%). AND/OR Stable near vertical. Erosion present on 60of banks are stable (60-80%). Channel or Poor due to lower bank slopes. banks. Streambed below average Erosion may be present on 40-60% of point bars / Vegetative protection or natural rock 80% of rooting depth, Condition bankfull benches are present. Access prominent (60-80%) AND/OR both banks. Vegetative protection on banks. Vegetative protection presen majority of banks vertical/undercut. Depositional features contribute to to their original floodplain or fully 40-60% of banks. Streambanks may on 20-40% of banks, and is Vegetative protection present on less developed wide bankfull benches. stability. The bankfull and low flow be vertical or undercut. AND/OR insufficient to prevent erosion. than 20% of banks, is not preventing channels are well defined. Stream channel bars and transverse bars few. 40-60% Sediment may be temporary. the stream is covered by sediment. erosion. Obvious bank sloughing Transient sediment deposition covers present. Erosion/raw banks on 80likely has access to bankfull transient, contribute instability. Sediment is temporary / transient in 100%. AND/OR Aggrading channel. less than 10% of bottom. Deposition that contribute to stability, nature, and contributing to instability benches, or newly developed portions of the reach. Transient than 80% of stream bed is covered by may be forming/present. AND/OR V-AND/OR V-shaped channels have sediment covers 10-40% of the shaped channels have vegetative vegetative protection is present on > deposition, contributing to instability. protection on > 40% of the banks and 40% of the banks and stable sediment Multiple thread channels and/or stream bottom. depositional features which contribute CI to stability. 3 2.4 2.40 2 1.6 **Scores** Assessment is limited to areas within the temporary ROW. NOTES>> RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) **Conditional Category** NOTES>> **Suboptimal Optimal** Marginal **Poor** Low Marginal: **High Poor:** \_ow Suboptimal: Non-maintained, **High Suboptimal** Lawns, mowed, Riparian areas **High Marginal:** dense herbaceous Riparian areas and maintained Low Poor: with tree stratum Non-maintained, vegetation, with tree stratum Impervious areas, nurseries; (dbh > 3 inches) dense herbaceous riparian areas (dbh > 3 inches) no-till cropland; surfaces, mine acking shrub and present, with 30% vegetation with Tree stratum (dbh > 3 inches) present present, with 30% actively grazed spoil lands, Riparian tree stratum, hay to 60% tree either a shrub with > 60% tree canopy cover. to 60% tree denuded surfaces pasture, sparsely production, ponds layer or a tree canopy cover and **Buffers** Wetlands located within the riparian canopy cover and vegetated nonrow crops, active a maintained layer (dbh > 3 open water. If feed lots, trails, or containing both maintained area, areas. inches) present, understory. present, tree recently seeded herbaceous and other comparable stratum (dbh >3 Recent cutover with <30% tree and stabilized, or shrub layers or a conditions. canopy cover. inches) present, (dense non-maintained other comparable with <30% tree vegetation). condition. understory. canopy cover with maintained understory. High High High Low Low Low 1.5 1.2 1.1 0.85 0.75 0.6 0.5 Scores Delineate riparian areas along each stream bank into Condition Categories and Condition Scores using the Ensure the sums descriptors. 2. Determine square footage for each by measuring or estimating length and width. Calculators are provided for you of % Riparian below. Blocks equal 100 3. Enter the % Riparian Area and Score for each riparian category in the blocks below. Assessment is limited to 10% 90% 100% % Riparian Area> areas within the **Right Bank** 0.6 0.85 Score > temporary ROW. CI= (Sum % RA \* Scores\*0.01)/2 10% 90% 100% CI % Riparian Area> Rt Bank CI > 0.83 **Left Bank** 0.6 0.85 0.83 Lt Bank CI > 0.83 Score > 3. INSTREAM HABITAT: Varied substrate sizes, water velocity and depths; woody and leafy debris; stable substrate; low embededness; shade; undercut banks; root mats; SAV; riffle/pool complexes, stable features. NOTES>> **Conditional Category** Marginal **Suboptimal Optimal Poor** Instream Habitat/ Stable habitat elements are typically Stable habitat elements are typically Habitat elements listed above are **Available** present in 10-30% of the reach and present in 30-50% of the reach and Habitat elements are typically present lacking or are unstable. Habitat Cover in greater than 50% of the reach. are adequate for maintenance of elements are typically present in less are adequate for maintenance of than 10% of the reach. populations. populations. **Stream Gradient** CI High / Low 1.5 1.2 0.9 0.5 0.90 Scores

Stream Impact Assessment Form Page 2									
Project #	Project Name (Applicant)		Locality	Cowardin Class.	HUC	Date	SAR#	Impact length	Impact Factor
22865.06	Mountain Valley Pipeline (Mountain Valley Pipeline, LLC)		Franklin County	R4	03010101	8/26/2021	S-A38	30	1
4. CHANNEL ALTERATION: Stream crossings, riprap, concrete, gabions, or concrete blocks, straightening of channel, channelization, embankments, spoil piles, constrictions, livestock  Conditional Category  NOTES>>									
	Negligible		Moderate				INO I LOPP		
	itogrigibio	IVI I	nor	Mode	erate	Sev	ere		

recovered.

0.7

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

recovered.

0.9

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

1.3

1.1

1.5

THE REACH CONDITION INDEX (RCI) >> 1.09

RCI= (Sum of all Cl's)/5, except if stream is ephemeral RCI = (Riparian Cl/2)

CI

1.30

COMPENSATION REQUIREMENT (CR) >> 33

 $CR = RCI X L_I X IF$ 

0.5


## **INSERT PHOTOS:**


**Scores** 




DESCRIBE PROPOSED IMPACT:

PROVIDED UNDER SEPARATE COVER







CL STAKEOUT POINTS: S-A38 CROSS SECTION B (PIPE CL)								
	PR	PRE-CROSSING						
PT. LOC.	NORTHING	EASTING	ELEV	VERT.	HORZ.			
P1. LOC.	NORTHING	EASTING	CLEV	DIFF.	DIFF.			
TS-L	13450013.20	1990684.58	1073.22					
BS-L	13450013.46	1990683.00	1069.81					
THW	13450013.55	1990681.89	1069.37					
BS-R	13450013.67	1990680.17	1070.22					
TS-R	13450015.57	1990664.58	1075.74					

TYPICAL 5-POINT CROSS-SECTION
(FACING DOWNSTREAM)


TS-L

TS-R

BS-L

THW

TS: TOP OF SLOPE
BS: BOTTOM OF SLOPE
THW: THALWEG (INVERT)



### SURVEY NOTES:

- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on February 19, 2019.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).
- 4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.
- 5. All section views shown are left to right facing downstream.
- 6. Cross-section B shot at location of pipe centerline (based on best professional judgement).



PHOTO TAKEN LOOKING DOWNSTREAM TO THE SOUTHEAST ON 02/19/2019



PHOTO TAKEN FROM BRIDGE LOOKING UPSTREAM TO THE NORTHWEST ON 02/19/2019

PENDING CROSSING

PHOTO TAKEN LOOKING

PENDING CROSSING

PHOTO TAKEN LOOKING

SN							SCALE: AS N
REVISIONS	Description						DATE: October 2021 SCA
	No. Date						TE: Oct
	No.						DAT
Horiz	zontal l	Dati	ım:	NAD :	1983 U	TM ZC	NE 17N

 $\infty$ 

Horizontal Datum: NAD 1983 UTM ZONE 177

Vertical Datum: NAVD 88

Boundary and Topo Source:

MVP

WSSI 2' C.I. Topo

Design Draft Approved

JSF NAS
Sheet #

1 of 1

Computer File Name:
L:\Survey\22000s\22800\22865.03\Spread I Work Dwgs
22865\_03 S-I MP 268-278 Sheets\_2.dwg

EJC