Baseline Assessment – Stream Attributes

Reach S-C19 (Pipeline ROW) Perennial Spread I Franklin County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope >4%)
RBP Physical Characteristics Form	✓
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A - No Riffles
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	✓
Longitudinal Profile and Cross Sections	✓



Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking SE downstream, RAH



Photo Type: LB DS VIEW

Location, Orientation, Photographer Initials: Upstream at ROW/LOD on left bank looking SW downstream, RAH



Location, Orientation, Photographer Initials: On ROW/LOD on left bank looking N upstream, RAH



Photo Type: LB CL

Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking NW at right streambank, RAH



Location, Orientation, Photographer Initials: Upstream at ROW/LOD on right bank looking SE downstream, RAH



Photo Type: RB US VIEW


Location, Orientation, Photographer Initials: Downstream a ROW/LOD on right bank looking NE upstream, RAH



Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking E at left streambank, RAH



Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking N upstream, RAH



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME	LOCATION					
STATION # RIVERMILE	STREAM CLASS					
LAT LONG	RIVER BASIN					
STORET#	AGENCY					
INVESTIGATORS						
FORM COMPLETED BY	DATE	REASON FOR SURVEY				

WEATHER CONDITIONS SITE LOCATION/MAP	Now Past 24 hours Yes No storm (heavy rain) rain (steady rain) showers (intermittent) %cloud cover clear/sunny Draw a map of the site and indicate the areas sampled (or attach a photograph)
	WY S PIPE CL
	ROW/BRIDGE
	LOD
STREAM CHARACTERIZATION	Stream Subsystem Perennial Intermittent Tidal Stream Type Coldwater Warmwater Stream Origin Glacial Spring-fed Non-glacial montane Mixture of origins Swamp and bog Other

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Fores Field/ Agric	Pasture Industria	rcial	Local Watershed NPS Pollution No evidence ☐ Some potential sources Obvious sources Local Watershed Erosion None Moderate Heavy	
RIPARIA VEGETA (18 meter	TION	Trees	SI SI	hrubs	Ominant species present Grasses Herbaceous	
INSTREA FEATURI		Estimat Estimat Samplin Area in Estimat	ed Reach Length ed Stream Width g Reach Area km² (m²x1000) ed Stream Depth Velocity m	m m m² km²	Canopy Cover Partly open Partly shaded Shaded High Water Markm Proportion of Reach Represented by Stream Morphology Types Riffle % Run% Pool% Channelized Yes No Dam Present Yes No	
LARGE V DEBRIS	VOODY		m²	n ² /km ² (LWD/	reach area)	
AQUATION VEGETA		Roote Floati Domin a	ominant species present ent Rooted floating Free floating			
WATER (QUALITY	Specific Dissolve pH Turbidi	cature0 C Conductance ed Oxygen ty trument Used		Water Odors Normal/None Sewage Petroleum Chemical Fishy Other	
SEDIMEN SUBSTRA		Odors Norm Chem Other Oils Abser			Relict shells Other	_
INC	ORGANIC SUBS		COMPONENTS 00%)		ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%)	
Substrate Type	Diamete	er	% Composition in Sampling Reach	Substrate Type	Characteristic % Composition in Sampling Area	
Bedrock Boulder	> 256 mm (10")			Detritus	sticks, wood, coarse plant materials (CPOM)	
Cobble Gravel	64-256 mm (2.5 2-64 mm (0.1"-2			Muck-Mud	black, very fine organic (FPOM)	
Sand	0.06-2mm (gritt	y)		Marl	grey, shell fragments	

Silt

Clay

0.004-0.06 mm

< 0.004 mm (slick)

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION	
STATION # RIVERMILE	STREAM CLASS	
LAT LONG	RIVER BASIN	
STORET#	AGENCY	
INVESTIGATORS		
FORM COMPLETED BY	DATE AM PM	REASON FOR SURVEY

	Habitat		Condition	ı Category						
	Parameter	Optimal	Suboptimal	Marginal	Poor					
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.					
ted in	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/depth regime (usually slow-deep).					
ıram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Pa	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	n Category						
	Parameter	Optimal	Suboptimal	Marginal	Poor					
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
oling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.					
samp	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing downstream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.					
e eva	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0					
to be	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0					
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.					
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0					
	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0					
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.					
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0					
ĺ	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0					

Total	Caama	
i otai	Score	

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME	LOCATION						
STATION # RIVERMILE	STREAM CLASS						
LAT LONG	RIVER BASIN						
STORET#	AGENCY	AGENCY					
INVESTIGATORS		LOT NUMBER					
FORM COMPLETED BY	DATETIME	REASON FOR SURVEY					
HADITAT TYPES Indicate the percentage of	and habitat type present						

HABITAT TYPES	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%
SAMPLE COLLECTION	Gear used D-frame kick-net Other How were the samples collected? wading from bank from boat
	Indicate the number of jabs/kicks taken in each habitat type. Cobble Snags Vegetated Banks Sand Submerged Macrophytes Other ()
GENERAL COMMENTS	

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3 = Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3 = Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

WOLMAN PEBBLE COUNT FORM

County: Franklin County Stream ID: S-C19

Stream Name:

Maggodee Creek 03010101 Upper Roanoke HUC Code: Basin:

Survey Date: 8/3/2021 Surveyors: RC, RH

Representative/Riffle Type:

			LE COUNT				
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cun
	Silt/Clay	< .062	S/C	A	32	32.00	32.00
	Very Fine	.062125		^	5	5.00	37.00
	Fine	.12525		~	9	9.00	46.00
	Medium	.255	SAND	^	7	7.00	53.00
	Coarse	.50-1.0		^	13	13.00	66.00
.0408	Very Coarse	1.0-2		•	16	16.00	82.00
.0816	Very Fine	2 -4		^		0.00	82.00
.1622	Fine	4 -5.7		A		0.00	82.00
.2231	Fine	5.7 - 8		A		0.00	82.00
.3144	Medium	8 -11.3		A	1	1.00	83.00
.4463	Medium	11.3 - 16	GRAVEL	A	1	1.00	84.00
.6389	Coarse	16 -22.6		A	2	2.00	86.00
.89 - 1.26	Coarse	22.6 - 32		^	1	1.00	87.00
1.26 - 1.77	Vry Coarse	32 - 45		^	4	4.00	91.00
1.77 -2.5	Vry Coarse	45 - 64		^	2	2.00	93.00
2.5 - 3.5	Small	64 - 90		^	4	4.00	97.00
3.5 - 5.0	Small	90 - 128		^	2	2.00	99.00
5.0 - 7.1	Large	128 - 180	COBBLE	^	1	1.00	100.0
7.1 - 10.1	Large	180 - 256		^		0.00	100.0
10.1 - 14.3	Small	256 - 362		A		0.00	100.0
14.3 - 20	Small	362 - 512	1	A		0.00	100.0
20 - 40	Medium	512 - 1024	BOULDER	A		0.00	100.0
40 - 80	Large	1024 -2048	1	A		0.00	100.0
80 - 160	Vry Large	2048 -4096	1	A		0.00	100.0
	Bedrock		BDRK	A		0.00	100.0
				Totals:	100		

RIVERMORPH PARTICLE SUMMARY

Maggodee Creek S-C19 Representative 08/31/2021 River Name: Reach Name: Sample Name: Survey Date:

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	32 5 9 7 13 16 0 0 0 1 1 1 2 1 4 2 4 2 1 0 0 0 0	32.00 5.00 9.00 7.00 13.00 16.00 0.00 0.00 1.00 1.00 2.00 1.00 4.00 2.00 4.00 2.00 1.00 0.00 0.00 0.00 0.00	32.00 37.00 46.00 53.00 66.00 82.00 82.00 82.00 83.00 84.00 86.00 87.00 91.00 93.00 97.00 99.00 100.00 100.00 100.00 100.00 100.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.03 0.1 0.39 16 77 179.99 32 50 11 7		

Total Particles = 100.

			Strear	n Ass	essm	ent Fo	orm (F	orm 1)			
				Unified S	tream Method	lology for use	in Virginia		,			
				For use in wade	able channels cla	ssified as interm	ittent or perennia	al	Impact	Impact		
Project #	-	ct Name (App	•	Locality	Class.	HUC	Date	SAR#	Impact Length	Impact Factor		
22865.06		Mountain Valley Pipeline (Mountain Franklin Valley Pipeline, LLC) County		R3 or R4	03010101	8/31/21	S-C19	75	1			
Nam	e(s) of Evalua	tor(s)	Stream Name and Information						SAR Length			
	RC, RH		Maggodee C	reek					75			
1. Channel Condition: Assess the cross-section of the stream and prevailing condition (erosion, aggradation)												
	Ont	imal	Subo	ptimal	Conditional Catego	ory ginal	D _i	oor	Sev	rere		
Channel	100% stable banks.	r active erosion; 80- Vegetative surface al rock, prominent	erosion or unproted	ew areas of active cted banks. Majority table (60-80%).	Poor. Banks more s	less than Severe or table than Severe or ank slopes. Erosion	laterally unstabl	cised. Vertically / e. Likely to widen both banks are near		(or excavated), stability. Severe led within the banks.		
Condition	(80-100%). AND/OI bankfull benches a to their original f developed wide bar channel bars and to Transient sedimen	R Stable point bars / ure present. Access floodplain or fully nkfull benches. Mid- ransverse bars few.	Vegetative protect prominent (60 Depositional feal stability. The ban channels are well d has access to bankl developed flo portions of the 1 sediment covers 1	tion or natural rock -80%) AND/OR -80%) AND/OR tures contribute to htfull and low flow efined. Stream likely ull benches, or newly odplains along reach. Transient 0-40% of the stream tom.	may be preser both banks. Vege 40-60% of banks. S vertical or und 40-60% Sediment transient, cont Deposition that co may be forming/p shaped channels protection on > 40 depositional feature	nt on 40-60% of tative protection on treambanks may be errout. AND/OR may be temporary / intuition to stability, ntribute to stability, ntribute to stability, resent. AND/OR V-5 have vegetative % of the banks and s which contribute to stility.	vertical. Erosion pi banks. Vegetative 20-40% of banks, prevent erosion. It the stream is cov Sediment is temp nature, and contri AND/OR V-shal vegetative protect 40% of the banks.	resent on 60-80% of protection present or and is insufficient to AND/OR 60-80% of ered by sediment. porary / transient in buting to instability, bed channels have tion is present on > and stable sediment in is absent.	Streambed below av majority of banks Vegetative protect than 20% of banks erosion. Obviou present. Erosion/rav AND/OR Aggradin	erage rooting depth, vertical/undercut. ion present on less s, is not preventing s bank sloughing v banks on 80-100%. g channel. Greater n bed is covered by uting to instability. channels and/or	CI	
Scores	;	3	2	.4	:	2	1	.6	1	l	2.40	
2. RIPARIAN	N BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) Conditional Category Optimal Suboptimal Marginal Poor					io limited to						
Riparian Buffers		e canopy cover. within the riparian	High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory.	Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation).	High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover.	Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory.	High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition.	Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions.	Assessment areas within t RO	he temporary		
			High	Low	High	Low	High	Low	1			
Scores	1	.5	1.2	1.1	0.85	0.75	0.6	0.5]			
. Determine squ	rian areas along e uare footage for ea iparian Area and S % Riparian Area> Score >	ach by measuring	or estimating lengt	th and width. Cald			of % I	the sums Riparian equal 100 100%				
	55516 -	0.0	0.10	0.00					CI= (Sum % RA * Sc	ores*0.01)/2		
Left Bank	% Riparian Area>	10%	15%	75%				100%	Rt Bank CI >	0.81	CI	
Leit Dalik	Score >	0.5	0.75	0.85					Lt Bank CI >	0.80	0.80	
3. INSTREAM		ried substrate size	es, water velocity a	nd depths; woody	and leafy debris;	stable substrate; lo	ow embededness;	shade; undercut l	oanks; root mats; S	AV; riffle/pool		
					al Category				NOTES>>			
Instream Habitat/ Available Cover	Habitat elements an	imal e typically present in % of the reach.	Stable habitat ele present in 30-50% adequate for r	ptimal ments are typically of the reach and are maintenance of ations.	Stable habitat ele present in 10-30% adequate for r	rginal Poor lements are typically 6 of the reach and are lacking or are unstable. Habitat elements listed above are lacking or are unstable. Habitat elements are typically present in less than 10% of the reach.			CI			
	I			populations.		populations.		1070 01 410 104011.		Stream Gradient		

Scores

1.5

1.2

0.9

0.5

High / Low

0.90

Stream Impact Assessment Form Page 2									
Project #	Project Name (App	Locality	Cowardin Class.	HUC	Date	SAR#	Impact Length	Impact Factor	
22865.06	Mountain Valley Pipeline (Mountain Valley Pipeline, LLC)		Franklin County	R3 or R4	03010101	8/31/21	S-C19	75	1
4. CHANNEL ALTERATION: Stream crossings, riprap, concrete, gabions, or concrete blocks, straightening of channel, channelization, embankments, spoil piles, constrictions, livestock									
	Conditional Category					NOTES>>			
	Negligible Mi		nor Moderate		Severe				
Channel Alteration	Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized.	Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	20-40% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	40 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	80 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	by any of the chanr in the parameter g	of reach is disrupted nel alterations listed uidelines AND/OR ored with gabion, cement.		
Scores	1.5	1.3	1.1	0.9	0.7	0	.5		
PEACH CONDITION INDEX and STDEAM CONDITION LIMITS FOR THIS DEACH									

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

THE REACH CONDITION INDEX (RCI) >> 1.04

RCI= (Sum of all CI's)/5, except if stream is ephemeral RCI = (Riparian CI/2)

COMPENSATION REQUIREMENT (CR) >> 78

CR = RCI X L_i X IF

INSERT PHOTOS:


(WSSI Photo Location L:\22000s\22800\22865.06\Admin\05-ENVR\Field Data\)




CAPTION. Assessment is limited to areas within the temporary $\ensuremath{\mathsf{ROW}}$.


ח	FSCI	RIRE	PRO	POSE	IMP/	CT.

PROVIDED UNDER SEPARATE COVER






CL ST	CL STAKEOUT POINTS: S-C19 CROSS SECTION B (PIPE CL)							
	PRE-CROSSING							
PT. LOC.	NORTHING	EASTING	ELEV	VERT.	HORZ.			
P1. LOC.	NOKIHING	EASTING	ELEV	DIFF.	DIFF.			
TS-L	13456754.97	1981648.05	869.62					
BS-L	13456757.42	1981658.67	855.55					
THW	13456760.79	1981673.27	855.18					
BS-R	13456769.23	1981707.65	856.04					
TS-R	13456774.87	1981731.37	866.44					




LEGEND STUDY AREA (EASEMENT) EXISTING SURVEY-LOCATED THALWEG EXISTING SURVEY-LOCATED EDGE OF WATER (AS NECESSARY) EXISTING CONTOUR LINE (MAJOR) EXISTING CONTOUR LINE (MINOR) EXISTING SURVEYED GROUND SHOT ELEVATION BENCHMARK POINT (WSSI)

SURVEY NOTES:

- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on September 4, 2018.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).
- 4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.
- 5. All section views shown are left to right facing downstream.





CROSS SECTION LEGEND EXISTING GRADE

NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.

CROSS SECTION H: 1"=10' V: 1"=5'



PHOTO TAKEN LOOKING DOWNSTREAM ALONG LEFT BANK ON 03/26/2021

Wetland

19



PHOTO TAKEN LOOKING UPSTREAM FROM ALONG LEFT BANK ON 03/26/2018






PHOTO TAKEN LOOKING DOWNSTREAM



PHOTO TAKEN LOOKING UPSTREAM ALONG RIGHT BANK ON 03/26/2018





Vertical Datum: NAVD 88 Boundary and Topo Source:

WSSI 2' C.I. Topo							
Design	Draft	Approved					
PFS	NAS/SIH	NAS					
Sheet #							
1 of 1							

Computer File Name: :\Survey\22000s\22800\22865.03\Spread I Work Dwgs 2865_03 S-I MP 268-278 Sheets.dwg