Baseline Assessment – Stream Attributes

Reach S-F10 (Timber Mat Crossing) Ephemeral Spread I Franklin County, Virginia

| Data | Included |
|---|---------------|
| Photos | ✓ |
| SWVM Form | ✓ |
| FCI Calculator and HGM Form | ✓ |
| RBP Physical Characteristics Form | ✓ |
| Water Quality Data | N/A – No flow |
| RBP Habitat Form | ✓ |
| RBP Benthic Form | ✓ |
| Benthic Identification Sheet | N/A – No flow |
| Wolman Pebble Count | ✓ |
| RiverMorph Data Sheet | ✓ |
| USM Form (Virginia Only) | ✓ |
| Longitudinal Profile and Cross Sections | ✓ |

Spread I Stream S-F10 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking NE upstream, CL



Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking SW downstream, CL

Spread I Stream S-F10 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking N at right streambank, CL



Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking SE at left streambank, CL

Spread I Stream S-F10 (Timber Mat Crossing) Franklin County



Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking SW downstream, CL

| (v2.1, Sept 2015) | | Woul | паш уапеу гірешіе | | cimal Degrees) | Lat. | 37.046037 | Lon. | -73.013334 | | WEATHER. | | Sullily | | DATE. | August 2 | 6, 2021 |
|--|---------------------|-----------------------|---|--------------------|-------------------------------|------|--|--------------------|---------------|-----|--|--------------------|----------------|--|---|----------------------|-------------|
| IMPACT STREAM/SITE ID (watershed size (acreage), | | | S-F10 | / 2.3 ac | | | MITIGATION STREAM CLASS. (watershed size {acreage | | | N: | | | | | comments: | | |
| STREAM IMPACT LENGTH: | 20 | FORM OF MITIGATION | : RESTORATION (Levels I-III) | | OORDINATES: cimal Degrees) | Lat. | | Lon. | | | PRECIPITATION PAST 48 HRS: | | | Mitig | ation Length: | | |
| Column No. 1- Impact Existing | g Condition (De | bit) | Column No. 2- Mitigation Existing C | ondition - Base | eline (Credit) | ľ | Column No. 3- Mitigation Pr Post Completion | | Years | | Column No. 4- Mitigation Proje Post Completion (6 | | ears | Columi | No. 5- Mitigation Projecte | ed at Maturity (Cr | redit) |
| Stream Classification: | Ephe | meral | Stream Classification: | | | | Stream Classification: | | 0 | Str | tream Classification: | | 0 | Stream Classifie | ation: | 0 | |
| Percent Stream Channel SI | | 13.15 | Percent Stream Channel SI | | | | Percent Stream Channel S | | 0 | | Percent Stream Channel Sle | | 0 | Per | cent Stream Channel SI | | 0 |
| HGM Score (attach d | ata forms): | | HGM Score (attach | data forms): | | | HGM Score (attach | data forms): | | | HGM Score (attach da | ata forms): | | | HGM Score (attach da | ata forms): | |
| | | Average | | | Average | | | | Average | | | | Average | | | | Average |
| Hydrology Biogeochemical Cycling Habitat | 0.3 0.32 0.09 | 0.23666667 | Hydrology Biogeochemical Cycling Hahitat | | 0 | | Hydrology Biogeochemical Cycling | | 0 | Bio | ydrology iogeochemical Cycling | | 0 | Hydrology Biogeochemica Habitat | Cycling | | 0 |
| PART I - Physical, Chemical and | | cators | PART I - Physical, Chemical ar | d Biological In | dicators | | PART I - Physical, Chemical at | nd Biological In | dicators | па | PART I - Physical, Chemical and | Biological Indi | icators | | - Physical, Chemical and | Biological Indica | ators |
| | Points Scale Range | Site Score | | Points Scale Range | Site Score | | | Points Scale Range | Site Score | | | Points Scale Range | Site Score | | | Points Scale Range | Site Score |
| PHYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | | PHYSICAL INDICATOR (Applies to all streams | s classifications) | | РН | HYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDI | CATOR (Applies to all streams | classifications) | |
| USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Low Gradient Data Sheet) | | | | USEPA RBP (High Gradient Data Sheet) | | | us | SEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Hi | gh Gradient Data Sheet) | | |
| | 0-20 | 0 | Epifaunal Substrate/Available Cover | 0-20 | | | Epifaunal Substrate/Available Cover | 0-20 | | | Epifaunal Substrate/Available Cover | 0-20 | | | trate/Available Cover | 0-20 | |
| Embeddedness Velocity/ Depth Regime | 0-20 | 8 | Pool Substrate Characterization Pool Variability | 0-20 | | | Embeddedness Velocity/ Depth Regime | 0-20 | | | Embeddedness Velocity/ Depth Regime | 0-20 | | Embeddednes Velocity/ Depti | | 0-20 | |
| Sediment Deposition | 0-20 | 16 | Sediment Deposition | 0-20 | | | 4. Sediment Deposition | 0-20 | | 4. | Sediment Deposition | 0-20 | | 4. Sediment Dep | osition | 0-20 | |
| Channel Flow Status | 0-20 | 0 | 5. Channel Flow Status | 0-20 | | | 5. Channel Flow Status | 0-20 | | | Channel Flow Status | 0-20 | | Channel Flow | Status | 0-20 | |
| 6. Channel Alteration | 0-20 | 14 | Channel Alteration | 0-20 | | | 6. Channel Alteration | 0-20 | | | Channel Alteration | 0-20 | | Channel Altera | | 0-20 | |
| 7. Frequency of Riffles (or bends) | 0-20 | 0 | 7. Channel Sinuosity | 0-20 | | | 7. Frequency of Riffles (or bends) | 0-20 | | | Frequency of Riffles (or bends) | 0-20 | | | Riffles (or bends) | 0-20 | |
| 8. Bank Stability (LB & RB) | 0-20 | 16 | 8. Bank Stability (LB & RB) | 0-20 | | | 8. Bank Stability (LB & RB) | 0-20 | | 8. | Bank Stability (LB & RB) | 0-20 | | Bank Stability | LB & RB) | 0-20 | |
| Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | 0-20 | 16 | Vegetative Protection (LB & RB) Regetative Zone Width (LB & RB) | 0-20 | | | Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB) | 0-20 | | | Vegetative Protection (LB & RB) D. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | | tection (LB & RB) ative Zone Width (LB & RB) | 0-20 | |
| Total RBP Score | Suboptimal | 88 | Total RBP Score | Poor | 0 | | Total RBP Score | Poor | 0 | To | otal RBP Score | Poor | 0 | Total RBP Score | ative Zone Width (LB & RB) | Poor | 0 |
| Sub-Total | Odboptima | 0.73333333 | Sub-Total | 1 001 | 0 | | Sub-Total | 1 001 | Ö | | ub-Total | 1 001 | ő | Sub-Total | | 1 001 | 0 |
| CHEMICAL INDICATOR (Applies to Intermittee | nt and Perennial S | • | CHEMICAL INDICATOR (Applies to Intermitter | t and Perennial S | treams) | | CHEMICAL INDICATOR (Applies to Intermitte | nt and Perennial S | treams) | СН | HEMICAL INDICATOR (Applies to Intermitter | nt and Perennial S | Streams) | CHEMICAL INDI | CATOR (Applies to Intermitter | t and Perennial Stre | eams) |
| WVDEP Water Quality Indicators (General | 1) | | WVDEP Water Quality Indicators (General | | | | WVDEP Water Quality Indicators (General | 1) | | w | VDEP Water Quality Indicators (General |) | | WVDEP Water C | uality Indicators (General |) | / |
| Specific Conductivity | | | Specific Conductivity | | 0 | | Specific Conductivity | | | Sp | pecific Conductivity | | | Specific Condu | tivity | | |
| 100-199 - 85 points | 0-90 | | | 0-90 | | | | 0-90 | | | | 0-90 | | | | 0-90 | |
| рп | 0-80 | | рп | 5-90 0-1 | | | рп | 5-90 0-1 | | pn | 1 | 5-90 0-1 | | pn | | 5-90 0-1 | |
| 5.6-5.9 = 45 points | 0-80 | | | 5-90 | | | | 5-90 | | _ | | 5-90 | | | | 5-90 | |
| DO | _ | | DO | _ | | | DO | _ | | DC | 0 | | | DO | | _ | |
| | 10-30 | | | 10-30 | | | | 10-30 | | | | 10-30 | | | | 10-30 | |
| Sub-Total | • | | Sub-Total | • | 0 | | Sub-Total | | 0 | Su | ub-Total | | 0 | Sub-Total | | • | 0 |
| BIOLOGICAL INDICATOR (Applies to Intermit | tent and Perennial | Streams) | BIOLOGICAL INDICATOR (Applies to Intermit | ent and Perennial | Streams) | | BIOLOGICAL INDICATOR(Applies to Interm | ittent and Perenr | nial Streams) | ви | IOLOGICAL INDICATOR (Applies to Interm | ittent and Peren | inial Streams) | BIOLOGICAL IN | DICATOR (Applies to Interm | ittent and Perennia | al Streams) |
| WV Stream Condition Index (WVSCI) | | | WV Stream Condition Index (WVSCI) | | | | WV Stream Condition Index (WVSCI) | | | w | /V Stream Condition Index (WVSCI) | | | WV Stream Con | dition Index (WVSCI) | | |
| | 0-100 0-1 | | | 0-100 0-1 | | | | 0-100 0-1 | | | | 0-100 0-1 | | | | 0-100 0-1 | |
| Sub-Total | | 0 | Sub-Total | - | 0 | | Sub-Total | | 0 | Su | ub-Total | | 0 | Sub-Total | | | 0 |
| | | | | | | | | | | | | | | | | | |
| PART II - Index and U | Init Score | | PART II - Index and | Unit Score | | | PART II - Index and | I Unit Score | | | PART II - Index and U | nit Score | | | PART II - Index and U | nit Score | |
| Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score | | Index | Linear Feet | Unit Score | | Index | Linear Feet | Unit Score | | Index | Linear Feet | Unit Score |
| 0.502 | 20 | 10.0333333 | 0 | 0 | 0 | | 0 | 0 | 0 | | 0 | 0 | 0 | | 0 | 0 | 0 |

| | | | High-G | | | | ms in Ap | - | a | | |
|--------|---|---------------|----------------|---------------|---------------|---------------|---------------------------|-------------------------|-----------------|--------------|---------|
| | Team: | RH, CL | | rieia L | Jala Sile | et and C | | | M Northing: | 37 0/18037 | |
| Dro | oject Name: | | /allev Pinelir | | | | | | M Easting: | | |
| 110 | | Franklin Co | | 10 | | | | - | pling Date: | | |
| 0.4 | | | | 1 (1 (6) | 0.0 | 01 T | | | | 00/20/21 | |
| SA | AR Number: | S-F10 | Reacn | Length (ft): | 68 | Stream Ty | /pe: Ephe | meral Stream | | | • |
| | Top Strata: | Shi | rub/Herb Str | ata | (determined | d from perce | ent calculate | d in V _{CCANO} | _{PY}) | | |
| Site | and Timing: | Project Site | | | | • | Before Projec | :t | | | |
| Sample | Variables | | | | | | | | | | |
| 1 | Average percent cover over channel by tree and sapling canopy. Measure at no fewer than 10 roughly equidistant points along the stream. Measure only if tree/sapling cover is at least 20%. (If less than 20%, enter at least one value between 0 and 19 to trigger Top Strata choice.) List the percent cover measurements at each point below: | | | | | | | | | | |
| | | | | | | | | | | | - |
| | 20 | 20 | 15 | 10 | 5 | 5 | 0 | 5 | 0 | 0 | |
| | | | | | | | | | | | |
| 2 | V_{EMBED} | | | | | | at no fewer | | | | 1.8 |
| | along the stream. Select a particle from the bed. Before moving it, determine the percentage of the surface and area surrounding the particle that is covered by fine sediment, and enter the rating | | | | | | | | | | |
| | according to the following table. If the bed is an artificial surface, or composed of fine sediments, use a | | | | | | | | | | |
| | | | | | | | a rating sco | | inio ocanin | orito, add a | |
| | | | | | • | | cles (rescale | | tts Menaha | n and | 1 |
| | | Minshall 19 | • | ioi giavoi, o | | ouldor parti | 0.00 (1.0004). | Ju 11011111110 | ito, mogana | ii, aiia | |
| | | Rating | Rating Des | crintion | | | | | | | |
| | | 5 | | | overed sur | rounded or | buried by fi | ne sedimen | t (or bedroc | :k) | |
| | | 4 | | | | | d, or buried | | | | |
| | | 3 | | | | | ed, or buried | | | | 1 |
| | | 2 | | | | | ed, or buried | | | | |
| | | 1 | | | covered, su | ırrounded, o | or buried by | fine sedime | nt (or artific | ial surface) | |
| | List the rati | ngs at each | point below | <i>l</i> : | | | | | | | |
| | 1 | 1 | 1 | 1 | 1 | 4 | 1 | 4 | 3 | 2 | |
| | 1 | 1 | 4 | 1 | 4 | 1 | 1 | 1 | 1 | 4 | |
| | 1 | 3 | 1 | 4 | 1 | 1 | 1 | 1 | 1 | 1 | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| 3 | $V_{SUBSTRATE}$ | Median stre | eam channe | l substrate | oarticle size | . Measure | at no fewer t | han 30 rou | ghly equidis | tant points | 0.00: |
| | | along the s | tream; use t | he same po | ints and pa | rticles as us | sed in V _{EMBEI} |). | | | 0.08 in |
| | Enter partic | le size in in | ches to the | nearest 0.1 | inch at eacl | h point belo | w (bedrock : | should be c | ounted as 9 | 9 in, | |
| | | | 0.0 in, sand | | | | ` | | | , | |
| | 0.08 | 0.08 | 0.08 | 0.08 | 0.08 | 5.90 | 6.60 | 5.10 | 0.60 | 5.90 | |
| | 0.08 | 0.08 | 10.60 | 0.08 | 1.50 | 0.08 | 0.08 | 5.00 | 0.08 | 0.60 | |
| | 0.08 | 6.90 | 0.08 | 4.50 | 0.08 | 0.08 | 0.08 | 0.08 | 0.08 | 0.08 | |
| | 3.00 | 3.00 | 3.00 | 1.00 | 3.00 | 3.00 | 3.00 | 3.00 | 3.00 | 3.00 | |
| | | | | | | | | | | | |
| 4 | V_{BERO} | Total perce | nt of eroded | stream ch | annel bank | Enter the t | otal number | of feet of e | roded bank | on each | |
| • | · DEKU | | | | | | nks are ero | | | | 51 % |
| | | may be up | | 5 | | | | , - | | | J . /v |
| | | | Left Bank: | 15 | i ft | 1 | Right Bank: | 20 |) ft | | |

| Sampl | e Variables | 5-9 within t | the entire ı | riparian/buf | fer zone ad | jacent to th | ne stream c | hannel (25 | feet from e | ach bank). | |
|--------------|-------------|--------------|--------------|-----------------|---|---------------|-------------------------------|----------------|----------------|--------------|-------------|
| 5 | V_{LWD} | stream rea | ch. Enter t | | rom the enti ulated. | re 50'-wide | ter and 36 ir buffer and w | | | | 1.5 |
| | | A !! | 1 61 | , | | | oody stems: | 11 1000 | 1 | | |
| 6 | V_{TDBH} | | | | ily if V _{CCANOF} tree DBHs | | ng cover is a | at least 20% |). Trees ar | e at least 4 | Not Used |
| | | • | , | | | | n) within the | huffer on e | ach side of | | |
| | | the stream | | nents of indi | viduai ii ees | (at least 4 i | ii) wiliiii liie | : builei oii e | acii side di | | |
| | | | Left Side | | | | | Right Side | | | |
| | 0 | | | | | 0 | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| | | | | | | | | | | | |
| 7 | V_{SNAG} | | | | | | et of stream. | Enter num | ber of snag | s on each | 0.0 |
| | | side of the | siream, an | u trie amour | nt per 100 fe | et will be ca | liculated. | | | | 0.0 |
| | | | Left Side: | | 0 | | Right Side: | | 0 | | |
| 8 | V_{SSD} | | | | | | hes dbh) pei ibs on each | | | | 110.3 |
| | | | | | e calculated | | ibs on each | side of the | Sirearri, arro | ıııe | 110.5 |
| | | · | Left Side: | : 3 | 35 | | Right Side: | | 10 | | |
| 9 | V_{SRICH} | | | | | | am reach. C ive species ¡ | | | | 0.00 |
| | | | | | | | from these of | | ııı sırata. Sp | ecies | 0.00 |
| | | | p 1 = 1.0 | | | | | | | | |
| | Acer rubru | ım | | Magnolia t | ripetala | | Ailanthus a | ltissima | | Lonicera ja | ponica |
| | Acer sacci | harum | | Nyssa sylv | atica . | | Albizia julib | rissin | | Lonicera ta | tarica |
| | Aesculus | flava | | Oxydendrur | n arboreum | | Alliaria peti | olata | | Lotus corni | culatus |
| | Asimina tr | iloba | | Prunus sei | rotina | | Alternanthe | | | Lythrum sa | licaria |
| | Betula alle | ghaniensis | | Quercus a | lba | | philoxeroid | | <u> </u> | Microstegiun | |
| | Betula len | - | | Quercus c | | | Aster tatari | cus | | Paulownia | |
| | Carya alba | | | Quercus in | | | Cerastium | | | Polygonum o | |
| | Carya glal | | | Quercus p | | | Coronilla va | | | Pueraria m | |
| | Carya ova | | | Quercus ru | | | Elaeagnus u | | | Rosa multif | |
| | • | | | Quercus v | | | | | _ | Sorghum h | |
| | Carya ova | | | Sassafras | | | Lespedeza | | | • | • |
| | Cornus flo | | | | | | Lespedeza | | Ш | Verbena br | asilierisis |
| | Fagus gra | | | Tilia ameri | | | Ligustrum ob | | | | |
| | Fraxinus a | | | Tsuga can | | | Ligustrum s | sinense | | | |
| \checkmark | Liriodendro | | | Ulmus ame | ericana | | | | | | |
| | Magnolia a | acuminata | | | | | | | | | |
| | | 1 | Species in | Group 1 | | | | 1 | Species in | Group 2 | |

| - | ample Variables 10-11 within at least 8 subplots (40" x 40", or 1m x 1m) in the riparian/buffer zone within 25 feet from each ank. The four subplots should be placed roughly equidistantly along each side of the stream. 10 V _{DETRITIES} Average percent cover of leaves, sticks, or other organic material. Woody debris <4" diameter and | | | | | | | | | | |
|----------------|---|--|----------------------------|---------------|-------------------------------|--------------|---------------|-------------|------------------------------|------------------------|----------------------------------|
| 10 | V _{DETRITUS} | | | | sticks, or oth ercent cove | | | | | er and | 19.38 % |
| | | | Left | Side | | | Righ | t Side | | | |
| | | 20 | 15 | 10 | 20 | 35 | 20 | 5 | 30 | | |
| 11 | \/ | Average pe | reentage co | over of borb | account you | otation (may | acura anly if | troo cover | is <20%). D |)o not | |
| | V_{HERB} | include woo | ody stems a percentage: | t least 4" db | oh and 36" t | all. Because | e there may | be several | layers of gro of ground v | und cover | 81 % |
| | | | Left | Side | | | Righ | t Side | | | |
| | | 80 | 85 | 90 | 80 | 65 | 80 | 95 | 70 | | |
| | | | | | | | | | | | |
| Sample | e Variable 1 | 2 within the | e entire cat | chment of | the stream. | | | | | | |
| 12 | V_{WLUSE} | Weighted A | Average of F | Runoff Scor | e for waters | hed: | | | | | 0.47 |
| | | | Land | Use (Choos | e From Dro | p List) | | | Runoff Score | % in Catch- ment | Running Percent (not >100) |
| | Forest and n | ative range (< | 50% ground | cover) | | | | • | 0.5 | 36.74 | 36.74 |
| | Forest and n | ative range (> | >75% ground | cover) | | | | — | 1 | 14.25 | 50.99 |
| | Open space | Open space (pasture, lawns, parks, etc.), grass cover >75% | | | | | | | 0.3 | 49.01 | 100 |
| | | | | | | | | | | | |
| | | | | | | | | _ | | | |
| | | | | | | | | _ | | | |
| | | | | | | | | _ | | | |
| | | | | | | | | ▼ | | | |
| | S | S-F10 | | | | | No | tes: | _ | | |
| V | ariable | Value | VSI | Land Cov | er Analysis | was comp | oleted using | the 2019 | National La | and Cover | Database |
| | CANOPY | Not Used, | Not Used | (NLCD), f | rom Lands | at satellite | imagery ar | nd other su | upplementa | ry datasets | |
| | MBED | <20% 1.8 | 0.38 | | | | | | ted stream ed to the ne | • | number |
| | | 0.08 in | 0.04 | | | | | | | | |
| | UBSTRATE | 51 % | 0.80 | | | | | | | | |
| V _L | ERO | 1.5 | 0.18 | | | | | | | | |
| | | | Not Used | | | | | | | | |
| | DBH | Not Used | | | | | | | | | |
| | NAG | 0.0 | 0.10 | | | | | | | | |
| Vs | | 110.3 | 1.00 | | | | | | | | |
| | RICH | 0.00 | 0.00 | | | | | | | | |
| | ETRITUS | 19.4 % | 0.24 | | | | | | | | |
| | ERB | 81 % | 1.00 | | | | | | | | |
| V _W | LUSE | 0.47 | 0.49 | | | | | | | | |

Ver. 10-20-17

FCI Calculator for the High-Gradient Headwater Streams in Appalachia

To ensure accurate calculations, the <u>UPPERMOST STRATUM</u> of the plant community is determined based on the calculated value for V_{CCANOPY} (≥20% cover is required for tree/sapling strata). Go to the SAR Data Entry tab and enter site characteristics and data in the yellow cells. For information on determining how to split a project into SARs, see Chapter 5 of the Operational Draft Regional Guidebook for the Functional Assessment of High-Gradient Headwater Streams and Low-Gradient Perennial Streams in Appalachia (Environmental Laboratory U.S. Army Corps of Engineers 2017).

Project Name: Mountain Valley Pipeline

Location: Franklin County

Sampling Date: 08/26/21 Project Site Before Project

Subclass for this SAR:

Ephemeral Stream

Uppermost stratum present at this SAR: SAR number: S-F10

Shrub/Herb Strata

Functional Results Summary:

Enter Results in Section A of the Mitigation Sufficiency Calculator

| Function | Functional Capacity Index |
|------------------------|------------------------------|
| Hydrology | 0.30 |
| Biogeochemical Cycling | 0.32 |
| Habitat | 0.09 |

Variable Measure and Subindex Summary:

| Variable | Name | Average Measure | Subindex |
|------------------------|---|--------------------|----------|
| V _{CCANOPY} | Percent canpoy over channel. | Not Used, <20% | Not Used |
| V _{EMBED} | Average embeddedness of channel. | 1.77 | 0.38 |
| V _{SUBSTRATE} | Median stream channel substrate particle size. | 0.08 | 0.04 |
| V _{BERO} | Total percent of eroded stream channel bank. | 51.47 | 0.80 |
| V_{LWD} | Number of down woody stems per 100 feet of stream. | 1.47 | 0.18 |
| V _{TDBH} | Average dbh of trees. | Not Used | Not Used |
| V _{SNAG} | Number of snags per 100 feet of stream. | 0.00 | 0.10 |
| V _{SSD} | Number of saplings and shrubs per 100 feet of stream. | 110.29 | 1.00 |
| V _{SRICH} | Riparian vegetation species richness. | 0.00 | 0.00 |
| V _{DETRITUS} | Average percent cover of leaves, sticks, etc. | 19.38 | 0.24 |
| V _{HERB} | Average percent cover of herbaceous vegetation. | 80.63 | 1.00 |
| V _{WLUSE} | Weighted Average of Runoff Score for Catchment. | 0.47 | 0.49 |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME S-F10 | | LOCATION Franklin | |
|----------------------------|-----------------------------------|-------------------------------|---|
| STATION#R | IVERMILE 270.5 | STREAM CLASS Epheme | eral |
| LAT <u>37.048037</u> LO | ONG79.813934 | RIVER BASIN Upper Roa | anoke |
| STORET# | | AGENCY VADEQ | |
| INVESTIGATORS RH, CL | - | | |
| FORM COMPLETED BY | CL | DATE 08/26/21 TIME 1400 | REASON FOR SURVEY Baseline Assessment |
| | | | Has there been a heavy rain in the last 7 days? |
| WEATHER CONDITIONS | Now | Past 24 hours | ✓ Yes No |
| | rain (| (heavy rain) (steady rain) | Air Temperature 34.4 °C |
| | showers | s (intermittent) | Other |
| | | ear/sunny | |
| SITE LOCATION/MAP | Draw a map of the sit | te and indicate the areas sam | apled (or attach a photograph) |
| | | | Streath |
| | | | 0//- |
| | | Cho V | 1/ / 67 |
| | | V Colon | 1 1.60 |
| | 7 |) FIDEN. | Deuze |
| | , | 70)11/ | Dense. |
| | | 1 | 4/1/ |
| | | 1/4 | 1//(5 |
| | | | 7 |
| | | 11 | |
| | | | David- |
| | | | 1 LARCIN |
| | TIDP. | | STILAN |
| | 1170 | - W | <u> </u> |
| | | | |
| | | J | |
| | - | 160 | |
| | | 0 - 1 | 1001 |
| | | Bridge | e/ROW |
| | | | 1 |
| | - | | 7 |
| CTDEAM | Stroom Subt | | Stussen Time |
| STREAM CHARACTERIZATION | Stream Subsystem Perennial Inte | ermittent Tidal | Stream Type ☐Coldwater ✓ Warmwater |
| | Stream Origin | ☐Spring-fed | Catchment Area 0.01 km ² |
| | Non-glacial montane Swamp and bog | Mixture of origins Other | |
| | B wamp and oog | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSI FEATURE | | Predom Fores Field Agric Resid | Pasture Industrultural Other | ercial | Local Watershed NPS ☑ No evidence ☐ Son ☐ Obvious sources ☐ Local Watershed Erosi ☑ None ☐ Moderate | ne potential sources |
|---|--|--|---|---------------------------------------|--|--|
| RIPARIAN VEGETAT (18 meter l | ΓION | | e the dominant type and | | minant species present He | rbaceous |
| INSTREA FEATURE | | Estimat Samplin Area in Estimat | km² (m²x1000) 001 ed Stream Depth 0 Velocity 0 r | m m² km² m | _ | ly shaded Shaded state of the shaded Shaded represented by Stream Run% No No |
| LARGE W DEBRIS | VOODY | LWD Density | of LWD 1 1 | m ² /km ² (LWD/ | reach area) | |
| AQUATIC VEGETATION Indicate the dominant type and record the dominant species present Rooted emergent Attached Algae Dominant species present Portion of the reach with aquatic vegetation 40 % | | | | | Free floating | |
| WATER Q | QUALITY | Specific Dissolve pH N/A Turbidi | cature N/A 0 C Conductance N/A ed Oxygen N/A ty N/A ttrument Used N/A | - | | Chemical Other NO FLOW Globs Flecks Wred |
| SEDIMEN SUBSTRA | | Odors Norm Chem Other Oils | al Sewage ical Anaerobic The Slight Modera | Petroleum None | — Εροking at stones whic are the undersides blace | □Paper fiber □Sand Other □Sand h are not deeply embedded, k in color? |
| INO | | STRATE (| COMPONENTS 00%) | | ORGANIC SUBSTRATE C (does not necessarily add | |
| Substrate Type | Diamet | er | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area |
| Bedrock Boulder | > 256 mm (10") | | | Detritus | sticks, wood, coarse plant materials (CPOM) | 65 |
| Cobble Gravel | 64-256 mm (2.5 2-64 mm (0.1"-2 | | 5 10 | Muck-Mud | black, very fine organic (FPOM) | |
| Sand Silt Clay | 0.06-2mm (gritty) 5 0.004-0.06 mm 10 < 0.004 mm (slick) 70 | | | Marl | grey, shell fragments | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME S-F10 | LOCATION Franklin | | | | |
|---|----------------------------------|---------------------------------------|--|--|--|
| STATION # RIVERMILE 270.5 | STREAM CLASS Ephemeral | | | | |
| LAT <u>37.048037</u> LONG <u>-79.813934</u> | RIVER BASIN Upper Roanoke | | | | |
| STORET# | AGENCY VADEQ | | | | |
| INVESTIGATORS RH, CL | | | | | |
| FORM COMPLETED BY CL | DATE 08/26/21 TIME 1420 AM PM | REASON FOR SURVEY Baseline Assessment | | | |

| | Habitat | | Condition | ı Category | | |
|--|---|---|---|---|---|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | |
| | SCORE 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25-50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | |
| ted ir | SCORE 8 ▼ | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | |
| ıram | SCORE 0 ▼ | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| Paran | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | |
| | _{SCORE} 16 ▼ | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | |
| | SCORE U | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | |

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| | Habitat | | Condition | n Category | |
|--|---|--|--|--|---|
| | Parameter | Optimal | Suboptimal | Marginal | Poor |
| | 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. |
| | SCORE 14▼ | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| g reach | 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. |
| ampl | SCORE 0 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| Parameters to be evaluated broader than sampling reach | 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. |
| e ev | SCORE 8 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| to b | SCORE 8 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| Parameters to | 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. |
| | SCORE 8 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| | SCORE 8 ▼ | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| | 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. |
| | SCORE 9 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| | SCORE 9) | | | | 2 1 0 |

Total Score 88

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME S-F | -10 | | | | | | LOC | ATION | √ Fran | klir | 1 | | | | | | | | | |
|-------------------------------|----------|------|--------|-------|--------|----------------|-------------|------------------------|---------------------------|------------|--------------|------|---------------------|-------|--|--------|------|-------|-----|-----|
| STATION # RIVERMILE 270.5 | | | | | | | | STREAM CLASS Ephemeral | | | | | | | | | | | | |
| LAT 37.048037 | | | | | | | | | RIVER BASIN Upper Roanoke | | | | | | | | | | | |
| STORET# | | | | | | | AGE | NCY V | /ADEC | 2 | | | | | | | | | | |
| INVESTIGATORS R | tH, C | L | | | | • | | | | | | | I | LOT | NUMBER | | | | | |
| FORM COMPLETE | ЭBY | C | L | | | | DAT TIM | E 08/2 | | | | | I | REAS | SON FOR SURVEY Ba | aselir | ne A | .sse: | ssm | ent |
| HABITAT TYPES | | Col | oble_ | | % | tage of e | gs | | | Ve | geta | ited | Bani | ks | % | _% | | | | |
| SAMPLE COLLECTION | LLECTION | | | | | | | | | | r-net Other | | | | | | | | | |
| | Γ | ndic | ate th | ie ni | ımbe | r of jabs Sna | /kick gs | s taken | in eac | ch l Ve | nab geta | itat | type Banl | e. | _ | | | | | |
| GENERAL COMMENTS | N | o k | oen | thi | cs; | Dry s | stre | am | | | | | | | | | | | | |
| Dominant | | | | | 0 = 2 | Absent/ | Not | Obser | ved, | | | | | = C | ommon, 3= Abund | | | | | 4 |
| Periphyton | | | | | - | 1 2 | - | | | | | nes | | 4 1 | | - | 1 | | 3 | |
| Filamentous Algae Macrophytes | | | | | | 1 2 1 2 | | | | | viac Fish | | nve | rtebi | rates | - | 1 | | 3 | |
| FIELD OBSERV | | | | | 0 = | Absent | /Not | Obse | | | | | | | rganisms), 2 = Con , 4 = Dominant (>5 | | | -9 | | |
| Porifera | 0 | 1 | 2 | 3 | 4 | Aniso | _ | | 0 | | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygop | | | 0 | | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | | 3 | 4 | Hemi | • | | 0 | | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | _ | 3 | 4 | Coleo | _ | | 0 | | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | _ | 3 | 4 | Lepid | _ | ra | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 2 | 3 | 4 | Sialid | | | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda Amphipoda | 0 | 1 | 2 | 3 | 4 4 | Coryo Tipul | | ıc | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda Decapoda | 0 | 1 | 2 | 3 | 4 | Empi | | | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simul | | | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabin | | | 0 | | 1 | 2 | 3 | 4 | | | | | | |
| | | • | | ٦ | | Culci | | | 0 | | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

County: Franklin County Stream ID: S-F10

Stream Name: UNT to Blackwater River

HUC Code: 03010101 Basin: Upper Roanoke

HUC Code: 03010101
Survey Date: 8/26/2021
Surveyors: MVP Team
Type: Representative

| | | | LE COUNT | | | | |
|-------------|-------------|-------------|----------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cum |
| | Silt/Clay | < .062 | S/C | A | 62 | 62.00 | 62.00 |
| | Very Fine | .062125 | | A | 9 | 9.00 | 71.00 |
| | Fine | .12525 | 7 | A | 5 | 5.00 | 76.00 |
| | Medium | .255 | SAND | A | 7 | 7.00 | 83.00 |
| | Coarse | .50-1.0 | 7 | A | 4 | 4.00 | 87.00 |
| .0408 | Very Coarse | 1.0-2 | 7 | - | | 0.00 | 87.00 |
| .0816 | Very Fine | 2 -4 | | - | | 0.00 | 87.00 |
| .1622 | Fine | 4 -5.7 | 7 | • | | 0.00 | 87.00 |
| .2231 | Fine | 5.7 - 8 | 7 | • | | 0.00 | 87.00 |
| .3144 | Medium | 8 -11.3 | 7 | * | 1 | 1.00 | 88.00 |
| .4463 | Medium | 11.3 - 16 | GRAVEL | * | 1 | 1.00 | 89.00 |
| .6389 | Coarse | 16 -22.6 | | A | | 0.00 | 89.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | | 3 | 3.00 | 92.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | A | 1 | 1.00 | 93.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | 1 | A | 1 | 1.00 | 94.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ^ | | 0.00 | 94.00 |
| 3.5 - 5.0 | Small | 90 - 128 | 1 | ^ | 1 | 1.00 | 95.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | A | 2 | 2.00 | 97.00 |
| 7.1 - 10.1 | Large | 180 - 256 | 1 | ▲ | 2 | 2.00 | 99.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | △ | 1 | 1.00 | 100.00 |
| 14.3 - 20 | Small | 362 - 512 | 1 | A | | 0.00 | 100.00 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | A | | 0.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | 1 | A | | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 | 1 | A | | 0.00 | 100.00 |
| | Bedrock | | BDRK | A | | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |

WOLMAN PEBBLE COUNT FORM

County: Franklin County Stream ID: S-F10

Stream Name: UNT to Blackwater River

HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 8/26/2021 Surveyors: CL, RH Type: Representative

| | | | LE COUNT | | | | |
|-------------|-------------|-------------|----------|-------------------|---------|--------|-------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cun |
| | Silt/Clay | < .062 | S/C | ^ | 62 | 62.00 | 62.00 |
| | Very Fine | .062125 | | - | 9 | 9.00 | 71.00 |
| | Fine | .12525 | | + | 5 | 5.00 | 76.00 |
| | Medium | .255 | SAND | 4 | 7 | 7.00 | 83.00 |
| | Coarse | .50-1.0 | .50-1.0 | ^ | 4 | 4.00 | 87.00 |
| .0408 | Very Coarse | 1.0-2 | 1 | ^ | | 0.00 | 87.00 |
| .0816 | Very Fine | 2 -4 | | ^ | | 0.00 | 87.00 |
| .1622 | Fine | 4 -5.7 | | ^ | | 0.00 | 87.00 |
| .2231 | Fine | 5.7 - 8 | 7 | ^ | | 0.00 | 87.00 |
| .3144 | Medium | 8 -11.3 | GRAVEL | ^ | 1 | 1.00 | 88.00 |
| .4463 | Medium | 11.3 - 16 | | ^ | 1 | 1.00 | 89.00 |
| .6389 | Coarse | 16 -22.6 | | ^ | | 0.00 | 89.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | ^ | 3 | 3.00 | 92.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | ^ | 1 | 1.00 | 93.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | ^ | 1 | 1.00 | 94.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ^ | | 0.00 | 94.00 |
| 3.5 - 5.0 | Small | 90 - 128 | | ^ | 1 | 1.00 | 95.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | ^ | 2 | 2.00 | 97.00 |
| 7.1 - 10.1 | Large | 180 - 256 | 1 | ^ | 2 | 2.00 | 99.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | ^ | 1 | 1.00 | 100.0 |
| 14.3 - 20 | Small | 362 - 512 | 1 | ^ | | 0.00 | 100.0 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | ^ | | 0.00 | 100.0 |
| 40 - 80 | Large | 1024 -2048 | 7 | A | | 0.00 | 100.0 |
| 80 - 160 | Vry Large | 2048 -4096 | 7 | A | | 0.00 | 100.0 |
| | Bedrock | | BDRK | ^ | | 0.00 | 100.0 |
| | | | | Totals: | 100 | | |

Ephemeral Stream Assessment Form (Form 1a) Unified Stream Methodology for use in Virginia For use in ephemeral streams Cowardin **Impact Impact Project Name** Project # HUC SAR# Locality Date Class. Length Factor **Mountain Valley Pipeline (Mountain** Franklin S-F10 20 22865.06 R6 03010101 08/26/21 **Valley Pipeline, LLC)** County Name(s) of Evaluator(s) **Stream Name and Information** SAR Length CL, RH Spread I; UNT to Blackwater River 217.12 ft

2. RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable)

| | | Cor | nditional Cate | gory | | | | NOTES>> | |
|---------------------|--|--|--|--|--------------------------------------|--------------------|---|-------------------|---------------|
| | Optimal | Subo | ptimal | Mar | ginal | Po | oor | | |
| Riparian Buffers | Tree stratum (dbh > 3 inches) preser with > 60% tree canopy cover and a non-maintained understory. Wetlan areas. | Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with >30% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh | ponds, open water. If present, tree | nurseries; no-till | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | | |
| | | High | Low | High | Low | High | Low | | |
| Condition Scores | 1.5 | 1.2 | 1.1 | 0.85 | 0.75 | 0.6 | 0.5 | | |
| Determine squ | urian areas along each stream bank | or estimating length | and width. Calcul | | · | of % F | the sums Riparian | | |
| Enter the % R | Riparian Area and Score for each rip | | blocks below. | | | BIOCKS | equal 100 | | |
| Right Bank | % Riparian Area> 10% Score > 0.6 | 90% 0.85 | | | | | 100% | ! | |
| | | | | | | | | CI= (Sum % RA * S | cores*0.01)/2 |
| Left Bank | % Riparian Area> 10% | 90% | | | | | 100% | Rt Bank CI > | 0.83 |
| Leit Dalik | Score > 0.6 | 0.85 | | | | | | Lt Bank CI > | 0.83 |

REACH CONDITION INDEX and STREAM CONDITION UNITS FOR THIS REACH

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.

THE REACH CONDITION INDEX (RCI) >>

RCI= (Riparian CI)/2

COMPENSATION REQUIREMENT (CR) >>

0.42

8

CR = RCI X LF X IF


INSERT PHOTOS:




CAPTION. Assessment is limited to areas within the temporary ROW.


DESCRIBE PROPOSED IMPACT:

PROVIDED UNDER SEPARATE COVER






| CL STAKEOUT POINTS: S-F10 CROSS SECTION B (PIPE CL) | | | | | | | | | |
|---|---------------------------|---------------|---------|-------|-------|--|--|--|--|
| | PR | POST-CROSSING | | | | | | | |
| DT LOC | OC. NORTHING EASTING ELEV | | VERT. | HORZ. | | | | | |
| PT. LOC. | NORTHING | EASTING | ELEV | DIFF. | DIFF. | | | | |
| TS-L | 13454234.63 | 1986431.73 | 1000.84 | | | | | | |
| BS-L | 13454236.24 | 1986429.98 | 999.36 | | | | | | |
| THW | 13454237.48 | 1986428.62 | 999.11 | | | | | | |
| BS-R | 13454237.87 | 1986428.12 | 999.19 | | | | | | |
| TS-R | 13454241.16 | 1986424.82 | 1002.77 | | | | | | |





LEGEND STUDY AREA (EASEMENT) EXISTING SURVEY-LOCATED THALWEG EXISTING SURVEY-LOCATED EDGE OF WATER (AS NECESSARY) EXISTING CONTOUR LINE (MAJOR) EXISTING CONTOUR LINE (MINOR) EXISTING SURVEYED GROUND SHOT ELEVATION 1000.3 +BENCHMARK POINT (WSSI)

SURVEY NOTES:

- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on November 26, 2018.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline
- 4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.
- 5. All section views shown are left to right facing downstream.
- 6. Cross-section B shot at location of pipe centerline (based on best professional judgement).







CROSS SECTION H: 1"=10' V: 1"=5'

CROSS SECTION LEGEND EXISTING GRADE

NOTE: ALL SECTION VIEWS SHOWN LEFT TO RIGHT FACING DOWNSTREAM.



03/28/2018




PHOTO TAKEN LOOKING DOWNSTREAM ON 03/28/2018



FROM PIPE CENTERLINE ON 03/28/2018



FROM PIPE CENTERLINE ON 03/28/2018

POST-CROSSING PHOTOS PENDING CROSSING

PHOTO TAKEN LOOKING

PENDING CROSSING

PHOTO TAKEN LOOKING

Horizontal Datum: NAD 1983 UTM ZONE 1 Vertical Datum: NAVD 88

Boundary and Topo Source: WSSI 2' C.I. Topo

Approved EJC MGE NAS Sheet # 1 of 1

Computer File Name: Survey\22000s\22800\22865.03\Spread I Work Dwgs 865_03 S-I MP 268-278 Sheets.dwg