Baseline Assessment – Stream Attributes

Reach S-G6 (Pipeline ROW) Intermittent Spread I Pittsylvania County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	N/A – slope less than 4%
RBP Physical Characteristics Form	✓
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A – Low flow
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	✓
Longitudinal Profile and Cross Sections	✓

Spread I Stream S-G6 (Pipeline ROW) Pittsylvania County



Photo Type: US VIEW Location, Orientation, Photographer Initials: Downstream at LOC looking SE upstream, CB



Location, Orientation, Photographer Initials: Downstream at LOC looking NE downstream, CB

Spread I Stream S-G6 (Pipeline ROW) Pittsylvania County



Location, Orientation, Photographer Initials: On left bank at pipe centerline looking E at right streambank, CB



Location, Orientation, Photographer Initials: On right bank at pipe centerline looking SE at left streambank, CB


Spread I Stream S-G6 (Pipeline ROW) Pittsylvania County



Photo Type: US COND Location, Orientation, Photographer Initials: Upstream at LOC looking SE upstream, CB



Location, Orientation, Photographer Initials: Upstream at LOC looking NE downstream, CB



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME		LOCATION					
STATION# RIV	VERMILE	STREAM CLASS					
LATLONG		RIVER BASIN					
STORET#		AGENCY					
INVESTIGATORS							
FORM COMPLETED BY		DATE	REASON FOR SURVEY				

WEATHER CONDITIONS	rain (si showers (% %clo	Past 24 hours heavy rain) teady rain) (intermittent) oud cover ur/sunny	Has there been a heavy rain in the last 7 days? Yes No Air Temperature0 C Other
SITE LOCATION/MAP	Draw a map of the site	and indicate the areas sam	pled (or attach a photograph)
	Dense veg	Bridge	Stream LOD
STREAM	Stream Subsystem Perennial Inter		Stream Type Coldwater Warmwater
CHARACTERIZATION	Perennial Inter Stream Origin Glacial Non-glacial montane Swamp and bog	mittent Tidal Spring-fed Mixture of origins Other	Coldwater Warmwater Catchment Areakm ²

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Fores Field Agric	Pasture Industri	ercial	No evidence Son Obvious sources Local Watershed Erosi None Moderate	ne potential sources				
RIPARIA VEGETA (18 meter	ΓΙΟΝ	Trees	e the dominant type and S ant species present	hrubs		rbaceous				
INSTREA FEATURI			ted Reach Length		Canopy Cover Partly open Part	ly shaded Shaded				
				m	High Water Mark	m				
					Proportion of Reach Re	epresented by Stream				
			km² (m²x1000) ted Stream Depth	km²	Morphology Types Riffle Pool %	Run%				
			Velocity		Channelized Yes	No				
		(111 11111			Dam Present Yes	No				
LARGE V DEBRIS	VOODY		m² of LWDn	n ² /km ² (LWD /	reach area)					
AQUATIO VEGETA		Indicate Roote Floati Domina	e the dominant type and demergent R ng Algae A	l record the do ooted submerge ttached Algae	minant species present nt Rooted floating	C				
		Portion	of the reach with aqua	tic vegetation _	%					
WATER (QUALITY	Specific	rature0 C Conductance	-	Water Odors Normal/None Sewage Petroleum Fishy	Chemical Other				
		рН	ed Oxygen		Water Surface Oils Slick Sheen None Other	Globs Flecks				
			strument Used		Turbidity (if not measu Clear ☐ Slightly tur Opaque Stained	r ed) rbid Turbid Other				
SEDIMEN SUBSTRA		Odors Norm Chem		Petroleum None	Deposits Sludge Sawdust Relict shells	Sludge Sawdust Paper fiber Sand				
		Oils Abser		te Profu	are the undersides blac	h are not deeply embedded, k in color?				
INC	ORGANIC SUBS		COMPONENTS (00%)		ORGANIC SUBSTRATE C (does not necessarily add					
Substrate Type	Diamete	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area				
Bedrock				Detritus	sticks, wood, coarse plant					
Boulder	> 256 mm (10")				materials (CPOM)					
Cobble	64-256 mm (2.5	"-10")		Muck-Mud	black, very fine organic					

Gravel

Sand

Silt

Clay

2-64 mm (0.1"-2.5")

0.06-2mm (gritty)

< 0.004 mm (slick)

0.004-0.06 mm

grey, shell fragments

Marl

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION	
STATION # RIVERMILE	STREAM CLASS	
LAT LONG	RIVER BASIN	
STORET#	AGENCY	
INVESTIGATORS		
FORM COMPLETED BY	DATE AM PM	REASON FOR SURVEY

	Habitat		Condition	ı Category						
	Parameter	Optimal	Suboptimal	Marginal	Poor					
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.					
ted in	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/depth regime (usually slow-deep).					
ıram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
Ps	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.					
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0					

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	n Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor			
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.			
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
oling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.			
samp	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing downstream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.			
e eva	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
to be	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.			
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.			
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
ĺ	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			

Total	Caama	
i otai	Score	

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME		LOCATION					
STATION #	_ RIVERMILE	STREAM CLASS					
LAT	LONG	RIVER BASIN					
STORET#		AGENCY					
INVESTIGATORS			LOT NUMBER				
FORM COMPLETED	ВҮ	DATE REASON FOR SURVEY TIME					
	Indicate the percentage of						
HABITAT TYPES							

HABITAT TYPES	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%
SAMPLE COLLECTION	Gear used D-frame kick-net Other
	How were the samples collected? wading from bank from boat
	Indicate the number of jabs/kicks taken in each habitat type. Cobble Snags Vegetated Banks Sand
	Cobble Snags Vegetated Banks Sand Submerged Macrophytes Other ()
GENERAL COMMENTS	

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3 = Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3 = Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

WOLMAN PEBBLE COUNT FORM

County: Pittsylvania Stream ID: S-G6

Stream Name: UNT to Harpen Creek HUC Code: 03010101 Basin: Upper Roanoke

Survey Date: 8/25/2021 Surveyors: CB, BH

Type: Representative / Riffle

Y 1	D A DEVOY E	PEBB	1	D I	TD 4 3 "	T. A.	0/ ~
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	1 % % Cum
	Silt/Clay	< .062	S/C	^	35	35.00	35.00
	Very Fine	.062125		*	15	15.00	50.00
	Fine	.12525		•	4	4.00	54.00
	Medium	.255	SAND	•		0.00	54.00
	Coarse	.50-1.0		*		0.00	54.00
.0408	Very Coarse	1.0-2		^	3	3.00	57.00
.0816	Very Fine	2 -4		A	9	9.00	66.00
.1622	Fine	4 -5.7	_	*	1	1.00	67.00
.2231	Fine	5.7 - 8	GRAVEL	•	2	2.00	69.00
.3144	Medium	8 -11.3		*	2	2.00	71.00
.4463	Medium	11.3 - 16		*	3	3.00	74.00
.6389	Coarse	16 -22.6		*	3	3.00	77.00
.89 - 1.26	Coarse	22.6 - 32		-	2	2.00	79.00
1.26 - 1.77	Vry Coarse	32 - 45		*	5	5.00	84.00
1.77 -2.5	Vry Coarse	45 - 64		*	3	3.00	87.00
2.5 - 3.5	Small	64 - 90		*	4	4.00	91.00
3.5 - 5.0	Small	90 - 128	COBBLE	A	3	3.00	94.00
5.0 - 7.1	Large	128 - 180		A		0.00	94.00
7.1 - 10.1	Large	180 - 256		*		0.00	94.00
10.1 - 14.3	Small	256 - 362		*		0.00	94.00
14.3 - 20	Small	362 - 512	_	*		0.00	94.00
20 - 40	Medium	512 - 1024	BOULDER	•		0.00	94.00
40 - 80	Large	1024 -2048		•		0.00	94.00
80 - 160	Vry Large	2048 -4096		-		0.00	94.00
	Bedrock		BDRK	*	6	6.00	100.0
				Totals:	100		

RIVERMORPH PARTICLE SUMMARY

UNT to Harpen Creek

River Name: UNT to Harpen of Seach Name: S-G6 Sample Name: Representative 08/25/2021

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	35 15 4 0 0 3 9 1 2 2 3 3 2 5 3 4 3 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	35.00 15.00 4.00 0.00 0.00 3.00 9.00 1.00 2.00 2.00 3.00 3.00 3.00 4.00 3.00 4.00 3.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00	35.00 50.00 54.00 54.00 54.00 57.00 66.00 67.00 69.00 71.00 74.00 77.00 79.00 84.00 87.00 91.00 94.00 94.00 94.00 94.00 94.00 94.00 94.00 94.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.03 0.06 0.13 45 Bedrock Bedrock 35 22 30 7 0		

Total Particles = 100.

Stream Assessment Form (Form 1) Unified Stream Methodology for use in Virginia For use in wadeable channels classified as intermittent or perennial Cowardin Impact Impact **Project Name (Applicant)** Project # Locality HUC Date SAR# Class Length Factor Mountain Valley Pipeline (Mountain 22865.06 Pittslyvania R4 03010101 8/25/21 S-G6 80 1 Valley Pipeline, LLC) Stream Name and Information Name(s) of Evaluator(s) SAR Length CB, BH Spread I; UNT to Harpen Creek 80 1. Channel Condition: Assess the cross-section of the stream and prevailing condition (erosion, aggradation) Conditional Category Optimal Suboptimal Marginal Poor Severe Slightly incised, few areas of active ery little incision or active erosion; 8 Often incised, but less than Severe o Overwidened/incised. Vertically / Deeply incised (or excavated), sion or unprotected banks. Majorit of banks are stable (60-80%). laterally unstable. Likely to widen further. Majority of both banks are 100% stable banks. Vegetative Poor, Banks more stable than Sever vertical/lateral instability. Seve surface protection or natural rock, prominent (80-100%). AND/OR Stable or Poor due to lower bank slopes incision, flow contained within the Channel Vegetative protection or natural rock Erosion may be present on 40-60% of near vertical. Erosion present on 60 banks. Streambed below average Condition pankfull benches are present. Acces prominent (60-80%) AND/OR both banks. Vegetative protection on 40-60% of banks. Streambanks may banks. Vegetative protection present majority of banks vertical/undercut. to their original floodplain or fully Depositional features contribute to stability. The bankfull and low flow on 20-40% of banks, and is insufficien to prevent erosion. AND/OR 60-80% egetative protection present on less developed wide bankfull benches be vertical or undercut. AND/OR than 20% of banks, is not preventing 40-60% Sediment may be temporary transient, contribute instability. Deposition that contribute to stability, the stream is covered by sediment. Sediment is temporary / transient in nature, and contributing to instability erosion. Obvious bank sloughing present. Erosion/raw banks on 80-100%. AND/OR Aggrading channel. channel bars and transverse bars fe channels are well defined. Stream Transient sediment deposition covers less than 10% of bottom. likely has access to bankfull benches,or newly developed portions of the reach. Transient may be forming/present. AND/OR V-AND/OR V-shaped channels have than 80% of stream bed is covered by sediment covers 10-40% of the stream bottom. shaped channels have vegetative protection on > 40% of the banks and vegetative protection is present on > 40% of the banks and stable sedimen deposition, contributing to instability. Multiple thread channels and/or . depositional features which contribute deposition is absent subterranean flow CI to stability. 3 1 2.4 2 1.6 **Scores** 1.60 NOTES>> 2. RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) **Conditional Category** NOTES>> Optimal Suboptimal Marginal Poor Low Marginal: High Poor: w Suboptimal Non-maintained ligh Suboptima Lawns, mowed Riparian areas High Marginal lense herbaceou Riparian areas and maintained Low Poor: vegetation with tree stratum Non-maintained Impervious surfaces, mine with tree stratum areas, nurseries (dbh > 3 inches) ense herbaceous riparian areas (dbh > 3 inches) no-till cropland; present, with 30% vegetation with acking shrub and Tree stratum (dbh > 3 inches) presen actively grazed pasture, sparsely vegetated nonresent, with 30% spoil lands. tree stratum, hay production, ponds open water. If to 60% tree either a shrub layer or a tree Riparian with > 60% tree canopy cover. Wetlands located within the riparian to 60% tree canopy cover and uded surface anopy cover and **Buffers** row crops, active a maintained laver (dbh > 3 containing both maintained area eed lots, trails, o understory. inches) present present, tree herbaceous and recently seeded and stabilized, or other comparable conditions. with <30% tree Recent cutover stratum (dbh >3 shrub layers or a (dense canopy cover inches) present non-maintained other comparable vegetation). with <30% tree understory condition anopy cover wit maintained understory High Low High High Low Low Scores 1.5 1.2 1.1 0.85 0.75 0.6 0.5 Delineate riparian areas along each stream bank into Condition Categories and Condition Scores using the Ensure the sums 2. Determine square footage for each by measuring or estimating length and width. Calculators are provided for you of % Riparian 3. Enter the % Riparian Area and Score for each riparian category in the blocks below Blocks equal 100 100% % Riparian Area> 30% 70% Right Bank Score > 0.5 0.85 CI= (Sum % RA * Scores*0.01)/2 % Riparian Area> 30% 70% 100% Rt Bank CI > 0.75 CI Left Bank Score > 0.5 0.85 Lt Bank CI > 0.75 0.75 3. INSTREAM HABITAT: Varied substrate sizes, water velocity and depths; woody and leafy debris; stable substrate; low embeddeness; shade; undercut banks; root mats; SAV; riffle/pool complexes, stable feature Conditional Category NOTES>> Optimal Suboptimal Marginal Poor Instream Habitat/ Stable habitat elements are typically present in 30-50% of the reach and Stable habitat elements are typically present in 10-30% of the reach and Habitat elements listed above are **Available** lacking or are unstable. Habitat are adequate for maintenance of in greater than 50% of the reach. are adequate for maintenance of elements are typically present in les Cover populations. populations. than 10% of the reach. Stream Gradient CI

Scores

1.5

0.9

0.5

High / Low

1.50

1.2

	St	tream Ir	npact A	ssessn	nent Fo	rm Pag	e 2		
Project #	Project Name (App	licant)	Locality	Cowardin Class.	HUC	Date	SAR#	Impact Length	Impact Factor
22865.06	Mountain Valley Pipelin Valley Pipeline, I		Pittslyvania	R4	03010101	8/25/21	S-G6	80	1
I. CHANNE	L ALTERATION: Stream cross	sings, riprap, cond		concrete blocks, s	traightening of ch	annel, channeliza	ation, embankmer	its, spoil piles, cons	trictions,
	Negligible	Mi	nor		erate	Sev	vere	NOTES	
Channel Alteration	Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized.	Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	20-40% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines.	40 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.		by any of the chang in the parameter g 80% of banks sh	of reach is disrupted nel alterations listed juidelines AND/OR lored with gabion, r cement.		
Scores	1.5	1.3	1.1	0.9	0.7	0	.5		
	REACH (CONDITION	INDEX and S	STREAM CO	NDITION UN	IITS FOR TH	IIS REACH		
NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number. THE REACH CONDITION INDEX (RCI) >>									

RCI= (Sum of all Cl's)/5, except if stream is ephemeral RCI = (Riparian Cl/2)

COMPENSATION REQUIREMENT (CR) >>

CR = RCI X L_I X IF


INSERT PHOTOS:





CAPTION. Assessment is limited to areas within the temporary ROW.

DESCRIBE I	PROPOSED	IMPACT:
------------	----------	---------

PROVIDED UNDER SEPARATE COVER







CL STAKEOUT POINTS: S-G6 CROSS SECTION B (PIPE CL)								
	PR	POST-CROSSING						
DT LOC	NORTHING	EASTING	ELEV	VERT.	HORZ.			
PT. LOC.	NORTHING	EASTING	ELEV	DIFF.	DIFF.			
TS-L	13409163.51	2077034.43	739.64					
BS-L	13409162.73	2077037.77	735.89					
THW	13409162.46	2077038.43	735.60					
BS-R	13409162.22	2077039.17	735.97					
TS-R	13409159.83	2077050.46	746.58					
		-						

LEGEND STUDY AREA (EASEMENT) EXISTING SURVEY-LOCATED THALWEG EXISTING SURVEY-LOCATED EDGE OF WATER (AS NECESSARY) EXISTING CONTOUR LINE (MAJOR) EXISTING CONTOUR LINE (MINOR) T42.2 + EXISTING SURVEYED GROUND SHOT ELEVATION BENCHMARK POINT (WSSI)

SURVEY NOTES:

- 1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using real time DGPS. Field locations were completed on March 14, 2019.
- 2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.
- 3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).
- 4. WSSI Contour Interval = 2.0'. Interpolated from cross-section and thalweg points without additional breakline shots.
- 5. All section views shown left to right facing downstream.
- 6. Cross section B shot at location of pipe centerline (based on field stakes).

