### Reach S-GH3 (Timber Mat Crossing) Perennial Spread I Franklin County, Virginia

| Data                                    | Included                               |
|---|--|
| Photos                                  | $\checkmark$                           |
| SWVM Form                               | $\checkmark$                           |
| FCI Calculator and HGM Form             | N/A – Perennial stream (not shadeable) |
| RBP Physical Characteristics Form       | $\checkmark$                           |
| Water Quality Data                      | $\checkmark$                           |
| RBP Habitat Form                        | $\checkmark$                           |
| RBP Benthic Form                        | $\checkmark$                           |
| Benthic Identification Sheet            | $\checkmark$                           |
| Wolman Pebble Count                     | $\checkmark$                           |
| RiverMorph Data Sheet                   | $\checkmark$                           |
| USM Form (Virginia Only)                | $\checkmark$                           |
| Longitudinal Profile and Cross Sections | $\checkmark$                           |




Photo Type: US VIEW Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking NW upstream, RAH




Photo Type: DS COND Location, Orientation, Photographer Initials: Downstream at ROW/LOD looking SE downstream, RAH



Photo Type: LB CL

Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking SW at right streambank, RAH



Photo Type: RB CL Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking NE at left streambank, RAH

### **DEQ Permit #21-0416**




Photo Type: US COND Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking NW upstream, RAH



Photo Type: DS VIEW Location, Orientation, Photographer Initials: Upstream at ROW/LOD looking SE downstream, RAH

#### West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name:<br>(v2.1, Sept 2015)                             |                      | Mountain Valley Pipeline |   | IMPACT COORDINATES:<br>(in Decimal Degrees) | Lat. | 37.089745 Lor  | n79.956042                | WEATHER:   | 99% Cloud Cover                       | DATE:  | 08/31/21                       |
|--|----------------------|--------------------------|---|---|------|--|---------------------------|--|---------------------------------------|--|--------------------------------|
| IMPACT STREAM/SITE ID<br>(watershed size (acreage),                            |                      |                          | S-GH3 / 1   | 111.18 ac                                   |      | MITIGATION STREAM CLASS./SITE<br>(watershed size (acreage), unat               |                           |  |                                       | Comments:  |                                |
| STREAM IMPACT LENGTH:  | 20                   | FORM OF<br>MITIGATION:   | RESTORATION (Levels I-III)  | MIT COORDINATES:<br>(in Decimal Degrees)    | Lat. | Lor  | n.                        | PRECIPITATION PAST 48 HRS:   |                                       | Mitigation Length:   |                                |
| Column No. 1- Impact Existing  | g Condition (De      | bit)                     | Column No. 2- Mitigation Existing Co  | ondition - Baseline (Credit)                |      | Column No. 3- Mitigation Projecte<br>Post Completion (Cre                      | ed at Five Years<br>edit) | Column No. 4- Mitigation Proj<br>Post Completion (                             |                                       | Column No. 5- Mitigation Project   | ed at Maturity (Credit)        |
| Stream Classification:   | Pere                 | ennial                   | Stream Classification:  |   |      | Stream Classification:   | 0                         | Stream Classification:   | 0                                     | Stream Classification:   | 0                              |
| Percent Stream Channel SI  | lope                 | 2.76                     | Percent Stream Channel Slo  | pe  |      | Percent Stream Channel Slope   | 0                         | Percent Stream Channel SI  | ope 0                                 | Percent Stream Channel S   | lope 0                         |
| HGM Score (attach d  | ata forms):          |                          | HGM Score (attach d   | lata forms):                                |      | HGM Score (attach data   | forms):                   | HGM Score (attach da   | ata forms):                           | HGM Score (attach d  | ata forms):                    |
|  |                      | Average                  |   | Average                                     |      |  | Average                   |  | Average                               |  | Average                        |
| Hydrology<br>Biogeochemical Cycling  |                      | 0                        | Hydrology<br>Biogeochemical Cycling   | 0   |      | Hydrology<br>Biogeochemical Cycling  | 0                         | Hydrology<br>Biogeochemical Cycling  | 0                                     | Hydrology<br>Biogeochemical Cycling  | 0                              |
| Habitat<br>PART I - Physical, Chemical and                                     | Riological In dis    | ators                    | Habitat<br>PART I - Physical, Chemical and                                    |   |      | Habitat<br>PART I - Physical, Chemical and Bio                                 |                           | Habitat<br>PART I - Physical, Chemical and                                     | Rielegical Indicators                 | Habitat<br>PART I - Physical, Chemical and                                     | Rielogical Indicators          |
| PART I - Physical, Gremical and  |                      |                          | PART I - Physical, onemical and   | -   |      |  | -                         | PART I - Physical, Chemical and  | -                                     | PART I - Physical, onemical and  | -                              |
|  | Points Scale Range   | Site Score               |   | Points Scale Range Site Score               |      | Point  | s Scale Range Site Score  |  | Points Scale Range Site Score         |  | Points Scale Range Site Score  |
| PHYSICAL INDICATOR (Applies to all streams                                     | s classifications)   |                          | PHYSICAL INDICATOR (Applies to all streams of                                 | lassifications)                             |      | PHYSICAL INDICATOR (Applies to all streams classif                             | fications)                | PHYSICAL INDICATOR (Applies to all streams                                     | classifications)                      | PHYSICAL INDICATOR (Applies to all streams                                     | classifications)               |
| USEPA RBP (High Gradient Data Sheet)<br>1. Epifaunal Substrate/Available Cover |                      | 49                       | USEPA RBP (Low Gradient Data Sheet)<br>1. Epifaunal Substrate/Available Cover |   |      | USEPA RBP (High Gradient Data Sheet) 1. Epifaunal Substrate/Available Cover 0- | 20                        | USEPA RBP (High Gradient Data Sheet)<br>1. Epifaunal Substrate/Available Cover | 0-20                                  | USEPA RBP (High Gradient Data Sheet)<br>1. Epifaunal Substrate/Available Cover | 0-20                           |
| 2. Embeddedness  | 0-20                 | 13                       | 2. Pool Substrate Characterization  | 0-20  |      |  | -20                       | 2. Embeddedness  | 0-20                                  | 2. Embeddedness  | 0-20                           |
| 3. Velocity/ Depth Regime  | 0-20                 | 18                       | 3. Pool Variability   | 0-20  |      |  | -20                       | 3. Velocity/ Depth Regime  | 0-20                                  | 3. Velocity/ Depth Regime  | 0-20                           |
| 4. Sediment Deposition   | 0-20                 | 9                        | 4. Sediment Deposition  | 0-20  |      |  | -20                       | 4. Sediment Deposition   | 0-20                                  | 4. Sediment Deposition   | 0-20                           |
| 5. Channel Flow Status   | 0-20 0-1             | 16                       | 5. Channel Flow Status  | 0-20 0-1                                    |      | 5. Channel Flow Status 0-  | -20 0.1                   | 5. Channel Flow Status   | 0-20 0.1                              | 5. Channel Flow Status   | 0-20 0-1                       |
| 6. Channel Alteration  | 0-20                 | 19                       | 6. Channel Alteration   | 0-20  |      |  | -20                       | 6. Channel Alteration  | 0-20                                  | 6. Channel Alteration  | 0-20                           |
| 7. Frequency of Riffles (or bends)   | 0-20                 | 18                       | 7. Channel Sinuosity  | 0-20  |      | 7. Frequency of Riffles (or bends) 0-  | -20                       | 7. Frequency of Riffles (or bends)   | 0-20                                  | 7. Frequency of Riffles (or bends)   | 0-20                           |
| 8. Bank Stability (LB & RB)  | 0-20                 | 15                       | 8. Bank Stability (LB & RB)   | 0-20  |      |  | -20                       | 8. Bank Stability (LB & RB)  | 0-20                                  | 8. Bank Stability (LB & RB)  | 0-20                           |
| 9. Vegetative Protection (LB & RB)   | 0-20                 | 18                       | 9. Vegetative Protection (LB & RB)  | 0-20  |      |  | -20                       | 9. Vegetative Protection (LB & RB)   | 0-20                                  | 9. Vegetative Protection (LB & RB)   | 0-20                           |
| 10. Riparian Vegetative Zone Width (LB & RB)                                   | 0-20                 | 17                       | 10. Riparian Vegetative Zone Width (LB & RB)                                  | 0-20  |      |  | -20                       | <ol> <li>Riparian Vegetative Zone Width (LB &amp; RB)</li> </ol>               | 0-20                                  | <ol> <li>Riparian Vegetative Zone Width (LB &amp; RB)</li> </ol>               | 0-20                           |
| Total RBP Score  | Suboptimal           | 161                      | Total RBP Score   | Poor 0                                      |      |  | Poor 0                    | Total RBP Score  | Poor 0                                | Total RBP Score  | Poor 0                         |
| Sub-Total<br>CHEMICAL INDICATOR (Applies to Intermitter                        | nt and Perennial Str | 0.805                    | Sub-Total<br>CHEMICAL INDICATOR (Applies to Intermittent a                    | 0<br>and Perennial Streams)                 |      | Sub-Total<br>CHEMICAL INDICATOR (Applies to Intermittent and F                 | 0<br>Perennial Streams)   | Sub-Total<br>CHEMICAL INDICATOR (Applies to Intermitter                        | t and Perennial Streams)              | Sub-Total<br>CHEMICAL INDICATOR (Applies to Intermitter                        | 0                              |
| WVDEP Water Quality Indicators (General  |                      |                          | WVDEP Water Quality Indicators (General)                                      | ,   |      | WVDEP Water Quality Indicators (General)                                       | ,                         | WVDEP Water Quality Indicators (General  |                                       | WVDEP Water Quality Indicators (General  |                                |
| Specific Conductivity  |                      |                          | Specific Conductivity   |   |      | Specific Conductivity  |                           | Specific Conductivity  | · · · · · · · · · · · · · · · · · · · | Specific Conductivity  |                                |
|  | 0-90                 | 89.8                     |   | 0-90  |      | 0-   | -90                       |  | 0-90                                  |  | 0-90                           |
| <=99 - 90 points   | 1                    |                          | -11   |   |      | -11  |                           | -11  |                                       | - 11   |                                |
| pn   | 0-80 0-1             | 0.00                     | pn  | 5-90 0-1                                    |      | ph .   | .90 0-1                   | рл   | 5-90 0-1                              | ph   | 5-90 0-1                       |
| 6.0-8.0 = 80 points  | 0-00                 | 6.92                     |   |   |      | 5-   | ~~                        |  |                                       |  | ~~~                            |
| DO   | 1                    |                          | DO  |   |      | DO   |                           | DO   |                                       | DO   |                                |
| >5.0 = 30 points   | 10-30                | 5.58                     |   | 10-30                                       |      | 10   | -30                       |  | 10-30                                 |  | 10-30                          |
| Sub-Total  | - I                  | 1                        | Sub-Total   | 0   |      | Sub-Total  | 0                         | Sub-Total  | 0                                     | Sub-Total  | 0                              |
| BIOLOGICAL INDICATOR (Applies to Intermit                                      | tent and Perennial   | Streams)                 | BIOLOGICAL INDICATOR (Applies to Intermitter                                  | nt and Perennial Streams)                   |      | BIOLOGICAL INDICATOR (Applies to Intermittent                                  | and Perennial Streams)    | BIOLOGICAL INDICATOR (Applies to Interm  | ittent and Perennial Streams)         | BIOLOGICAL INDICATOR (Applies to Intern  | nittent and Perennial Streams) |
| WV Stream Condition Index (WVSCI)  |                      |                          | WV Stream Condition Index (WVSCI)   |   |      | WV Stream Condition Index (WVSCI)  |                           | WV Stream Condition Index (WVSCI)  |                                       | WV Stream Condition Index (WVSCI)  |                                |
|  | 0-100 0-1            | 64.2                     |   | 0-100 0-1                                   |      |  | 100 0-1                   |  | 0-100 0-1                             |  | 0-100 0-1                      |
| Grey Zone  |                      |                          |   | 0.00  |      |  |                           |  |                                       |  | 0.00                           |
| Sub-Total  |                      | 0.642                    | Sub-Total   | 0   |      | Sub-Total  | 0                         | Sub-Total  | 0                                     | Sub-Total  | 0                              |
| PART II - Index and U  | Jnit Score           |                          | PART II - Index and L   | Jnit Score                                  |      | PART II - Index and Unit   | Score                     | PART II - Index and U  | nit Score                             | PART II - Index and L  | Jnit Score                     |
| Index  | Linear Feet          | Unit Score               | Index   | Linear Feet Unit Score                      |      | Index Li   | near Feet Unit Score      | Index  | Linear Feet Unit Score                | Index  | Linear Feet Unit Score         |
| 0.816  | 20                   | 16.3133333               | 0   | 0 0   |      | 0  | 0 0                       | 0  | 0 0                                   | 0  | 0 0                            |
| μ  | 1                    | I                        | L   |   | ļ    | μ  |                           | L  |                                       | μ  | 1 1                            |

# PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME S-GH3                           | LOCATION Franklin County   | LOCATION Franklin County                 |  |  |  |  |  |
|---|----------------------------|--|--|--|--|--|--|
| STATION # RIVERMILE 258.5                   | STREAM CLASS Perennial     |  |  |  |  |  |  |
| LAT <u>37.089745</u> LONG <u>-79.956042</u> | RIVER BASIN Upper Roano    | ke                                       |  |  |  |  |  |
| STORET #                                    | AGENCY VADEQ               |  |  |  |  |  |  |
| INVESTIGATORS RH, RC                        |                            |  |  |  |  |  |  |
| FORM COMPLETED BY RH                        | DATE 8/31/21<br>TIME 11:38 | REASON FOR SURVEY<br>Baseline Assessment |  |  |  |  |  |

| WEATHER<br>CONDITIONS      | Now     Past 24<br>hours     Has there been a heavy rain in the last 7 days?       99 %     storm (heavy rain)<br>rain (steady rain)<br>showers (intermittent)<br>%cloud cover<br>clear/sunny     Air Temperature 28.3 ° C  |
|----------------------------|---|
| SITE LOCATION/MAP          | Dense Veg. Veg. Veg. Veg. Veg. Veg. Veg. Veg  |
| STREAM<br>CHARACTERIZATION | Stream Subsystem       Stream Type         Perennial       Intermittent       Tidal         Stream Origin       Coldwater       Warmwater         Glacial       Spring-fed       Catchment Area         Non-glacial montane       Mixture of origins       Mixture of origins |

# PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED<br>FEATURES<br>RIPARIAN<br>VEGETATION<br>(18 meter buffer) | Predominant Surrounding Landuse         Forest       Commercial         Field/Pasture       Industrial         Agricultural       Other         Residential       Other    Indicate the dominant type and record the domined th | Local Watershed NPS Pollution         ☑ No evidence       □ Some potential sources         □ Obvious sources         Local Watershed Erosion         ☑ None       □ Moderate         □ Heavy         tant species present         □ Grasses  |
|--|---|--|
| INSTREAM<br>FEATURES   | Estimated Reach Length17.98mEstimated Stream Width0.91mSampling Reach Area53.95m²Area in km² (m²x1000)0.05km²Estimated Stream Depth0.08mSurface Velocity0.25m/sec(at thalweg)0.25m/sec  | Canopy Cover       □Partly shaded □Shaded         □Partly open       □Partly shaded □Shaded         High Water Mark       0.30 m         Proportion of Reach Represented by Stream         Morphology Types         Riffle 40       % Run 30         Pool 30       %         Channelized       Yes         Dam Present       Yes                         |
| LARGE WOODY<br>DEBRIS  | LWDm <sup>2</sup><br>Density of LWDm <sup>2</sup> /km <sup>2</sup> (LWD/ read   | ch area)   |
| AQUATIC<br>VEGETATION  | Indicate the dominant type and record the domin<br>Rooted emergent<br>Floating Algae<br>Dominant species present<br>Portion of the reach with aquatic vegetation 4  | ☐Rooted floating ☐Free floating  |
| WATER QUALITY  | Temperature 21.9 D       0 C         Specific Conductance 89.8 D ms/cm         Dissolved Oxygen 5.58 D mg/L         pH 6.92 D su         Turbidity N/A         WQ Instrument Used YSI   | Water Odors         Normal/None       Sewage         Petroleum       Chemical         Fishy       Other         Water Surface Oils       Slick         Slick       Sheen       Globs         Vone       Other         Turbidity (if not measured)       Turbid         Clear       Slightly turbid       Turbid         Opaque       Stained       Other |
| SEDIMENT/<br>SUBSTRATE   | Odors       Sewage       Petroleum         Chemical       Anaerobic       None         Other       Oils       Profuse   | Deposits         □Sludge       □Sawdust       □Paper fiber       □Sand         □Relict shells       □Other         □ Epoking at stones which are not deeply embedded, are the undersides black in color?         □ Yes       ☑ No  |

| INC               | ORGANIC SUBSTRATE<br>(should add up to |                                    |                   | ORGANIC SUBSTRATE C<br>(does not necessarily add |                                   |  |  |  |  |
|-------------------|--|------------------------------------|-------------------|--|-----------------------------------|--|--|--|--|
| Substrate<br>Type | Diameter                               | % Composition in<br>Sampling Reach | Substrate<br>Type | Characteristic                                   | % Composition in<br>Sampling Area |  |  |  |  |
| Bedrock           | 10                                     |                                    | Detritus          | sticks, wood, coarse plant                       | Б                                 |  |  |  |  |
| Boulder           | > 256 mm (10")                         |                                    |                   | materials (CPOM)                                 | C                                 |  |  |  |  |
| Cobble            | 64-256 mm (2.5"-10") 30                |                                    | Muck-Mud          | black, very fine organic                         |                                   |  |  |  |  |
| Gravel            | 2-64 mm (0.1"-2.5")                    | 20                                 |                   | (FPOM)   |                                   |  |  |  |  |
| Sand              | 0.06-2mm (gritty)                      | 25                                 | Marl              | grey, shell fragments                            |                                   |  |  |  |  |
| Silt              | 0.004-0.06 mm                          | 15                                 | ]                 |  |                                   |  |  |  |  |
| Clay              | < 0.004 mm (slick)                     |                                    | ]                 |  |                                   |  |  |  |  |

#### HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME S-GH3                           | LOCATION Franklin County                           |  |  |  |  |  |
|---|--|--|--|--|--|--|
| STATION # RIVERMILE 258.5                   | STREAM CLASS Perennial                             |  |  |  |  |  |
| LAT <u>37.089745</u> LONG <u>-79.956042</u> | RIVER BASIN Upper Roanoke                          |  |  |  |  |  |
| STORET #                                    | AGENCY VADEQ                                       |  |  |  |  |  |
| INVESTIGATORS RH, RC                        |  |  |  |  |  |  |
| FORM COMPLETED BY RH                        | DATE 8/31/21 REASON FOR SURVEY Baseline Assessment |  |  |  |  |  |

|  | Habitat                                       |   | Condition   | Category  |   |  |  |  |  |
|--|---|---|---|---|---|--|--|--|--|
|  | Parameter                                     | Optimal   | Suboptimal  | Marginal  | Poor  |  |  |  |  |
|  | 1. Epifaunal<br>Substrate/<br>Available Cover | Greater than 70% of<br>substrate favorable for<br>epifaunal colonization and<br>fish cover; mix of snags,<br>submerged logs, undercut<br>banks, cobble or other<br>stable habitat and at stage<br>to allow full colonization<br>potential (i.e., logs/snags<br>that are <u>not</u> new fall and<br><u>not</u> transient). | 40-70% mix of stable<br>habitat; well-suited for<br>full colonization potential;<br>adequate habitat for<br>maintenance of<br>populations; presence of<br>additional substrate in the<br>form of newfall, but not<br>yet prepared for<br>colonization (may rate at<br>high end of scale). | 20-40% mix of stable<br>habitat; habitat<br>availability less than<br>desirable; substrate<br>frequently disturbed or<br>removed.   | Less than 20% stable<br>habitat; lack of habitat is<br>obvious; substrate<br>unstable or lacking.   |  |  |  |  |
|  | <sub>SCORE</sub> 18▼                          | 20 19 18 17 16  | 15 14 13 12 11  | 10 9 8 7 6  | 5 4 3 2 1 0   |  |  |  |  |
| ı sampling reach                             | 2. Embeddedness                               | Gravel, cobble, and<br>boulder particles are 0-<br>25% surrounded by fine<br>sediment. Layering of<br>cobble provides diversity<br>of niche space.  | Gravel, cobble, and<br>boulder particles are 25-<br>50% surrounded by fine<br>sediment.   | Gravel, cobble, and<br>boulder particles are 50-<br>75% surrounded by fine<br>sediment.   | Gravel, cobble, and<br>boulder particles are more<br>than 75% surrounded by<br>fine sediment.   |  |  |  |  |
| ed ir  | score 13 ▼                                    | 20 19 18 17 16  | 15 14 13 12 11  | 10 9 8 7 6  | 5 4 3 2 1 0   |  |  |  |  |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth<br>Regime                   | All four velocity/depth<br>regimes present (slow-<br>deep, slow-shallow, fast-<br>deep, fast-shallow).<br>(Slow is < 0.3 m/s, deep is<br>> 0.5 m.)  | Only 3 of the 4 regimes<br>present (if fast-shallow is<br>missing, score lower than<br>if missing other regimes).   | Only 2 of the 4 habitat<br>regimes present (if fast-<br>shallow or slow-shallow<br>are missing, score low).   | Dominated by 1 velocity/<br>depth regime (usually<br>slow-deep).  |  |  |  |  |
| Iram   | <sub>SCORE</sub> 18 ▼                         | 20 19 18 17 16  | 15 14 13 12 11  | 10 9 8 7 6  | 5 4 3 2 1 0   |  |  |  |  |
| P  | 4. Sediment<br>Deposition                     | Little or no enlargement<br>of islands or point bars<br>and less than 5% of the<br>bottom affected by<br>sediment deposition.   | Some new increase in bar<br>formation, mostly from<br>gravel, sand or fine<br>sediment; 5-30% of the<br>bottom affected; slight<br>deposition in pools.   | Moderate deposition of<br>new gravel, sand or fine<br>sediment on old and new<br>bars; 30-50% of the<br>bottom affected; sediment<br>deposits at obstructions,<br>constrictions, and bends;<br>moderate deposition of<br>pools prevalent. | Heavy deposits of fine<br>material, increased bar<br>development; more than<br>50% of the bottom<br>changing frequently;<br>pools almost absent due to<br>substantial sediment<br>deposition. |  |  |  |  |
|  | score 9                                       | 20 19 18 17 16  | 15 14 13 12 11  | 10 9 8 7 6  | 5 4 3 2 1 0   |  |  |  |  |
|  | 5. Channel Flow<br>Status                     | Water reaches base of<br>both lower banks, and<br>minimal amount of<br>channel substrate is<br>exposed.   | Water fills >75% of the<br>available channel; or<br><25% of channel<br>substrate is exposed.  | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.   | Very little water in<br>channel and mostly<br>present as standing pools.  |  |  |  |  |
|  | SCORE 16                                      | 20 19 18 17 16  | 15 14 13 12 11  | 10 9 8 7 6  | 5 4 3 2 1 0   |  |  |  |  |

### HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

|  | Habitat  |  | Condition  | Category   |   |  |  |  |  |
|--|--|--|--|--|---|--|--|--|--|
|  | Parameter  | Optimal  | Suboptimal   | Marginal   | Poor  |  |  |  |  |
|  | 6. Channel<br>Alteration   | Channelization or<br>dredging absent or<br>minimal; stream with<br>normal pattern.   | Some channelization<br>present, usually in areas<br>of bridge abutments;<br>evidence of past<br>channelization, i.e.,<br>dredging, (greater than<br>past 20 yr) may be<br>present, but recent<br>channelization is not<br>present.   | Channelization may be<br>extensive; embankments<br>or shoring structures<br>present on both banks;<br>and 40 to 80% of stream<br>reach channelized and<br>disrupted.   | Banks shored with gabion<br>or cement; over 80% of<br>the stream reach<br>channelized and<br>disrupted. Instream<br>habitat greatly altered or<br>removed entirely.   |  |  |  |  |
|  | SCORE 19▼  | 20 19 18 17 16   | 15 14 13 12 11   | 10 9 8 7 6   | 5 4 3 2 1 0   |  |  |  |  |
| ling reach   | 7. Frequency of<br>Riffles (or bends)  | Occurrence of riffles<br>relatively frequent; ratio<br>of distance between riffles<br>divided by width of the<br>stream <7:1 (generally 5<br>to 7); variety of habitat is<br>key. In streams where<br>riffles are continuous,<br>placement of boulders or<br>other large, natural<br>obstruction is important.           | Occurrence of riffles<br>infrequent; distance<br>between riffles divided by<br>the width of the stream is<br>between 7 to 15.  | Occasional riffle or bend;<br>bottom contours provide<br>some habitat; distance<br>between riffles divided by<br>the width of the stream is<br>between 15 to 25.   | Generally all flat water or<br>shallow riffles; poor<br>habitat; distance between<br>riffles divided by the<br>width of the stream is a<br>ratio of >25.  |  |  |  |  |
| samp   | <sub>SCORE</sub> 18▼   | 20 19 18 17 16   | 15 14 13 12 11   | 10 9 8 7 6   | 5 4 3 2 1 0   |  |  |  |  |
| Parameters to be evaluated broader than sampling reach | 8. Bank Stability<br>(score each bank)<br>Note: determine left<br>or right side by<br>facing deurostant. | Banks stable; evidence of<br>erosion or bank failure<br>absent or minimal; little<br>potential for future<br>problems. <5% of bank<br>affected.  | Moderately stable;<br>infrequent, small areas of<br>erosion mostly healed<br>over. 5-30% of bank in<br>reach has areas of erosion.   | Moderately unstable; 30-<br>60% of bank in reach has<br>areas of erosion; high<br>erosion potential during<br>floods.  | Unstable; many eroded<br>areas; "raw" areas<br>frequent along straight<br>sections and bends;<br>obvious bank sloughing;<br>60-100% of bank has<br>erosional scars.   |  |  |  |  |
| e ev   | SCORE 8  | Left Bank 10 9   | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |
| s to b   | SCORE /  | Right Bank 10 9  | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |
| Parameter  | 9. Vegetative<br>Protection (score<br>each bank)   | More than 90% of the<br>streambank surfaces and<br>immediate riparian zone<br>covered by native<br>vegetation, including<br>trees, understory shrubs,<br>or nonwoody<br>macrophytes; vegetative<br>disruption through<br>grazing or mowing<br>minimal or not evident;<br>almost all plants allowed<br>to grow naturally. | 70-90% of the<br>streambank surfaces<br>covered by native<br>vegetation, but one class<br>of plants is not well-<br>represented; disruption<br>evident but not affecting<br>full plant growth potential<br>to any great extent; more<br>than one-half of the<br>potential plant stubble<br>height remaining. | 50-70% of the<br>streambank surfaces<br>covered by vegetation;<br>disruption obvious;<br>patches of bare soil or<br>closely cropped vegetation<br>common; less than one-<br>half of the potential plant<br>stubble height remaining. | Less than 50% of the<br>streambank surfaces<br>covered by vegetation;<br>disruption of streambank<br>vegetation is very high;<br>vegetation has been<br>removed to<br>5 centimeters or less in<br>average stubble height. |  |  |  |  |
|  | SCORE 9  | Left Bank 10 9   | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |
|  | SCORE 9  | Right Bank 10 9  | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |
|  | <b>10. Riparian</b><br><b>Vegetative Zone</b><br><b>Width</b> (score each<br>bank riparian zone)         | Width of riparian zone<br>>18 meters; human<br>activities (i.e., parking<br>lots, roadbeds, clear-cuts,<br>lawns, or crops) have not<br>impacted zone.   | Width of riparian zone<br>12-18 meters; human<br>activities have impacted<br>zone only minimally.  | Width of riparian zone 6-<br>12 meters; human<br>activities have impacted<br>zone a great deal.  | Width of riparian zone <6<br>meters: little or no<br>riparian vegetation due to<br>human activities.  |  |  |  |  |
|  | SCORE 8  | Left Bank 10 9   | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |
|  | SCORE 9  | Right Bank 10 9  | 8 7 6  | 5 4 3  | 2 1 0   |  |  |  |  |

Total Score 161

#### BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME S-G      | iH3                               | LOCATION Franklin County   |  |  |  |  |  |  |  |  |
|----------------------|-----------------------------------|--|--|--|--|--|--|--|--|--|
| STATION #            | RIVERMILE                         | STREAM CLASS Perennial   |  |  |  |  |  |  |  |  |
| LAT37.089745         | LONG79.956042                     | RIVER BASIN None   | RIVER BASIN None                         |  |  |  |  |  |  |  |
| STORET #             |                                   | AGENCY VADEQ   |  |  |  |  |  |  |  |  |
| INVESTIGATORS SE     | B, KD                             |  | LOT NUMBER                               |  |  |  |  |  |  |  |
| FORM COMPLETED       | <sup>BY</sup> SB                  | DATE 9/2/2021<br>TIME 8:00 AM  | REASON FOR SURVEY<br>Baseline Assessment |  |  |  |  |  |  |  |
|                      |                                   |  |  |  |  |  |  |  |  |  |
| HABITAT TYPES        | ✓Cobble 100 <sup>°</sup> % Sn     | Indicate the percentage of each habitat type present         Cobble 100 %       Snags %         Vegetated Banks %       Sand %         Submerged Macrophytes %       Other ( |  |  |  |  |  |  |  |  |
| SAMPLE<br>COLLECTION |                                   | lected? ☑ wading ☐ fi<br>ps/kicks taken in each habitat ty<br>lags ☐ Vegetated B.  | rom bank                                 |  |  |  |  |  |  |  |
| GENERAL<br>COMMENTS  | 4 kicks in cobble.<br>vegetation. | Stream shaded by   | overhanging scrub/shrub                  |  |  |  |  |  |  |  |

#### QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton        | 0 | 1 | 2 | 3 | 4 | Slimes             | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes       | 0 | 1 | 2 | 3 | 4 | Fish               | 0 | 1 | 2 | 3 | 4 |

#### FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera        | 0 | 1 | 2 | 3 | 4 | Anisoptera  | 0 | 1 | 2 | 3 | 4 | Chironomidae  | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa        | 0 | 1 | 2 | 3 | 4 | Zygoptera   | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera   | 0 | 1 | 2 | 3 | 4 | Trichoptera   | 0 | 1 | 2 | 3 | 4 |
| Turbellaria     | 0 | 1 | 2 | 3 | 4 | Coleoptera  | 0 | 1 | 2 | 3 | 4 | Other         | 0 | 1 | 2 | 3 | 4 |
| Hirudinea       | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Oligochaeta     | 0 | 1 | 2 | 3 | 4 | Sialidae    | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Isopoda         | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Amphipoda       | 0 | 1 | 2 | 3 | 4 | Tipulidae   | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Decapoda        | 0 | 1 | 2 | 3 | 4 | Empididae   | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Gastropoda      | 0 | 1 | 2 | 3 | 4 | Simuliidae  | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
| Bivalvia        | 0 | 1 | 2 | 3 | 4 | Tabinidae   | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |
|                 |   |   |   |   |   | Culcidae    | 0 | 1 | 2 | 3 | 4 |               |   |   |   |   |   |

#### Mountain Valley Pipeline Data are not adjusted for subsampling

ECO ANALYSTS, INC.

|                      | Sample ID<br>Collection Date      | S-GH3<br>09-02-2021 |
|----------------------|-----------------------------------|---------------------|
|                      | Concerten Date                    | 00 02 2021          |
| ORDER                | GENUS/SPECIES                     | COUNT               |
| Ephemeroptera        | Baetis sp.                        | 1                   |
| Ephemeroptera        | Caenis sp.                        | 1                   |
| Ephemeroptera        |                                   | 3                   |
|                      | Maccaffertium sp.                 | 21                  |
| Ephemeroptera        |                                   | 4                   |
|                      | Leuctra sp.                       | 3                   |
|                      | Cheumatopsyche sp.                | 9                   |
|                      | Chimarra sp.                      | 4                   |
|                      | Diplectrona sp.                   | 1                   |
|                      | Hydropsyche sp.                   | 4                   |
|                      | Neophylax sp.                     | 2                   |
|                      | Calopteryx sp.                    | 2                   |
|                      | Gomphidae                         | 1                   |
|                      | Stylogomphus sp.                  | 1                   |
| •                    | Anchytarsus bicolor               | 1                   |
|                      | Ectopria sp.                      | 1                   |
|                      | Helichus sp.                      | 6                   |
|                      | Macronychus glabratus             | 1                   |
|                      | Optioservus sp.                   | 1                   |
|                      | Oulimnius sp.<br>Stenelmis sp.    | 3<br>7              |
| Megaloptera          |                                   | 2                   |
| Diptera-Chironomidae |                                   | 1                   |
| •                    |                                   |                     |
| Diptera-Chironomidae |                                   | 1                   |
| Diptera-Chironomidae |                                   | 1                   |
| Diptera-Chironomidae |                                   | 2                   |
| Diptera-Chironomidae |                                   | 4                   |
| Diptera-Chironomidae | Parametriocnemus sp.              | 2                   |
| Diptera-Chironomidae | Paraphaenocladius sp.             | 3                   |
| Diptera-Chironomidae | Paratendipes sp.                  | 1                   |
| Diptera-Chironomidae | Polypedilum sp.                   | 14                  |
| Diptera-Chironomidae |                                   | 9                   |
| Diptera-Chironomidae |                                   | 1                   |
|                      |                                   |                     |
| Diptera-Chironomidae | 5                                 | 4                   |
|                      | Thienemannimyia gr. sp.           | 47                  |
| Diptera              | Hemerodromia sp.                  | 1                   |
| Diptera              | Tabanidae                         | 2                   |
| Annelida             | Enchytraeidae                     | 1                   |
| Annelida             | Lumbriculidae                     | 1                   |
| Annelida             | Naididae                          | 23                  |
|                      | tubificoid Naididae w/o cap setae | 1                   |
|                      | Sphaeriidae                       | 3                   |
|                      | Ferrissia sp.                     | 20                  |
|                      | Lymnaeidae                        | 1                   |
| Acari                | Hygrobates sp.                    | 1                   |
| Other Organisms      |                                   | 1                   |
|                      | TOTAL                             | 224                 |

Mountain Valley Pipeline WV SCI Metrics

## ECO ANALYSTS, INC.

| Sample ID<br>Collection Date  |   |
|---|---|
| WVSCI Metric Values<br>Total taxa<br>EPT taxa<br>% EPT<br>% Chironomidae<br>% 2 Dominant<br>HBI | 26<br>8<br>23.7<br>40.2<br>51.3<br>5.62       |
| WVSCI Metric Scores<br>Total taxa<br>EPT taxa<br>% EPT<br>% Chironomidae<br>% 2 Dominant<br>HBI | 123.8<br>61.5<br>25.7<br>60.4<br>76.0<br>61.7 |
| WVSCI Metric Scores<br>Total taxa<br>EPT taxa<br>% EPT<br>% Chironomidae<br>% 2 Dominant<br>HBI | 100.0<br>61.5<br>25.7<br>60.4<br>76.0<br>61.7 |
| WVSCI Total Score   | 64.2  |

#### WVSCI Thresholds

Unimpaired = > 68.00 Gray Zone = 60.61 to 68.00

Impaired = <60.61

#### WOLMAN PEBBLE COUNT FORM

Basin:

County: Franklin County Stream Name: UNT to Teels Creek HUC Code: 03010101 Survey Date: 8/31/2021 Surveyors: RH, RC Type: Representative

Inches

Stream ID:

Upper Roanoke

% Cum

S-GH3

PEBBLE COUNT PARTICLE Millimeters Particle Total # Item % Count < .062 S/C ٠ I 15 15.00

|             | Silt/Clay   | < .062     | S/C     | •<br>•   | 15  | 15.00 | 15.00  |
|-------------|-------------|------------|---------|----------|-----|-------|--------|
|             | Very Fine   | .062125    |         | ÷        | 1   | 1.00  | 16.00  |
|             | Fine        | .12525     | 1       | ÷        | 5   | 5.00  | 21.00  |
|             | Medium      | .255       | SAND    | <b>^</b> | 3   | 3.00  | 24.00  |
|             | Coarse      | .50-1.0    | -       | <b></b>  | 2   | 2.00  | 26.00  |
| .0408       | Very Coarse | 1.0-2      | 1       | <b></b>  | 4   | 4.00  | 30.00  |
| .0816       | Very Fine   | 2 -4       |         |          | 1   | 1.00  | 31.00  |
| .1622       | Fine        | 4 -5.7     | 1       | <b>^</b> | 2   | 2.00  | 33.00  |
| .2231       | Fine        | 5.7 - 8    | 1       | ÷        | 6   | 6.00  | 39.00  |
| .3144       | Medium      | 8 -11.3    | 1       | <b></b>  | 6   | 6.00  | 45.00  |
| .4463       | Medium      | 11.3 - 16  | GRAVEL  | ÷        |     | 0.00  | 45.00  |
| .6389       | Coarse      | 16 -22.6   | 1       | ÷        | 1   | 1.00  | 46.00  |
| .89 - 1.26  | Coarse      | 22.6 - 32  | 1       | ▲<br>▼   | 2   | 2.00  | 48.00  |
| 1.26 - 1.77 | Vry Coarse  | 32 - 45    | 1       | <b></b>  | 4   | 4.00  | 52.00  |
| 1.77 -2.5   | Vry Coarse  | 45 - 64    | 1       | ÷        | 8   | 8.00  | 60.00  |
| 2.5 - 3.5   | Small       | 64 - 90    |         |          | 4   | 4.00  | 64.00  |
| 3.5 - 5.0   | Small       | 90 - 128   | 1       | ÷        | 14  | 14.00 | 78.00  |
| 5.0 - 7.1   | Large       | 128 - 180  | COBBLE  | ÷        | 3   | 3.00  | 81.00  |
| 7.1 - 10.1  | Large       | 180 - 256  | 1       | ÷        | 5   | 5.00  | 86.00  |
| 10.1 - 14.3 | Small       | 256 - 362  |         | ÷        |     | 0.00  | 86.00  |
| 14.3 - 20   | Small       | 362 - 512  | 1       | ÷        |     | 0.00  | 86.00  |
| 20 - 40     | Medium      | 512 - 1024 | BOULDER | <br>▼    |     | 0.00  | 86.00  |
| 40 - 80     | Large       | 1024 -2048 | 1       | <br>↓    |     | 0.00  | 86.00  |
| 80 - 160    | Vry Large   | 2048 -4096 | 1       | ▲<br>▼   |     | 0.00  | 86.00  |
|             | Bedrock     |            | BDRK    |          | 14  | 14.00 | 100.00 |
|             |             |            | 1       | Totals:  | 100 |       |        |

\_\_\_\_\_

| River Name:<br>Reach Name:<br>Sample Name:<br>Survey Date:  | Representative  |  |   |  |
|---|---|--|---|--|
| Size (mm)   | TOT #   | ITEM %   | CUM %   |  |
| 0 - 0.062<br>0.062 - 0.125<br>0.125 - 0.25<br>0.25 - 0.50<br>0.50 - 1.0<br>1.0 - 2.0<br>2.0 - 4.0<br>4.0 - 5.7<br>5.7 - 8.0<br>8.0 - 11.3<br>11.3 - 16.0<br>16.0 - 22.6<br>22.6 - 32.0<br>32 - 45<br>45 - 64<br>64 - 90<br>90 - 128<br>128 - 180<br>180 - 256<br>256 - 362<br>362 - 512<br>512 - 1024<br>1024 - 2048<br>Bedrock | 15<br>1<br>5<br>3<br>2<br>4<br>1<br>2<br>6<br>6<br>0<br>1<br>2<br>4<br>8<br>4<br>14<br>3<br>5<br>0<br>0<br>0<br>0<br>14 | $\begin{array}{c} 15.00\\ 1.00\\ 5.00\\ 3.00\\ 2.00\\ 4.00\\ 1.00\\ 2.00\\ 6.00\\ 0.00\\ 1.00\\ 2.00\\ 4.00\\ 1.00\\ 2.00\\ 4.00\\ 1.00\\ 3.00\\ 5.00\\ 0.00\\ 0.00\\ 0.00\\ 0.00\\ 0.00\\ 14.00\\ 14.00\end{array}$ | $ \begin{array}{c} 16.00\\ 21.00\\ 24.00\\ 26.00\\ 30.00\\ 31.00\\ 33.00\\ 39.00\\ 45.00\\ 45.00\\ 45.00\\ 45.00\\ 45.00\\ 45.00\\ 60.00\\ 64.00\\ 52.00\\ 60.00\\ 86.00\\ 8$ |  |
| D16 (mm)<br>D35 (mm)<br>D50 (mm)<br>D84 (mm)<br>D95 (mm)<br>D100 (mm)<br>Silt/Clay (%)<br>Sand (%)<br>Gravel (%)<br>Gravel (%)<br>Boulder (%)<br>Bedrock (%)<br>Total Particles = 100   | 0.13<br>6.47<br>38.5<br>225.6<br>Bedrock<br>15<br>15<br>30<br>26<br>0<br>14   |  |   |  |

|  |   |  | F  | For use in wadea  | able channels cla  | ssified as interm   | nittent or perenni   | al   |  |  |          |
|--|---|--|--|---|--|---|--|--|--|--|----------|
| Project #  | Project Nan   | me (Applicant  |  | Locality  | Cowardin<br>Class.   | HUC   | Date   | SAR #  | Impact<br>Length   | Impact<br>Factor   |          |
| 22865.06   | Mountain Valley<br>Valley Pip   | v Pipeline (Mo<br>ipeline, LLC)  |  | Franklin<br>County  | R3   | 03010101  | 8/31/21  | S-GH3  | 20   | 1  |          |
| Nam  | e(s) of Evaluator(s)  |  |  | and Informa   | tion   | •   |  | •  | SAR Length   |  |          |
|  | RH, RC  | Spre   | ead I; UNI   | ۲ to Teels Cre  | ek   |   |  |  | 74   |  |          |
| Channel C  | ondition: Assess the c  | cross-section of th  | the stream ar  | nd prevailing cond  | dition (erosion, ago   | gradation)  |  |  |  |  |          |
|  | 1   |  |  |   | Conditional Catego   | ory   |  |  | Sov  | <u></u>  |          |
|  | Optimal   |  | Subop  |   | IVIAI  | ginal   |  | Dor  | Seve   |  |          |
|  | Very little incision or active e  |  |  | w areas of active   | · · ·  | less than Severe or   |  | cised. Vertically /  | Deeply incised (   |  |          |
| Channel<br>Condition   | <ul> <li>100% stable banks. Vegetar<br/>protection or natural rock,</li> <li>(80-100%). AND/OR Stable<br/>bankfull benches are presen<br/>to their original floodplain<br/>developed wide bankfull ben<br/>channel bars and transverse<br/>Transient sediment deposit<br/>less than 10% of both</li> </ul>  | prominent of<br>e point bars / Vege<br>ent. Access p<br>in or fully Dep<br>enches. Mid-<br>se bars few.<br>ition covers has<br>ottom. new  | of banks are sta<br>etative protecti<br>prominent (60-<br>positional featu<br>bility. The ban<br>nels are well de<br>s access to ban<br>wly developed f  | ted banks. Majority<br>able (60-80%).<br>ion or natural rock<br>80%) AND/OR<br>ures contribute to<br>kfull and low flow<br>fined. Stream likely<br>nkfull benches,or<br>floodplains along<br>each. Transient  | or Poor due to lo<br>Erosion may be pro<br>both banks. Veget<br>40-60% of banks. S<br>vertical or unde<br>40-60% Sediment<br>transient, contr<br>Deposition that co          | stable than Severe<br>ower bank slopes.<br>esent on 40-60% of<br>tative protection on<br>Streambanks may be<br>ercut. AND/OR<br>may be temporary /<br>fibute instability.<br>ntribute to stability,<br>resent. AND/OR V-  | further. Majority of<br>vertical. Erosion pro-<br>banks. Vegetative<br>on 20-40% of bank<br>to prevent erosion.<br>the stream is cov<br>Sediment is temp<br>nature, and contri   |  | vertical/lateral ins<br>incision, flow containe<br>Streambed below ave<br>majority of banks v<br>Vegetative protection<br>than 20% of banks,<br>erosion. Obvious<br>present. Erosion/raw<br>AND/OR Aggrading<br>than 80% of stream | ed within the banks.<br>erage rooting depth,<br>vertical/undercut.<br>on present on less<br>, is not preventing<br>bank sloughing<br>banks on 80-100%.<br>g channel. Greater |          |
|  |   | sedime   | ent covers 10<br>botto   | )-40% of the stream   | l '  | s have vegetative<br>% of the banks and   | 40% of the banks a   | tion is present on > and stable sediment   | deposition, contribu<br>Multiple thread c  |  |          |
|  |   |  |  |   |  | es which contribute<br>ability.   | · ·  | n is absent.   | subterrane   | ean flow.  | С        |
| Scores<br>NOTES>>  | BLIEFERS: Access is   | both bonk's 100 f  | 2.   |   | to sta   | ability.<br><b>2</b>  | 1  | .6   | subterrane<br>1  | ean flow.  |          |
| NOTES>>  | 3 BUFFERS: Assess b Optimal   | both bank's 100 fc   | <b>2.</b>  | areas along the er<br>ditional Cate   | ntire SAR. (rough  | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:   | f length & width ma<br>Po<br>High Poor: Lawns,   | .6<br>ay be acceptable)  | subterrane<br>1  | ean flow.  |          |
| NOTES>>  |   | nes) present,<br>by cover.<br>the riparian<br>tree ca<br>and cou<br>herba<br>shrub<br>non-r  | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both   | areas along the er<br>ditional Cate   | to stand<br>to stand<br>tire SAR. (rough<br>gory<br>Marg<br>Marg<br>Marg<br>Marg<br>vegetation with<br>either a shrub layer<br>or a tree layer (dbh<br>> 3 inches)           | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,  | f length & width ma<br>f length & width ma<br>Point A stabilized, or other<br>comparable   | A be acceptable)<br><b>Dor</b><br><b>Low Poor:</b><br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable                        | 1  | ean flow.  | C<br>2.4 |
| NOTES>>  | <b>Optimal</b><br>Tree stratum (dbh > 3 inche<br>with > 60% tree canopy<br>Wetlands located within th   | nes) present,<br>by cover.<br>the riparian<br>tree ca<br>and cou<br>herba<br>shrub<br>non-r<br>und   | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both<br>baceous and<br>b layers or a<br>-maintained  | areas along the er<br>ditional Categotimal<br>Dimal<br>Low Suboptimal:<br>Riparian areas with<br>tree stratum (dbh ><br>3 inches) present,<br>with 30% to 60%<br>tree canopy cover<br>and a maintained<br>understory. Recent<br>cutover (dense                      | to sta<br>to sta<br>tire SAR. (rough<br>gory<br>Marg<br>Marg<br>Marg<br>vegetation with<br>either a shrub layer<br>or a tree layer (dbh<br>> 3 inches)<br>present, with <30% | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,<br>hay production,<br>ponds, open water.<br>If present, tree<br>stratum (dbh >3<br>inches) present,<br>with <30% tree<br>canopy cover with<br>maintained                               | f length & width ma<br>f length & width ma<br>Point A stabilized, or other<br>comparable   | A be acceptable)<br><b>Dor</b><br><b>Low Poor:</b><br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable                        | 1  | ean flow.  |          |
| NOTES>>  | <b>Optimal</b><br>Tree stratum (dbh > 3 inche<br>with > 60% tree canopy<br>Wetlands located within th   | nes) present,<br>by cover.<br>the riparian<br>tree ca<br>and cou<br>herba<br>shrub<br>non-r<br>und   | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both<br>baceous and<br>b layers or a<br>-maintained<br>nderstory.  | areas along the er<br>ditional Categotimal<br>Dimal<br>Low Suboptimal:<br>Riparian areas with<br>tree stratum (dbh ><br>3 inches) present,<br>with 30% to 60%<br>tree canopy cover<br>and a maintained<br>understory. Recent<br>cutover (dense<br>vegetation).      | to sta<br>to sta<br>to sta<br>tire SAR. (rough<br>gory<br>Marg<br>Marg<br>Marg<br>Marg<br>sta<br>sta<br>sta<br>sta<br>sta<br>sta<br>sta<br>sta<br>sta<br>sta                 | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,<br>hay production,<br>ponds, open water.<br>If present, tree<br>stratum (dbh >3<br>inches) present,<br>with <30% tree<br>canopy cover with<br>maintained<br>understory.                | f length & width ma<br>f length & width ma<br>Particular<br>High Poor: Lawns,<br>mowed, and<br>maintained areas,<br>nurseries; no-till<br>cropland; actively<br>grazed pasture,<br>sparsely vegetated<br>non-maintained<br>area, recently<br>seeded and<br>stabilized, or other<br>comparable<br>condition.  | ay be acceptable)<br><b>Dor</b><br>Low Poor:<br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable<br>conditions.               | 1  | ean flow.  |          |
| NOTES>><br>RIPARIAN<br>Riparian Buffers<br>Scores<br>Delineate ripa<br>Determine squ               | Optimal         Tree stratum (dbh > 3 inche with > 60% tree canopy Wetlands located within th areas.         Wetlands located within th areas.         1.5         trian areas along each stree uare footage for each by mare footage | hes) present,<br>by cover.<br>the riparian<br>tree str<br>3 inch<br>with 3<br>tree ca<br>and cou<br>herba<br>shrub<br>non-r<br>und<br>ream bank into Co<br>measuring or estir<br>for each riparian ca<br>20%                         | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both<br>baceous and<br>b layers or a<br>-maintained<br>nderstory.<br>High<br>1.2<br>ondition Cate<br>imating lengt<br>asom | areas along the er<br>ditional Categotimal<br>Low Suboptimal:<br>Riparian areas with<br>tree stratum (dbh ><br>3 inches) present,<br>with 30% to 60%<br>tree canopy cover<br>and a maintained<br>understory. Recent<br>cutover (dense<br>vegetation).<br>Low<br>1.1 | to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>sta<br>sta<br>titon Scores using   | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,<br>hay production,<br>ponds, open water.<br>If present, tree<br>stratum (dbh >3<br>inches) present,<br>with <30% tree<br>canopy cover with<br>maintained<br>understory.<br>Low<br>0.75 | f length & width ma<br>f length & width ma<br>Provide the second of the seco | ay be acceptable)<br><b>Dor</b><br>Low Poor:<br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable<br>conditions.               | 1  | ean flow.  |          |
| NOTES>><br>RIPARIAN<br>Riparian Buffers<br>Scores<br>Delineate ripa<br>Determine squ Enter the % R | Optimal         Tree stratum (dbh > 3 inche with > 60% tree canopy Wetlands located within th areas.         Wetlands located within th areas.         1.5         trian areas along each stree uare footage for each by mare footage | hes) present,<br>by cover.<br>the riparian<br>tree str<br>3 inch<br>with 3<br>tree ca<br>and cou<br>herba<br>shrub<br>non-r<br>und<br>ream bank into Co<br>measuring or estir<br>for each riparian ca<br>20%                         | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both<br>baceous and<br>b layers or a<br>-maintained<br>nderstory.<br>High<br>1.2<br>ondition Cate                          | areas along the er<br>ditional Categotimal<br>Low Suboptimal:<br>Riparian areas with<br>tree stratum (dbh ><br>3 inches) present,<br>with 30% to 60%<br>tree canopy cover<br>and a maintained<br>understory. Recent<br>cutover (dense<br>vegetation).<br>Low<br>1.1 | to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>sta<br>sta<br>titon Scores using   | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,<br>hay production,<br>ponds, open water.<br>If present, tree<br>stratum (dbh >3<br>inches) present,<br>with <30% tree<br>canopy cover with<br>maintained<br>understory.<br>Low<br>0.75 | f length & width ma<br>f length & width ma<br>Provide the second of the seco | .6<br>ay be acceptable)<br>Cor<br>Low Poor:<br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable<br>conditions.<br>Low<br>100% | NOTES>>  |  |          |
| NOTES>><br>RIPARIAN<br>Riparian Buffers<br>Scores<br>Delineate ripa<br>Determine squ Enter the % R | Optimal         Tree stratum (dbh > 3 inche with > 60% tree canopy Wetlands located within th areas.         Wetlands located within th areas.         1.5         trian areas along each stree uare footage for each by mare footage  | hes) present,<br>by cover.<br>the riparian<br>the riparian<br>tree str<br>3 inch<br>with 3<br>tree ca<br>and cor<br>herba<br>shrub<br>non-r<br>und<br>ream bank into Cor<br>measuring or estir<br>for each riparian ca<br>20%<br>0.5 | foot riparian a<br>Con<br>Suboptimal:<br>ian areas with<br>stratum (dbh ><br>hes) present,<br>30% to 60%<br>canopy cover<br>ontaining both<br>baceous and<br>b layers or a<br>-maintained<br>nderstory.<br>High<br>1.2<br>ondition Cate<br>imating lengt<br>asom | areas along the er<br>ditional Categotimal<br>Low Suboptimal:<br>Riparian areas with<br>tree stratum (dbh ><br>3 inches) present,<br>with 30% to 60%<br>tree canopy cover<br>and a maintained<br>understory. Recent<br>cutover (dense<br>vegetation).<br>Low<br>1.1 | to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>to sta<br>sta<br>sta<br>titon Scores using   | ability.<br>2<br>measurements o<br>ginal<br>Low Marginal:<br>Non-maintained,<br>dense herbaceous<br>vegetation, riparian<br>areas lacking shrub<br>and tree stratum,<br>hay production,<br>ponds, open water.<br>If present, tree<br>stratum (dbh >3<br>inches) present,<br>with <30% tree<br>canopy cover with<br>maintained<br>understory.<br>Low<br>0.75 | f length & width ma<br>f length & width ma<br>Provide the second of the seco | .6<br>ay be acceptable)<br>Cor<br>Low Poor:<br>Impervious<br>surfaces, mine<br>spoil lands,<br>denuded surfaces,<br>row crops, active<br>feed lots, trails, or<br>other comparable<br>conditions.<br>Low<br>100% | 1  |  |          |

|                                |  | Conditional Category  |  |   |                 |      |  |  |  |  |
|--------------------------------|--|---|--|---|-----------------|------|--|--|--|--|
| Instream                       | Optimal Suboptimal Marginal Poor   |   |  |   |                 |      |  |  |  |  |
| Habitat/<br>Available<br>Cover | Habitat elements are typically present in greater than 50% of the reach. | Stable habitat elements are typically present in 30-50% of the reach and are adequate for maintenance of populations. | Stable habitat elements are typically<br>present in 10-30% of the reach and are<br>adequate for maintenance of<br>populations. | Habitat elements listed above are<br>lacking or are unstable. Habitat<br>elements are typically present in less<br>than 10% of the reach. | Stream Gradient | CI   |  |  |  |  |
| Scores                         | 1.5  | 1.2   | 0.9  | 0.5   | High / Low      | 1.50 |  |  |  |  |

Reach R3-R4

File: https://tetratechinc.sharepoint.com/teams/MVPStreamWetlandAssessment/Shared Documents/General/01. Virginia Field Data Management/03. Preliminary QAQC (working files)/S-GH3\_20211006SS/9. S-GH3\_USM\_20211006SS.xlsx

| Project #             | Project Name (App  | licant)  | Locality   | Cowardin<br>Class.   | HUC  | Date   | SAR #   | Impact<br>Length         | Impact<br>Factor |            |
|-----------------------|--|--|--|--|--|--|---|--------------------------|------------------|------------|
| 22865.06              | Mountain Valley Pipeline (MountainFranklinValley Pipeline, LLC)County  |  |  | R3   | 03010101   | 8/31/21  | S-GH3   | 20                       | 1                |            |
| . CHANNEI             | ALTERATION: Stream crossin   | igs, riprap, concret   | e, gabions, or con   | ncrete blocks, strai   | ghtening of chann  | el, channelization   |   | •                        | ons, livestock   |            |
|                       |  |  | Conditiona   | al Category  |  |  |   | NOTES>>                  |                  |            |
|                       | Negligible   | Mir  | nor  |  | erate  | Sev  | /ere  |                          |                  |            |
| Channel<br>Alteration | Channelization, dredging, alteration, or<br>hardening absent. Stream has an<br>unaltered pattern or has naturalized. | the channel<br>alterations listed in<br>the parameter<br>guidelines. | the channel<br>alterations listed in<br>the parameter<br>guidelines. | is disrupted by any<br>of the channel<br>alterations listed in<br>the parameter<br>guidelines. If<br>stream has been<br>channelized,<br>normal stable<br>stream meander<br>pattern has not<br>recovered. | the parameter<br>guidelines. If<br>stream has been<br>channelized,<br>normal stable<br>stream meander<br>pattern has not<br>recovered. | by any of the chanr<br>in the parameter g<br>80% of banks sh<br>riprap, or | of reach is disrupted<br>nel alterations listed<br>uidelines AND/OR<br>ored with gabion,<br>cement. |                          |                  | CI         |
| Scores                | 1.5  | 1.3  | 1.1  | 0.9  | 0.7  | 0  | .5  |                          |                  | 1.10       |
|                       | REACH  | CONDITION  | INDEX and S  | STREAM CO  | NDITION UN   | ITS FOR THI  | S REACH   |                          |                  |            |
| OTE: The Cls a        | nd RCI should be rounded to 2 decir  | mal places. The Cl   | R should be round  | led to a whole nun   | nber.  |  | THE REACH   | H CONDITION IN           | DEX (RCI) >>     | 1.16       |
|                       |  |  |  |  |  | RCI= (Sum of   | all CI's)/5, exce   | ept if stream is ep      | hemeral RCI = (F | ≀iparian C |
|                       |  |  |  |  |  |  | COMPENSA  | TION REQUIRE             | MENT (CR) >>     | 23         |
|                       |  |  |  |  |  |  | CR = RC   | CI X L <sub>I</sub> X IF |                  |            |



37.08968°N


CAPTION. Assessment is limited to areas within the temporary ROW.


DESCRIBE PROPOSED IMPACT:

PROVIDED UNDER SEPARATE COVER


Reach R3-R4

File: https://tetratechinc.sharepoint.com/teams/MVPStreamWetlandAssessment/Shared Documents/General/01. Virginia Field Data Management/03. Preliminary QAQC (working files)/S-GH3\_20211006SS/9. S-GH3\_USM\_20211006SS.xlsx





| CL STAKEOUT POINTS: S-GH3 CROSS SECTION B (PIPE CL) |             |            |         |       |       |  |  |  |  |
|---|-------------|------------|---------|-------|-------|--|--|--|--|
|   | PRE         | POST-CF    | ROSSING |       |       |  |  |  |  |
|   | NODTUINC    | FACTINIC   |         | VERT. | HORZ. |  |  |  |  |
| PT. LOC.  | NORTHING    | EASTING    | ELEV    | DIFF. | DIFF. |  |  |  |  |
| TS-L  | 13468929.23 | 1944811.21 | 1157.56 |       |       |  |  |  |  |
| BS-L  | 13468925.82 | 1944808.30 | 1155.90 |       |       |  |  |  |  |
| THW   | 13468924.02 | 1944806.17 | 1155.44 |       |       |  |  |  |  |
| BS-R  | 13468922.09 | 1944804.05 | 1155.96 |       |       |  |  |  |  |
| TS-R  | 13468919.30 | 1944802.45 | 1159.72 |       |       |  |  |  |  |



SURVEY NOTES:

1. This map has been oriented to NAD 1983 UTM ZONE 17N, and vertically to The North American Vertical Datum of 1988 (NAVD 88), using a Real Time Network (RTN) GPS. Field locations were completed on September 12, 2018.

2. Monumentation, including traverse stations and fly points, shown on this drawing should be used to orient any future boundary, topographic, or location survey.

3. Easement lines shown on plan view were provided by Mountain Valley Pipeline (MVP).

4. WSSI Contour Interval = 2.0'. Contours within the channel were interpolated using stream channel breaklines (i.e. top of slopes, toe of slopes, thalweg) and cross-sectional points. Contours outside the channel were interpolated using cross-sectional spot shots.

5. All section views shown are left to right facing downstream.

6. Cross-section B shot at location of pipe centerline (based on best professional judgement).

