Reach S-GH44 (Timber Mat Crossing) Perennial Spread I Franklin County, Virginia

| Data | Included |
|---|--|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | N/A – Perennial stream (not shadeable, slope |
| | <4%) |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | \checkmark |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | \checkmark |
| Wolman Pebble Count | \checkmark |
| RiverMorph Data Sheet | \checkmark |
| USM Form (Virginia Only) | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |

Spread I Stream S-GH44 (Timber Mat Crossing) Franklin County



Photo Type: US VIEW Location, Orientation, Photographer Initials: Downstream at ROW/LOC looking SW upstream, VM




Photo Type: DS COND DS Location, Orientation, Photographer Initials: Downstream at ROW/LOC looking NW downstream, VM

Spread I Stream S-GH44 (Timber Mat Crossing) Franklin County



Photo Type: LB CL Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking SE at left streambank, VM



Photo Type: RB CL Location, Orientation, Photographer Initials: On thalweg at pipe centerline looking SW at right streambank, VM

DEQ Permit #21-0416

Spread I Stream S-GH44 (Timber Mat Crossing) Franklin County



Photo Type: US COND Location, Orientation, Photographer Initials: Upstream at ROW/LOC looking SE upstream, VM




Photo Type: DS VIEW Location, Orientation, Photographer Initials: Upstream at ROW/LOC looking NW downstream, VM

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| | USACE FILE NO./ Project Name: (v2.1, Sept 2015) | Mountain | Valley Pipeline | IMPACT COORDINATES: (in Decimal Degrees) | Lat. | 37.028392 | Lon. | -79.773359 | WEATHER: | Sunny | DATE: | August 26, 2021 |
|---|--|-----------------------------|---|---|------|--|--------------------|---------------|--|--------------------------------|---|--------------------------------|
| Image: Description of the strength of t | | | S-GH44; | ; 56.67 ac | | | | | | | Comments: | |
| | STREAM IMPACT LENGTH: | | RESTORATION (Levels I-III) | | Lat. | | Lon. | | PRECIPITATION PAST 48 HRS: | No | Mitigation Length: | |
| | Column No. 1- Impact Existing | g Condition (Debit) | Column No. 2- Mitigation Existing Co | ondition - Baseline (Credit) | | | | Years | | | Column No. 5- Mitigation Project | ted at Maturity (Credit) |
| | Stream Classification: | Perennial | Stream Classification: | | | Stream Classification: | | 0 | Stream Classification: | 0 | Stream Classification: | 0 |
| | Percent Stream Channel SI | ope 2.76 | Percent Stream Channel Slo | ope | | Percent Stream Channel S | lope | 0 | Percent Stream Channel St | ope 0 | Percent Stream Channel S | Slope 0 |
| | HGM Score (attach da | ata forms): | HGM Score (attach o | data forms): | | HGM Score (attach | data forms): | | HGM Score (attach d | ata forms): | HGM Score (attach o | lata forms): |
| | | Average | | Average | | | | Average | | Average | | Average |
| | | | | | | | | | | | | |
| | Habitat | 0 | Habitat | 0 | | Habitat | | | Habitat | 0 | Habitat | 0 |
| | PART I - Physical, Chemical and | | PART I - Physical, Chemical and | - | | PART I - Physical, Chemical a | - | | PART I - Physical, Chemical and | - | PART I - Physical, Chemical and | |
| | | | | | | | | Site Score | | | | |
| | PHYSICAL INDICATOR (Applies to all streams | s classifications) | PHYSICAL INDICATOR (Applies to all streams of | classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | PHYSICAL INDICATOR (Applies to all streams | s classifications) | PHYSICAL INDICATOR (Applies to all stream | s classifications) |
| | | | USEPA RBP (Low Gradient Data Sheet) | | | | 1 | | | | | T T |
| | | | | | | | | | | | | |
| | | 0.20 11 | | | | | | | | | | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| | | | 8. Bank Stability (LB & RB) | | | | | | | | | |
| Total RBP Score Por 0 Sub-Total Total RBP Score 0< | | | | | | | | | | | | |
| | | | | | | | | 0 | | | | |
| WDEP Water Quality indicators (General) WDEP Water Quality indicator | Total Ttbi Oobio | | | Poor U | | | POOT | 0 | | Poor 0 | | Poor U |
| Specific Conductivity | CHEMICAL INDICATOR (Applies to Intermitten | | CHEMICAL INDICATOR (Applies to Intermittent | and Perennial Streams) | | CHEMICAL INDICATOR (Applies to Intermittee | nt and Perennial S | treams) | CHEMICAL INDICATOR (Applies to Intermitter | nt and Perennial Streams) | CHEMICAL INDICATOR (Applies to Intermitte | nt and Perennial Streams) |
| | WVDEP Water Quality Indicators (General |) | WVDEP Water Quality Indicators (General) | | | WVDEP Water Quality Indicators (General |) | | WVDEP Water Quality Indicators (General | 1) | WVDEP Water Quality Indicators (General | I) |
| pic pic <td>Specific Conductivity</td> <td></td> <td>Specific Conductivity</td> <td></td> <td></td> <td>Specific Conductivity</td> <td></td> <td></td> <td>Specific Conductivity</td> <td></td> <td>Specific Conductivity</td> <td></td> | Specific Conductivity | | Specific Conductivity | | | Specific Conductivity | | | Specific Conductivity | | Specific Conductivity | |
| pH < | | 0-90 67.1 | | 0-90 | | 1 | 0-90 | | | 0-90 | | 0-90 |
| b. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. | <=99 - 90 points | | - | | | nH | | | | | | |
| b. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. 0. | pri | 0-1 | pri | 0-1 | | pri | 0-1 | | pri | 0-1 | pri l | 0.1 |
| DO Set Total DO Set Total DO DO <th< td=""><td>6 0-8 0 = 80 points</td><td>0-80 6.5</td><td></td><td>5-90</td><td></td><td>1</td><td>5-90</td><td></td><td></td><td>5-90</td><td></td><td>5-90</td></th<> | 6 0-8 0 = 80 points | 0-80 6.5 | | 5-90 | | 1 | 5-90 | | | 5-90 | | 5-90 |
| 10-30 0.8.8 6.8.9 Sub_Total 0.10 0.00 <td></td> <td></td> <td>DO</td> <td></td> <td></td> <td>DO</td> <td>•</td> <td></td> <td>DO</td> <td></td> <td>DO</td> <td></td> | | | DO | | | DO | • | | DO | | DO | |
| ×6.0 = 3D points ✓ </td <td></td> <td>10.30 6.95</td> <td></td> <td>10-30</td> <td></td> <td></td> <td>10-30</td> <td></td> <td></td> <td>10-30</td> <td></td> <td>10.30</td> | | 10.30 6.95 | | 10-30 | | | 10-30 | | | 10-30 | | 10.30 |
| BIOLOGICAL INDICATOR (Applies to Intermittent and Permital Stream) BIOLOGICAL INDICATOR (Applies to Intermittent a | | | | | | | 10-50 | | | | | |
| W Stream Condition Index (WVSC) W Stream Conditindex (WVSC) W Stream | | | | 0 | | | | | | 0 | | 0 |
| Good 0-10 0-1 0-10 0-1 0-10 0-1 0-10 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 0-1 0-100 <td></td> <td>tent and Perennial Streams)</td> <td></td> <td>ent and Perennial Streams)</td> <td></td> <td></td> <td>nittent and Peren</td> <td>nial Streams)</td> <td></td> <td>nittent and Perennial Streams)</td> <td></td> <td>nittent and Perennial Streams)</td> | | tent and Perennial Streams) | | ent and Perennial Streams) | | | nittent and Peren | nial Streams) | | nittent and Perennial Streams) | | nittent and Perennial Streams) |
| Good | WV Stream Condition Index (WVSCI) | 0,100 0,1 70 7 | WV Stream Condition Index (WVSCI) | 0.100 0.1 | | WV Stream Condition Index (WVSCI) | 0.100 0.1 | | WV Stream Condition Index (WVSCI) | 0.100 0.1 | WV Stream Condition Index (WVSCI) | 0-100 0-1 |
| | | | | | | | 2.20 0.1 | | | | | |
| PART II - Index and Unit Score | Sub-Total | 0.707 | Sub-Total | 0 | | Sub-Total | | 0 | Sub-Total | 0 | Sub-Total | 0 |
| | PART II - Index and U | Jnit Score | PART II - Index and | Unit Score | | PART II - Index and | I Unit Score | | PART II - Index and L | Init Score | PART II - Index and | Unit Score |
| Index Linear Feet Unit Score Index Linear Feet Unit Score Index Linear Feet Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet Unit Score | | Index | Linear Feel | Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet Unit Score |
| 0.804 103 82.812 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | 0.804 | 103 82.812 | 0 | 0 0 | | 0 | 0 | 0 | 0 | 0 0 | 0 | 0 0 |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)



PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential Other Indicate the dominant type and record the domin Indicate the dominant type and record the domin Trees Dominant species present Sycamore, American beech, rose, jewely | ✓ Grasses ✓ Herbaceous |
|--|---|--|
| INSTREAM FEATURES | Estimated Reach Length 57.9 m Estimated Stream Width 0.6 m Sampling Reach Area m² Area in km² (m²x1000) km² Estimated Stream Depth 0.1 m Surface Velocity (at thalweg) 1.2 m/sec | Canopy Cover □ Partly shaded □ Shaded □ Partly open □ Partly shaded □ Shaded High Water Mark 0.2 m Proportion of Reach Represented by Stream Morphology Types Riffle ±0 % Pool % Channelized Yes Dam Present Yes |
| LARGE WOODY DEBRIS | LWDm ² Density of LWDm ² /km ² (LWD/ reac | ch area) |
| AQUATIC VEGETATION | Indicate the dominant type and record the domin Rooted emergent Floating Algae Dominant species present None Portion of the reach with aquatic vegetation | ☐Rooted floating ☐Free floating |
| WATER QUALITY | Temperature 19.5 0 C Specific Conductance 67.1 D ms/cm Dissolved Oxygen 6.95 D mg/L pH 6.5 D su Turbidity N/A WQ Instrument Used YSI | Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Slick Sheen None Other Turbidity (if not measured) Turbid Clear Slightly turbid Opaque Stained |
| SEDIMENT/ SUBSTRATE | Odors Normal Chemical Other Oils Absent Slight | Deposits □Sludge □Sawdust □Paper fiber □Sand □Relict shells □Other □Lpoking at stones which are not deeply embedded, are the undersides black in color? □Yes ☑No |

| INC | ORGANIC SUBSTRATE (should add up to | | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | | | | | |
|-------------------|--|----|-------------------|---|-----------------------------------|--|--|--|--|--|--|
| Substrate Type | | | Substrate Type | Characteristic | % Composition in Sampling Area | | | | | | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | F | | | | | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | 5 | | | | | | |
| Cobble | 64-256 mm (2.5"-10") | 40 | Muck-Mud | black, very fine organic (FPOM) | 10 | | | | | | |
| Gravel | 2-64 mm (0.1"-2.5") | 30 | | (FPOM) | 10 | | | | | | |
| Sand | 0.06-2mm (gritty) | 10 | Marl | grey, shell fragments | | | | | | | |
| Silt | 0.004-0.06 mm | 20 |] | | | | | | | | |
| Clay | < 0.004 mm (slick) | | | | | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME S-GH44 | LOCATION Franklin County | | | | |
|---|---|--|--|--|--|
| STATION # RIVERMILE | STREAM CLASS Perennial | | | | |
| LAT <u>37.028392</u> LONG <u>-79.773359</u> | RIVER BASIN Upper Roanoke | | | | |
| STORET # | AGENCY VADEQ | | | | |
| INVESTIGATORS AJ, VM | | | | | |
| FORM COMPLETED BY AJ, VM | DATE 8-26-21 TIME 930 AM PM REASON FOR SURVEY Baseline Assessment | | | | |

| | Habitat | | Condition | a Category | | | |
|--|---|---|---|---|---|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | |
| | _{score} 11 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| ı sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | |
| ted ir | score 9 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | | |
| Iram | score 11 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | |
| | _{SCORE} 16 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | | |
| | _{score} 13 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| | Habitat | | Condition | n Category | | | |
|--|--|--|--|--|---|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | |
| | 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | |
| | _{score} 16 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| ling reach | 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | |
| amp | score 15 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | |
| Parameters to be evaluated broader than sampling reach | 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | |
| e ev | SCORE 9 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |
| top | SCORE 9 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |
| Parameters | 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | |
| | SCORE 7 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |
| | SCORE 7 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |
| | 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. | | |
| | SCORE 9 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |
| | SCORE 9 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | |

Total Score 141

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAMES-GH44 | | LOCATION Franklin County | , | | | | | | |
|----------------------|--|---|--|--|--|--|--|--|--|
| STATION # | RIVERMILE 258.5 | STREAM CLASS Perennial | STREAM CLASS Perennial | | | | | | |
| LAT | LONG79.773359 | RIVER BASIN Upper Roano | ke | | | | | | |
| STORET # | | AGENCY VADEQ | | | | | | | |
| INVESTIGATORS A | J, VM | | LOT NUMBER | | | | | | |
| FORM COMPLETED | ^{BY} AJ | DATE <u>9/09/21</u> TIME <u>8:30</u> | REASON FOR SURVEY Baseline Assessmnet | | | | | | |
| | | | | | | | | | |
| HABITAT TYPES | S Indicate the percentage of each habitat type present □ Cobble% □Snags% □Vegetated Banks% □Sand% □ Submerged Macrophytes% □Other ()% | | | | | | | | |
| SAMPLE COLLECTION | Gear used D-frame | | rom bank 🗌 from boat | | | | | | |
| | Indicate the number of jabs/kicks taken in each habitat type. ✓ Cobble 4 □Snags □Vegetated Banks □Sand □ Submerged Macrophytes □Other () | | | | | | | | |
| GENERAL COMMENTS | Yes, benthics sampled 9/9/2021. | | | | | | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

Mountain Valley Pipeline Data are not adjusted for subsampling

ECO ANALYSTS, INC.

| | Sample ID Collection Date | S-GH44 09-09-2021 |
|----------------------|------------------------------------|----------------------|
| | | |
| ORDER | GENUS/SPECIES | COUNT |
| Ephemeroptera | | 1 |
| Ephemeroptera | | 2 |
| Ephemeroptera | | 1 |
| | Eccoptura xanthenes | 5 5 |
| Plecoptera | | 5 |
| | Cheumatopsyche sp. Chimarra sp. | 2 |
| | Diplectrona sp. | 11 |
| Trichoptera | | 1 |
| | Hydropsyche sp. | 4 |
| | Rhyacophila sp. | 6 |
| | Anchytarsus bicolor | 1 |
| Coleoptera | Ectopria sp. | 11 |
| Coleoptera | Helichus sp. | 2 |
| Coleoptera | Optioservus sp. | 1 |
| Coleoptera | Oulimnius sp. | 1 |
| Coleoptera | Psephenus sp. | 95 |
| Coleoptera | Stenelmis sp. | 1 |
| Diptera-Chironomidae | Micropsectra sp. | 8 |
| Diptera-Chironomidae | | 1 |
| Diptera-Chironomidae | | 5 |
| Diptera-Chironomidae | | 3 |
| Diptera-Chironomidae | | 1 |
| • | Thienemannimyia gr. sp. | 2 |
| Diptera | Antocha sp. | 2 |
| Diptera | Ceratopogoninae | 3 |
| Diptera | Dixa sp. | 1 |
| Diptera | Simulium sp. | 1 |
| Diptera | Tabanidae | 2 |
| Annelida | tubificoid Naididae w/ cap setae | 2 |
| Bivalvia | Pisidium sp. | 1 |
| Gastropoda | Elimia sp. | 18 |
| Gastropoda | Ferrissia sp. | 2 |
| Gastropoda | Lymnaeidae | 2 |
| Other Organisms | - | 2 |
| | TOTAL | 209 |

Mountain Valley Pipeline WV SCI Metrics



| Sample ID Collection Date | |
|---|---|
| WVSCI Metric Values Total taxa EPT taxa % EPT % Chironomidae % 2 Dominant HBI | 25 9 19.6 9.6 60.3 4.31 |
| WVSCI Metric Scores Total taxa EPT taxa % EPT % Chironomidae % 2 Dominant HBI | 119.0 69.2 21.3 91.3 62.1 80.1 |
| WVSCI Metric Scores Total taxa EPT taxa % EPT % Chironomidae % 2 Dominant HBI | 100.0 69.2 21.3 91.3 62.1 80.1 |
| WVSCI Total Score | 70.7 |

WVSCI Thresholds

Unimpaired = > 68.00 Gray Zone = 60.61 to 68.00 Impaired = <60.61

WOLMAN PEBBLE COUNT FORM

Basin:

County:Franklin CountyStream Name:UNT to Foul Ground CreekHUC Code:03010101Survey Date:8/26/2021Surveyors:AJ,VMType:Representative

Stream ID: S-GH44

Upper Roanoke

| | | | LE COUNT | · · · | | | |
|-------------|-------------|-------------|-------------|-------------------|---------|--------|--------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cun |
| | Silt/Clay | < .062 | S/C | * * | 20 | 20.00 | 20.00 |
| | Very Fine | .062125 | | ÷ | | 0.00 | 20.00 |
| | Fine | .12525 | | • | | 0.00 | 20.00 |
| | Medium | .255 | SAND | ÷ | 7 | 7.00 | 27.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | | 0.00 | 27.00 |
| .0408 | Very Coarse | 1.0-2 | | * * | 3 | 3.00 | 30.00 |
| .0816 | Very Fine | 2 -4 | | - | | 0.00 | 30.00 |
| .1622 | Fine | 4 -5.7 | | * * | | 0.00 | 30.00 |
| .2231 | Fine | 5.7 - 8 | | * * | | 0.00 | 30.00 |
| .3144 | Medium | 8 -11.3 | | - | 8 | 8.00 | 38.00 |
| .4463 | Medium | 11.3 - 16 | G R A V E L | * * | 5 | 5.00 | 43.00 |
| .6389 | Coarse | 16 -22.6 | | * * | 12 | 12.00 | 55.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | * * | | 0.00 | 55.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | | 5 | 5.00 | 60.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | * * | | 0.00 | 60.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | * * | 5 | 5.00 | 65.00 |
| 3.5 - 5.0 | Small | 90 - 128 | COBBLE | | 25 | 25.00 | 90.00 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | ÷ | 7 | 7.00 | 97.00 |
| 7.1 - 10.1 | Large | 180 - 256 | | * * | 3 | 3.00 | 100.00 |
| 10.1 - 14.3 | Small | 256 - 362 | | * * | | 0.00 | 100.00 |
| 14.3 - 20 | Small | 362 - 512 | | * * | | 0.00 | 100.00 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | * * | | 0.00 | 100.00 |
| 40 - 80 | Large | 1024 -2048 | | ÷ | | 0.00 | 100.00 |
| 80 - 160 | Vry Large | 2048 -4096 |] | * * | | 0.00 | 100.00 |
| | Bedrock | | BDRK | * * | | 0.00 | 100.00 |
| | | | | Totals: | 100 | | |


| River Name: U Reach Name: S Sample Name: R Survey Date: 0 | Representative | | | | | | |
|--|---|--------|--|--|--|--|--|
| Size (mm) | тот # | ITEM % | CUM % | | | | |
| | 20 | | 20.00 20.00 27.00 27.00 30.00 30.00 30.00 30.00 30.00 35.00 55.00 60.00 60.00 65.00 90.00 97.00 100.00 100.00 100.00 100.00 | | | | |
| D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Gravel (%) Boulder (%) Bedrock (%) | $\begin{array}{c} 0.05 \\ 10.06 \\ 19.85 \\ 118.88 \\ 165.14 \\ 256 \\ 20 \\ 10 \\ 30 \\ 40 \\ 0 \\ 0 \\ \end{array}$ | | | | | | |

Total Particles = 100.

| | S | Strean | | | | | | | | |
|--|---|---|--|--|--|--|--|--|---|------------|
| | | | | tream Method able channels cla | | • | | | | |
| Ducie of # | Droiget Norme (Ann | | | Cowardin | | | | Impact | Impact | |
| Project # | Project Name (App | , | Locality | Class. | HUC | Date | SAR # | Length | Factor | |
| 22865.06 | Mountain Valley Pipeline Valley Pipeline, I | | Franklin | R3 | 03010101 | 8/26/2021 | S-GH44 | 103 | 1 | |
| Nam | e(s) of Evaluator(s) | | County e and Inform | ation | | | | SAR Length | | |
| | AJ, VM | S-GH44 | | | | | | 190 | | |
| | · | | | | | | | 150 | | |
| Channel C | Condition: Assess the cross-sec | tion of the stream | | | | | | | | |
| | Optimal | Subo | ptimal | Conditional Catego | ginal | Po | or | Sev | vere | |
| Channel Condition | Very little incision or active erosion; 80 100% stable banks. Vegetative surface protection or natural rock, prominent (80-100%). AND/OR Stable bankfull benches are present. Access to their original floodpilain or fully developed wide bankfull benches. Mid channel bars and transverse bars few. Transient sediment deposition covers less than 10% of bottom. | erosion or unprotec of banks are s Vegetative protec prominent (60 Depositional feat stability. The bar channels are wel likely has acc benches,or ne | ew areas of active ted banks. Majority table (60-80%). tion or natural rock -80%) AND/OR ures contribute to htfull and low flow I defined. Stream ess to bankfull wby developed each. Transient | Poor. Banks more or Poor due to lo Erosion may be pro- both banks. Vege 40-60% of banks. be vertical or un 40-60% Sediment transient, contr | ibute instability. ntribute to stability, | laterally unstable further. Majority near vertical. Eros banks. Vegetative on 20-40% of bank to prevent erosion the stream is cov Sediment is temp nature, and contri | cised. Vertically / e. Likely to widen of both banks are sion present on 60- protection present , AND/OR 60-80% ered by sediment. orary / transient in buting to instability. Ved channels have | present. Erosion/ 100%. AND/OR A | stability. Severe tained within the ed below average vertical/undercut. on present on less s, is not preventing s bank sloughing 'raw banks on 80- | |
| | | sediment covers stream | s 10-40% of the bottom. | shaped channels protection on > 40 depositional featur to sta | s have vegetative % of the banks and es which contribute ability. | vegetative protect 40% of the banks a deposition | ion is present on > and stable sediment n is absent. | deposition, contrib Multiple thread o subterran | uting to instability. channels and/or ean flow. | CI |
| Scores | 3 | 2 | .4 | | 2 | 1 | .6 | 1 | 1 | 2.40 |
| RIPARIAN | N BUFFERS: Assess both bank | | | | gh measurements | of length & width | may be acceptab | | | |
| . RIPARIAN | N BUFFERS: Assess both bank | Con Subo High Suboptimal: Riparian areas | n areas along the ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum | gory | ginal Low Marginal: Non-maintained, dense herbaceous | Pc High Poor: Lawns, mowed, and maintained areas, | Low Poor: | ie) NOTES>> | | |
| . RIPARIAN Riparian Buffers | 1 | Con Subo High Suboptimal: | ditional Cate ptimal Low Suboptimal: Riparian areas | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub | ginal Low Marginal: Non-maintained, | Pc High Poor: Lawns, mowed, and | por | | | |
| Riparian | Optimal Tree stratum (dbh > 3 inches) present, with > 60% tree canopy cover. Wetlands located within the riparian areas. | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understoy Low | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. | Dor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low | | | |
| Riparian | Optimal Tree stratum (dbh > 3 inches) present, with > 60% tree canopy cover. Wetlands located within the riparian | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | | | |
| Riparian Buffers Scores Delineate ripa Determine sq low. | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian | NOTES>> | | |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 | Dor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 | NOTES>> | is limited to | |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F | Optimal Tree stratum (dbh > 3 inches) present, with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 | Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian | NOTES>> Assessment areas wi | ithin the | |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 | Dor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 | NOTES>> | ithin the ry ROW. | |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F Right Bank | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 | Dor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 | NOTES>> Assessment areas wi tempora | ithin the ry ROW. | CI |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F Light Bank | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> 100% Score > 0.85 | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- ogth and width. Ca | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F | Door Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 | NOTES>> Assessment areas wi tempora CI= (Sum % RA * Sc | thin the ry ROW. cores*0.01)/2 | |
| Riparian Buffers Scores Delineate ripe Determine sq ow. Enter the % F Right Bank Left Bank | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> 100% Score > 0.85 % Riparian Area> 100% | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) ro 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge arian category in | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Com gth and width. Cat the blocks below. | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using alculators are prov | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. ided for you | High Poor: Lawns, mowed, and maintained areas, nurseries, no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks e | Dor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100% | NOTES>> Assessment areas wi tempora CI= (Sum % RA * So Rt Bank CI > Lt Bank CI > | thin the ry ROW. pores*0.01)/2 0.85 0.85 | CI 0.85 |
| Riparian Buffers Scores Delineate ripa Determine sq ow. Enter the % F Right Bank Left Bank INSTREAI | Optimal Tree stratum (dbh > 3 inches) present, with > 60% tree canopy cover. Wetlands located within the riparian areas. 1.5 Train areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> 100% Score > 0.85 % Riparian Area> 100% Score > 0.85 % Riparian Area> 100% Score > 0.85 % HABITAT: Varied substrate sizexes, stable features. | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating leng arian category in | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- gth and width. Ca the blocks below. | High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using alculators are prov | ginal Low Marginal: Non-maintained, dense herbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. ided for you s; stable substrat | Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure of % F Blocks e | Door Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lost, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100% 100% | NOTES>> Assessment areas wi tempora CI= (Sum % RA * So Rt Bank CI > Lt Bank CI > | thin the ry ROW. pores*0.01)/2 0.85 0.85 | |
| Riparian Buffers Scores Delineate ripa Determine sq low. Enter the % F Right Bank Left Bank INSTREAI InSTREAI Habitat/ | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring Riparian Area and Score for each rip % Riparian Area> 100% Score > 0.85 % Riparian Area> 100% Score > 0.85 % Riparian Area> 100% Score > 0.85 % HABITAT: Varied substrate size | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge arian category in carian category in category | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Con- gth and width. Ca the blocks below. | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using alculators are prov | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. ided for you | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F Blocks e blocks e lacking or are u elements are typic | Door Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100% 100% Setting a state of the sums restable. Habitat ally present in less | NOTES>> Assessment areas wi tempora CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > Lt Bank CI > | thin the ry ROW. pores*0.01)/2 0.85 0.85 | |
| Riparian Buffers Scores Delineate ripa Determine sq low. Enter the % F Right Bank Left Bank INSTREAI Instream Habitat/ Available | Optimal Tree stratum (dbh > 3 inches) present, with > 60% free canopy cover. Wetlands located within the riparian areas. 1.5 arian areas along each stream bank uare footage for each by measuring % Riparian Area and Score for each rip % Riparian Area> 100% Score > 0.85 MHABITAT: Varied substrate sizes, stable features. Optimal Habitat elements are typically present | Con Subo High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 into Condition Ca or estimating lenge arian category in carian category in category | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 tegories and Cono gth and width. Cat the blocks below. | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 dition Scores using alculators are prov | ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory Low 0.75 g the descriptors. ided for you | High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F Blocks e blocks e lacking or are u elements are typic | Cor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 the sums Riparian equal 100 100% 100% Series; shade; under conditions stable. Habitat | NOTES>> Assessment areas wi tempora CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > Lt Bank CI > | ithin the ry ROW. cores*0.01)/2 0.85 0.85 ts; SAV; | |

Reach R3-R4 File: C:\Users\dan.weidenhof\Documents\Documents\VA Stream Sampling\0 QAQC SUBMITTALS\QAQC working 1st submittal\Ready for Submittal\Needs Benthics\S-GH44_20210913KEH benthics & LP\9. S-GH44_USM_MVP_20210913KEH.xlsx


| | St | ream In | npact A | ssessn | nent Fo | rm Pag | e 2 | | | |
|---|--|--|---------------------|--------------------|---|--|---|-------------------------|-------------------|------------|
| Project # | Project Name (App | licant) | Locality | Cowardin Class. | HUC | Date | SAR # / Data Point | Impact / SAR length | Impact Factor | |
| 22865.06 | | Mountain Valley Pipeline (Mountain Franklin Valley Pipeline, LLC) County | | R3 | 03010101 | 44434 | S-GH44 | 103 | 1 | |
| I. CHANNE | LALTERATION: Stream cross | ings, riprap, concr | rete, gabions, or c | oncrete blocks, st | raightening of cha | annel, channelizat | ion, embankment | s, spoil piles, constri | ctions, livestock | |
| | | | | al Category | | | | NOTES>> | | |
| | Negligible | Mi | nor | | erate | Se | vere | | | |
| Channel Alteration | Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized. | Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines. | the channel | of the channel | is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered | by any of the chan in the parameter of 80% of banks sh | of reach is disrupted nel alterations listed juidelines AND/OR nored with gabion, r cement. | | | CI |
| Scores | 1.5 | 1.5 1.3 1.1 0.9 0.7 | | 0 | .5 | | | 1.30 | | |
| | REACH C | ONDITION I | INDEX and S | TREAM CO | NDITION UN | | IIS REACH | | | |
| NOTE: The Cis and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number. THE REACH CONDITION INDEX (RCI) >> | | | | | | | | 1.15 | | |
| | | | | | | RCI= (Sum of | all CI's)/5, exce | pt if stream is epl | nemeral RCI = | Riparian C |
| COMPENSATION REQUIREMENT (CR) >> | | | | | | | | | 118 | |
| CR = RCI X L _I X IF | | | | | | | | | | |



DESCRIBE PROPOSED IMPACT:

PROVIDED UNDER SEPARATE COVER

Reach R3-R4 File: C:\Users\dan.weidenhof\Documents\Documents\VA Stream Sampling\0 QAQC SUBMITTALS\QAQC working 1st submittal\Ready for Submittal\Needs Benthics\S-GH44_20210913KEH benthics & LP\9. S-GH44_USM_MVP_20210913KEH.xlsx



| S-GH44 CROSS SECTION B (PIPE CL) | | | | | | | | | |
|----------------------------------|---------|---------------|-------|--|--|--|--|--|--|
| CROSSING | | POST-CROSSING | | | | | | | |
| | | VERT. | HORZ. | | | | | | |
| ASTING | ELEV | DIFF. | DIFF. | | | | | | |
| 98346.45 | 1015.46 | | | | | | | | |
| 98349.79 | 1013.92 | | | | | | | | |
| 98355.15 | 1014.20 | | | | | | | | |
| 98358.49 | 1014.29 | | | | | | | | |
| 98360.58 | 1015.82 | | | | | | | | |
| | | | | | | | | | |

