Reach S-GH6 (Timber Mat Crossing) Perennial Spread I Franklin County, Virginia

| Data | Included | | | | | |
|---|---|--|--|--|--|--|
| Photos | \checkmark | | | | | |
| SWVM Form | \checkmark | | | | | |
| FCI Calculator and HGM Form | Perennial stream (not shadeable, slope >4%) | | | | | |
| RBP Physical Characteristics Form | \checkmark | | | | | |
| Water Quality Data | \checkmark | | | | | |
| RBP Habitat Form | ✓ | | | | | |
| RBP Benthic Form | \checkmark | | | | | |
| Benthic Identification Sheet | N/A – Lack of habitat | | | | | |
| Wolman Pebble Count | \checkmark | | | | | |
| RiverMorph Data Sheet | \checkmark | | | | | |
| USM Form (Virginia Only) | \checkmark | | | | | |
| Longitudinal Profile and Cross Sections | \checkmark | | | | | |

Stream S-GH6 (TMC) Franklin County



Photo Type: DS VIEW Location, Orientation, Photographer Initials: Downstream view of ROW looking S, KB



Photo Type: US VIEW Location, Orientation, Photographer Initials: Upstream view of ROW looking N, KB

Stream S-GH6 (TMC)



Photo Type: LB CL Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking W, KB



Photo Type: RB CL Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking E, KB

DEQ Permit #21-0416

Stream S-GH6 (TMC) Franklin County



Photo Type: DS COND Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking S, KB

L:\22000s\22860\22865.06\Admin\05-ENVR\Field Data\Spread I\Field Forms\S-GH6\0_Potesta Submission\Docs\Photos_S-GH6.docx

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | Mountain | Valley Pipeline | IMPACT COORDINATES: (in Decimal Degrees) | Lat. | 37.092397 | .on. | -79.983227 | WEATHER: | Sunny, Partly Cloudy | DATE: | August 2 | 27, 2021 |
|--|-------------------------------|--|--|------|--|--|---------------------------|---|-------------------------------|---|---------------------|-----------------------|
| IMPACT STREAM/SITE ID (watershed size (acreage), s | | S-0 | GH6 | | MITIGATION STREAM CLASS./SI (watershed size (acreage), u | | | | | Comments: | | |
| STREAM IMPACT LENGTH: | 20 FORM OF MITIGATION: | RESTORATION (Levels I-III) | MIT COORDINATES: (in Decimal Degrees) | Lat. | | .on. | | PRECIPITATION PAST 48 HRS: | 0 | Mitigation Length: | | |
| Column No. 1- Impact Existing | Condition (Debit) | Column No. 2- Mitigation Existing Co | ondition - Baseline (Credit) | | Column No. 3- Mitigation Proje Post Completion (6 | | e Years | Column No. 4- Mitigation Proje Post Completion (| cted at Ten Years Credit) | Column No. 5- Mitigation Project | ed at Maturity (Cr | redit) |
| Stream Classification: | Perennial | Stream Classification: | Perennial | | Stream Classification: | | Perennial | Stream Classification: | Perennial | Stream Classification: | Peren | nnial |
| Percent Stream Channel Sic | ppe 1.7 | Percent Stream Channel Slo | ope | | Percent Stream Channel Slop | e | 0 | Percent Stream Channel SI | ope 0 | Percent Stream Channel S | lope | 0 |
| HGM Score (attach da | ita forms): | HGM Score (attach o | data forms): | | HGM Score (attach da | ita forms) | | HGM Score (attach da | ata forms): | HGM Score (attach d | ata forms): | |
| | Average | | Average | | | | Average | | Average | | | Average |
| Hydrology Biogeochemical Cycling Habitat | 0 | Hydrology Biogeochemical Cycling Habitat | 0 | | Hydrology Biogeochemical Cycling Habitat | | 0 | Hydrology Biogeochemical Cycling Habitat | 0 | Hydrology Biogeochemical Cycling Habitat | | 0 |
| PART I - Physical, Chemical and I | Biological Indicators | PART I - Physical, Chemical and | d Biological Indicators | | PART I - Physical, Chemical and | Biological | Indicators | PART I - Physical, Chemical and | Biological Indicators | PART I - Physical, Chemical and | Biological Indica | ators |
| | Points Scale Range Sile Score | | Points Scale Range Site Score | | 1 | Points Scale Ran | ge Site Score | | Points Scale Range Site Score | | Points Scale Range | Site Score |
| PHYSICAL INDICATOR (Applies to all streams | classifications) | PHYSICAL INDICATOR (Applies to all streams of | classifications) | | PHYSICAL INDICATOR (Applies to all streams cla | assifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | PHYSICAL INDICATOR (Applies to all streams | s classifications) | - |
| USEPA KRB (High Gradient Data Sheet) L Enfland Schutz Available Cover 2. Entback Schutz Available Cover 3. Entback Depath Regime 5. Schurzel Frey States 5. Charavel F | | USEFA RAP (Low Gradient Data Sheet) I - Enflavant Substrate Available Cover 2 Pool Substrate Characterization 4 - Sediment Deposition 5 - Charanel Flow Status 5 - Charanel Available 5 - Charan | 0.20 0.21 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 0.20 0.1 | | 2. Embeddedness 3. Velocity/ Depth Regime 4. Sediment Deposition 5. Channel How Status 6. Channel Alkraration 7. Frequency of Riffles (or bends) 8. Bank Stability (LB & RB) 9. Vegetative Protection (LB & RB) | 0-20 0-20 0-20 0-20 0-20 0-20 0-20 0-20 | 1 0 0 0 53teans) | USEPA RBP (tright Gradient Data Sheet) 1. Epflurant Substration/Validate Cover 2. Embeddedness 3. Velocity (Opth Regime 4. Sediment Deposition 5. Charnel Attention 1. Englurant Pro Status 6. Charnel Attention 1. Englurant of Refiles (or kands) 3. Vench Balter Pr (Labor) 3. Vench Balter Pr (Labor) 10. Repart and Vegetable Zover Wolft, (LB & RE) 10. Repart of Vegetable Zover Wolft, (LB & RE) 10. | | USEPAREP (High Gradient Data Sheet) 1. Epiferunal Substrate/Available.Cover 2. Enheaddedness 3. Velocity/ Depit Regime 4. Sediment Deposition 5. Channel Free Status 6. Channel Free Status 6. Channel Free Status 6. Channel Free Status 6. Channel Free Status 10. Tresumor of Riffles (or bords) 1. Tresumor of Riffles (or bords) 10. Research vegetates Zone Worth (LB & RE) 1. Catal REP Score Sub-Total ChEMICAL INDICATOR (Applies to Intermitted Specific Conductivity pH DO | | 0 0 0 teams) |
| Sub-Total BIOLOGICAL INDICATOR(Applies to Intermitte | 0.825 | Sub-Total BIOLOGICAL INDICATOR (Applies to Intermitte | 0 | | Sub-Total BIOLOGICAL INDICATOR (Applies to Intermitte | ent and Pere | 0 nnial Streams) | Sub-Total BIOLOGICAL INDICATOR (Applies to Interm | 0 | Sub-Total BIOLOGICAL INDICATOR (Applies to Interm | ittent and Perennia | 0 |
| WV Stream Condition Index (WVSCI) | , | WV Stream Condition Index (WVSCI) | , | | WV Stream Condition Index (WVSCI) | | , | WV Stream Condition Index (WVSCI) | , | WV Stream Condition Index (WVSCI) | | , |
| 0 Sub-Total | 0-100 0-1 0 | Sub-Total | 0-100 0-1 0 | | Sub-Total | 0-100 0- | 1 0 | Sub-Total | 0-100 0-1 O | Sub-Total | 0-100 0-1 | 0 |
| PART II - Index and U | nit Score | PART II - Index and I | Unit Score | | PART II - Index and U | nit Score | | PART II - Index and U | nit Score | PART II - Index and U | Init Score | |
| Index | Linear Feet Unit Score | Index | Linear Feet Unit Score | | Index | Linear Fee | t Unit Score | Index | Linear Feet Unit Score | Index | Linear Feet | Unit Score |
| 0.685 | 20 13.7 | 0 | 0 0 | | 0 | 0 | 0 | 0 | 0 0 | 0 | 0 | 0 |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME S-GH6 | | LOCATION Franklin County | | | | | | |
|-------------------------|------------------------------|---|---|--|--|--|--|--|
| STATION # R | IVERMILE | STREAM CLASS Perennial | | | | | | |
| LAT <u>37.092397</u> LO | ONG79.983227 | RIVER BASIN Upper Roand | bke | | | | | |
| STORET # | | AGENCY VADEQ | | | | | | |
| INVESTIGATORS JB, AW | / | - | | | | | | |
| FORM COMPLETED BY | JB, AW | DATE 8/27/2021 TIME 1:30 PM | REASON FOR SURVEY Baseline Assessment | | | | | |
| | | 1 | Has there been a heavy rain in the last 7 days? | | | | | |
| WEATHER CONDITIONS | Now | Past 24 hours | Yes \checkmark No | | | | | |
| | 10 % $10 %$ $10 %$ $10 %$ | (steady rain) | Air Temperature <u>33.3</u> ⁰ C Dther | | | | | |
| SITE LOCATION/MAP | Draw a map of the sit | te and indicate the areas sample | d (or attach a photograph) | | | | | |
| STREAM | CO CO Stream Subsystem | S ROCK PILLE HERBAU SIGH HERBAU HERBAU NOWED (| EOUSVEZ PRIAN TREA TASS JEOINE TREATINE | | | | | |
| CHARACTERIZATION | Stream Subsystem | ermittent Tidal [Spring-fed | Coldwater ØWarmwater | | | | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse ✓ Forest Commercial ✓ Field/Pasture Industrial ✓ Agricultural ✓ Other readway ■ Residential ✓ Other state Indicate the dominant type and record the domin □ Trees □ Trees □ Shrubs □ Dominant species present Sorghastrum nutans, Lespedeza cuneata | |
|--|---|---|
| INSTREAM FEATURES | Estimated Reach Length 11.6 m Estimated Stream Width 0.9 m Sampling Reach Area 10.4 m² Area in km² (m²x1000) km² Estimated Stream Depth 0.3 m Surface Velocity (at thalweg) 0.1 m/sec | Canopy Cover □Partly shaded □Shaded I Partly open □Partly shaded □Shaded High Water Mark 0.5 m Proportion of Reach Represented by Stream Morphology Types Riffle 20 % Pool 40 % Channelized Yes Dam Present Yes |
| LARGE WOODY DEBRIS | LWD <u>•</u> m ² Density of LWD <u>•</u> m ² /km ² (LWD/ reac | ch area) |
| AQUATIC VEGETATION | Indicate the dominant type and record the domin Rooted emergent Floating Algae Dominant species present Portion of the reach with aquatic vegetation | ☐Rooted floating ☐Free floating |
| WATER QUALITY (DS, US) | Temperature 20.1, 20 0 C Specific Conductance 84.6, 84.1 Dissolved Oxygen 8.68, 9.38 pH 8.75, 8.53 Turbidity 56.25, 54.60 WQ Instrument Used YSI | Water Odors Normal/None Sewage Petroleum Chemical Fishy Other Slick Sheen None Other Turbidity (if not measured) Turbid Clear Slightly turbid Opaque Stained |
| SEDIMENT/ SUBSTRATE | Odors Sewage Petroleum Other Anaerobic None Oils Absent Slight Moderate Profuse | Deposits □Sludge □Sawdust □Paper fiber □Sand □Relict shells □Other □Lpoking at stones which are not deeply embedded, are the undersides black in color? □Yes ☑No |

| INC | ORGANIC SUBSTRATE (should add up to | | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | | | | |
|-------------------|--|------------------------------------|-------------------|---|-----------------------------------|--|--|--|--|--|
| Substrate Type | Diameter | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | | | | | |
| Bedrock | | 0 | Detritus | sticks, wood, coarse plant | F | | | | | |
| Boulder | > 256 mm (10") | 0 | | materials (CPOM) | 5 | | | | | |
| Cobble | 64-256 mm (2.5"-10") | 25 | Muck-Mud | black, very fine organic | 0 | | | | | |
| Gravel | 2-64 mm (0.1"-2.5") | 20 | | (FPOM) | 0 | | | | | |
| Sand | 0.06-2mm (gritty) | 25 | Marl | grey, shell fragments | 0 | | | | | |
| Silt | 0.004-0.06 mm | 30 |] | | 0 | | | | | |
| Clay | < 0.004 mm (slick) | 0 |] | | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME S-GH6 | LOCATION Franklin County | | | | |
|---|---|--|--|--|--|
| STATION # RIVERMILE | STREAM CLASS Perennial | | | | |
| LAT <u>37.092397</u> LONG <u>-79.983227</u> | RIVER BASIN Upper Roanoke | | | | |
| STORET # | AGENCY VADEQ | | | | |
| INVESTIGATORS JB, AW | | | | | |
| FORM COMPLETED BY JB, AW | DATE 8/27/2021 REASON FOR SURVEY TIME 1:30 PM AM PM Baseline Assessment | | | | |

| | Habitat | | Condition | Category | | | | | |
|--|---|---|---|---|---|--|--|--|--|
| | Parameter | Optimal | Suboptimal | Marginal | Poor | | | | |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. | | | | |
| | _{SCORE} 8 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. | | | | |
| ted i | _{score} 6 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). | | | | |
| ıram | _{SCORE} 8 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. | | | | |
| | _{score} 6 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. | | | | |
| | SCORE 17 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |

Notes:

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Conditio | on Category | | | | | |
|--|--|---|--|--|--|--|--|--|
| Parameter | Optimal | Suboptimal | Marginal | Poor | | | | |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabio or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. | | | | |
| _{SCORE} 19 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water of shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. | | | | |
| score 3 | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 | | | | |
| 8. Bank Stability (score each bank) Note: determine left or right side by facing deurstream. | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. | | | | |
| SCORE 8 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE 9 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potentia to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streamban vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. | | | | |
| SCORE 4 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE 4 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone < meters: little or no riparian vegetation due t human activities. | | | | |
| SCORE 9 | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |
| SCORE 8 | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 | | | | |

Total Score 109 Notes:

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME S-G | GH6 | LOCATION Franklin County | LOCATION Franklin County | | | | | | |
|----------------------|--|--------------------------------|--|--|--|--|--|--|--|
| STATION # | RIVERMILE | STREAM CLASS Perennial | | | | | | | |
| LAT <u>37.092397</u> | LONG79.983227 | RIVER BASIN Upper Roand | ke | | | | | | |
| STORET # | | AGENCY VADEQ | | | | | | | |
| INVESTIGATORS JE | 3, AW | | LOT NUMBER | | | | | | |
| FORM COMPLETED | JB, AW | DATE 8/27/2021 TIME 1:30 PM | REASON FOR SURVEY Baseline Assessment | | | | | | |
| | | | | | | | | | |
| HABITAT TYPES | Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()% | | | | | | | | |
| SAMPLE COLLECTION | Gear used D-frame | kick-net Other_ | | | | | | | |
| | How were the samples coll | lected? wading fi | rom bank 🔲 from boat | | | | | | |
| | Indicate the number of jabs/kicks taken in each habitat type. CobbleSnags Vegetated Banks Sand Submerged Macrophytes Other () | | | | | | | | |
| GENERAL COMMENTS | No benthics collected due to lack of habitat. | | | | | | | | |
| | | | | | | | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

Basin:

County:Franklin CountyStream Name:UNT to Little CreekHUC Code:03010101Survey Date:8/27/2021Surveyors:JB AWType:Representative

Stream ID: S-GH6

Upper Roanoke

PEBBLE COUNT Inches PARTICLE Millimeters Particle Total # Item % % Cum Count Silt/Clay < .062 S/C ۸ 32 31.68 31.68 • Very Fine .062-.125 ۸ 0 0.00 31.68 • .125-.25 Fine ٠ 0.99 1 32.67 • Medium .25-.5 ۸ SAND 0 0.00 32.67 • .50-1.0 Coarse ۸ 6 5.94 38.61 • .04-.08 1.0-2 Very Coarse ۸ 12 11.88 50.50 • .08 -.16 Very Fine 2 -4 ٠ 0 0.00 50.50 • .16 - .22 Fine 4 - 5.7 ۸ 0 0.00 50.50 • .22 - .31 Fine 5.7 - 8 ۸ 1 0.99 51.49 • .31 - .44 Medium 8 - 11.3 ۸ 2 1.98 53.47 • .44 - .63 Medium 11.3 - 16 ۸ GRAVEL 2 1.98 55.45 • .63 - .89 Coarse 16 - 22.6 ۸ 2 1.98 57.43 • .89 - 1.26 Coarse 22.6 - 32 ۸ 2 1.98 59.41 -32 - 45 1.26 - 1.77 Vry Coarse ٠ 5 4.95 64.36 • 1.77 -2.5 Vry Coarse 45 - 64 ۸ 9 73.27 8.91 • 2.5 - 3.5 64 - 90 Small ۸ 11 10.89 84.16 • 3.5 - 5.0 Small 90 - 128 ۲ 12 11.88 96.04 • COBBLE 5.0 - 7.1 Large 128 - 180 ۸ 4 3.96 100.00 -7.1 - 10.1 Large 180 - 256 ۸ 0 0.00 100.00 • 10.1 - 14.3 Small 256 - 362 100.00 0 0.00 • 14.3 - 20 Small 362 - 512 ۸ 0 0.00 100.00 • 20 - 40 512 - 1024 Medium ۸ BOULDER 0 0.00 100.00 • 40 - 80 Large 1024 - 2048 ۸ 0 0.00 100.00 • 80 - 160 Vry Large 2048 - 4096 0 0.00 100.00 • BDRK . Bedrock 0 0.00 100.00 • Totals 101 Total Tally:


| River Name: Reach Name: Sample Name: Survey Date: | UNT to Little S-GH6 Representative 08/27/2021 | Creek | |
|---|--|--|--|
| Size (mm) | тот # | ITEM % | CUM % |
| 0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock | 32 0 1 0 6 12 0 0 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 | 31.68 0.00 0.99 0.00 5.94 11.88 0.00 0.00 0.99 1.98 1.089 11.88 3.96 0.00 | 31.68 31.68 32.67 32.67 38.61 50.50 50.50 51.49 53.47 55.45 57.43 59.41 64.36 73.27 84.16 96.04 100.00 100.00 100.00 100.00 100.00 100.00 100.00 |
| D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Gravel (%) Boulder (%) Bedrock (%) | 0.03 0.7 1.96 89.62 124.67 180 31.68 18.82 22.77 26.73 0 0 | | |

Total Particles = 101.


| | | | | Unified St | tream Method | lology for use | in Minalala | | | | |
|---|---|---|---|---|---|--|--|--|---|--|------|
| | | | | | | 0, | • | | | | |
| | | | | | Cowardin | | nittent or perennia | | Impact | Impact | |
| Project # | • | t Name (App | , | Locality | Class. | HUC | Date | SAR # | Length | Factor | |
| 22865.06 | Mountain Va Valle | y Pipeline, L | • | Franklin County | R3 | 03010101 | 8/27/2021 | S-GH6 | 20 | 1 | |
| | | | | Stream Name and Information | | | | | SAR Length | | |
| | AW, JB | | Unnamed Tri | butary to Litt | le Creek | | | | 75 | | |
| Channel C | ondition: Assess | s the cross-secti | on of the stream a | nd prevailing cond | dition (erosion, ag | gradation) | | | | | |
| | | | | | Conditional Catego | ory | • | | 1 | | |
| | Optimal | | Suboptimal | | Marginal | | Poor | | Severe | | |
| | Very little incision or active erosion; 80- | | | | Often incised, but less than Severe or | | Overwidened/incised. Vertically / | | Deeply incised | <u>s</u> | |
| Channel | 100% stable banks. N protection or natural | egetative surface | | | Poor. Banks more stable than Severe | | laterally unstable. Likely to widen further. Majority of both banks are near | | vertical/lateral in: incision, flow contain | stability. Severe | |
| Channel Condition | (80-100%). AND/OR | Stable point bars / | Vegetative protect | tion or natural rock | | | vertical. Erosion pr | esent on 60-80% of | Streambed below av | erage rooting depth, | |
| | bankfull benches are to their original flo | odplain or fully | Depositional feat | -80%) AND/OR ures contribute to | | | banks. Vegetative protection present on 20-40% of banks, and is insufficient to prevent erosion. AND/OR 60-80% of the stream is covered by sediment. | | majority of banks vertical/undercut. Vegetative protection present on less | | |
| | developed wide bank channel bars and tra | nsverse bars few. | | efined. Stream likely | | | | | than 20% of banks erosion. Obvious | s bank sloughing | |
| | Transient sediment of less than 10% | | newly developed | nkfull benches,or floodplains along | Deposition that co | | nature, and contri | | present. Erosion/raw AND/OR Aggradin | g channel. Greater | |
| | | | | portions of the reach. Transient sediment covers 10-40% of the stream | | may be forming/present. AND/OR V- shaped channels have vegetative | | AND/OR V-shaped channels have vegetative protection is present on > | | than 80% of stream bed is covered by deposition, contributing to instability. | |
| | | | bottom. | | protection on > 40% of the banks and depositional features which contribute | | 40% of the banks and stable sediment deposition is absent. | | t Multiple thread channels and/or subterranean flow. | | |
| | | | | | to stability. | | 1.6 | | | | CI |
| Scores | 3 | | 2 | .4 | | 2 | 1 | | 1 | | 2.40 |
| | BUFFERS: Ass | sess both bank's | | | | | | | NOTESSS | | |
| NOTES>> | I BUFFERS: Ass | | Con | areas along the er ditional Cate ptimal | gory | | f length & width ma | | NOTES>> | | |
| | 1 | nal 3 inches) present, anopy cover. ithin the riparian | Con | ditional Cate | gory | measurements o | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till | ay be acceptable) | NOTES>> | | |
| RIPARIAN Riparian Buffers | Optin Tree stratum (dbh > with > 60% tree c Wetlands located w area | nal 3 inches) present, canopy cover. tithin the riparian s. | Con Subop High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintened understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | measurements o ginal Low Marginal: Non-maintained, dense herbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. Low | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-til cropland; actively grazed pasture, sparsely vegetated area, recently seeded and stabilized, or other comparable condition. High | Ay be acceptable) DOF Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, conditions, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. | NOTES>> | | |
| RIPARIAN | Optin Tree stratum (dbh > with > 60% tree c Wetlands located w | nal 3 inches) present, canopy cover. tithin the riparian s. | Con Subop High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). | High Marginal: Non-maintained, dense herbaccous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrut and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh > 3 inches) present, with <30% tree canopy cover with maintained understory. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actived grazed pasture, sparsely vegetated anon-maintained anon-maintained area, recently seeded and stabilized, or other comparable condition. | Ay be acceptable) DOF Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, conditions, active feed lots, trails, or other comparable conditions. | NOTES>> | | |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 Trian areas along ear uare footage for eaco | nal 3 inches) present, canopy cover. ithin the riparian s. 5 ch stream bank h by measuring | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy comparison and a maincaver and a maincaver and a maincaver (dense vegetation). Low 1.1 egories and Cond th and width. Cal | High Marginal: Non-maintained, dense hetbaceous vegetation with either a shrub layer or a tree layer (dbh either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F | ay be acceptable) bor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian | NOTES>> | | |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 Tian areas along ear uare footage for eac uare footage for eac uare footage for eac | nal 3 inches) present, canopy cover. within the riparian s. 5 6 6 6 6 6 6 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 | Con Subor High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy course and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calc he blocks below. | High Marginal: Non-maintained, dense hetbaceous vegetation with either a shrub layer or a tree layer (dbh either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F | ay be acceptable) bor Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 | NOTES>> | | |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 Trian areas along ear uare footage for eaco | nal 3 inches) present, canopy cover. ithin the riparian s. 5 ch stream bank h by measuring | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy comparison and a maincaver and a maincaver and a maincaver (dense vegetation). Low 1.1 egories and Cond th and width. Cal | High Marginal: Non-maintained, dense hetbaceous vegetation with either a shrub layer or a tree layer (dbh either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F | ay be acceptable) bor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian | NOTES>> | | |
| Riparian Buffers Scores | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 Trian areas along ear uare footage for eac uare footage for eac sparian Area and Sc % Riparian Area> | nal 3 inches) present, canopy cover. within the riparian s. 5 5 6 6 6 6 6 6 7 7 7 8 7 7 7 7 7 7 7 7 7 7 7 7 7 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy course and a maintained understory. Recent cutowr (dense vegetation). Low 1.1 egories and Cond th and width. Calk he blocks below. 10% | High Marginal: Non-maintained, dense hetbaceous vegetation with either a shrub layer or a tree layer (dbh either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F | ay be acceptable) bor Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 | NOTES>> | ores*0.01)/2 | |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 Trian areas along ear uare footage for eac uare footage for eac sparian Area and Sc % Riparian Area> | nal 3 inches) present, canopy cover. within the riparian s. 5 5 6 6 6 6 6 6 7 7 7 8 7 7 7 7 7 7 7 7 7 7 7 7 7 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy course and a maintained understory. Recent cutowr (dense vegetation). Low 1.1 egories and Cond th and width. Calk he blocks below. 10% | High Marginal: Non-maintained, dense hetbaceous vegetation with either a shrub layer or a tree layer (dbh either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F | ay be acceptable) bor Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 | | ores*0.01)/2 0.72 | CI |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 1.5 1.5 1.5 1.5 1.5 1.5 1.5 | nal 3 inches) present, canopy cover. within the riparian s. 5 6 6 6 6 6 6 7 7 7 8 8 7 8 8 8 8 8 8 8 8 8 8 8 8 8 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% 0.75 | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy course and a maintained understory. Recent cutowr (dense vegetation). Low 1.1 egories and Cond th and width. Cale the blocks below. 10% 0.5 | gory Marg Marg High Marginal: Non-maintained, dense herbaceous vegetation with https://www.seland.org vegetation with seland.org vegetation with seland.org"/seland.org"/seland.org"/seland.org"/seland.org"/seland.org"/seland.org"/sela | measurements o ginal Low Marginal: Non-maintained, dense hetbaccous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F | ay be acceptable) toor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 100% | CI= (Sum % RA * Sc | | |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 1.5 1.5 1.5 1.5 1.5 1.5 1.5 | a inches) present, canopy cover. tithin the riparian s. ch stream bank by measuring core for each ripa 5% 0.6 10% 0.5 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% 0.75 | Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calc th and width. Calc th and width. Calc th and width. Calc th and width. Calc th an | gory High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provid 50% 0.6 | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrutu and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and tabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e | ay be acceptable) Cor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, active feed lots, trails, or other comparable conditions. Low 0.5 tiparian qual 100 100% 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > | 0.72 0.63 | CI |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 1.5 1.5 1.5 1.5 1.5 1.5 1.5 | a inches) present, canopy cover. tithin the riparian s. ch stream bank by measuring core for each ripa 5% 0.6 10% 0.5 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% 0.75 | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy comparison and a maincained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cale the and width. Cale the blocks below. 10% 0.5 10% 0.5 and depths; woody | gory High Marginal: Non-maintained, dense herbaceous vegetation with either a shrub layer or a tree layer (dbh > 3 inches) present, with <30% tree canopy cover. High 0.85 ition Scores using culators are provid 50% 0.6 | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrutu and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and tabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e | ay be acceptable) Cor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, active feed lots, trails, or other comparable conditions. Low 0.5 tiparian qual 100 100% 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > | 0.72 0.63 | CI |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN | Optin Tree stratum (dbh > : with > 60% tree Wetlands located w area: 1.5 1.5 1.5 1.5 1.5 1.5 1.5 1.5 | nal 3 inches) present, canopy cover. ithin the riparian s. 5 6 6 7 6 7 7 8 7 7 7 8 7 7 7 7 7 7 7 7 7 7 7 7 7 | Con Suboy High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% 0.75 30% 0.75 as, water velocity a | ditional Cate ptimal Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy comparison and a maincained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Cale the and width. Cale the blocks below. 10% 0.5 10% 0.5 and depths; woody | gory Marg Velocity Velocity Velocity Velocity Marg Velocity Vel | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrutu and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below. | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-til cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e | ay be acceptable) Cor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, active feed lots, trails, or other comparable conditions. Low 0.5 tiparian qual 100 100% 100% | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; S | 0.72 0.63 | CI |
| RIPARIAN Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN mplexes, stabl | Optin Tree stratum (dbh > : with > 60% tree (Wetlands located w area: 1.5 rian areas along ear vare footage for eac iparian Area and Sc % Riparian Area and Sc % Riparian Area Score > % Riparian Area Score > 1 HABITAT: Varie e features. | nal a inches) present, canopy cover. ithin the riparian s. ch stream bank th by measuring core for each ripr 5% 0.6 10% 0.5 ed substrate size nal et ypically present | Con Subop High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Catu or estimating leng arian category in th 85% 0.75 30% 0.75 as, water velocity a Stable habitat eler present in 30-50% c adequate for n | Low Suboptimal: Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calk the blocks below. 10% 0.5 10% 0.5 and depths; woody Conditional ptimal ments are typically of the reach and are understant cond the point of the reach and are ptimal | gory Marg Non-maintained, dense herbaceous vegetation with vegetation with s inches) present, with <30% tree canopy cover. High 0.85 diton Scores using culators are provid So0% 0.6 v and leafy debris; Stable habitat elep present in 10-30% c adequate for n | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with <30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below. stable substrate; ginal ments are typically of the reach and are | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-till cropland; actively grazed pasture, sparsely vegetated non-maintained area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure 1 of % F Blocks e low embededness Pc Habitat elements lacking or are typic | ay be acceptable) toor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 100% 100% ; shade; undercut coor elisted above are nstable. Habitat ally present in less | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; S | 0.72 0.63 | CI |
| Riparian Buffers Scores Delineate ripa Determine squ Enter the % R Right Bank Left Bank INSTREAN mplexes, stabl Instream Habitat/ Available | Optin Tree stratum (dbh > with > 60% tree (Wetlands located w area: 1.5 rian areas along ear uare footage for ear tiparian Area and Sc % Riparian Area> Score > % Riparian Area> % Riparian | nal a inches) present, canopy cover. ithin the riparian s. ch stream bank th by measuring core for each ripa 5% 0.6 10% 0.5 ed substrate size nal the present % of the reach. | Con Subop High Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and containing both herbaceous and shrub layers or a non-maintained understory. High 1.2 nto Condition Cate or estimating leng arian category in th 85% 0.75 30% 0.75 es, water velocity a Stable habitat eler present in 30-50% of adequate for n popula | Low Suboptimal: Low Suboptimal: Riparian areas with tree stratum (dbh > 3 inches) present, with 30% to 60% tree canopy cover and a maintained understory. Recent cutover (dense vegetation). Low 1.1 egories and Cond th and width. Calk the blocks below. 10% 0.5 10% 0.5 and depths; woody Conditional ptimal ments are typically of the reach and are understant cond the point of the reach and are ptimal | gory High Marginal: Non-maintained, dense hetbaceous vegetation with > 3 inches) present, with <30% | measurements o ginal Low Marginal: Non-maintained, dense herbaceous vegetation, riparian areas lacking shrub and tree stratum, hay production, ponds, open water. If present, tree stratum (dbh >3 inches) present, with ~30% tree canopy cover with maintained understory. Low 0.75 the descriptors. led for you below. stable substrate; ginal ments are typically of the reach and are | f length & width ma Pc High Poor: Lawns, mowed, and maintained areas, nurseries; no-til cropland; actively grazed pasture, sparsely vegetated area, recently seeded and stabilized, or other comparable condition. High 0.6 Ensure I of % F Blocks e Blocks e Habitat elements lacking or are u elements are typic than 10% c | ay be acceptable) oor Low Poor: Impervious surfaces, mine spoil lands, denuded surfaces, row crops, active feed lots, trails, or other comparable conditions. Low 0.5 he sums tiparian qual 100 100% 100% ; shade; undercut oor | CI= (Sum % RA * Sc Rt Bank CI > Lt Bank CI > banks; root mats; S | 0.72 0.63 GAV; riffle/pool | CI |

Reach R3 File: L\22000s\22800\22865.06\Admin\05-ENVR\Field Data\Spread I\Field Forms\S-GH6\1_QAQC\S-GH6 HGM_HG_R4R6_USM_Wolman.xlsx


| Project # | Stream Impac Project Name (Applicant) Local | | | ocality Cowardin Class. | HUC | Date | SAR # | Impact Length | Impact Factor | | |
|---|--|--|---|-------------------------|---|---------------------|-------------------|-------------------------|------------------|--------------|--|
| 22865.06 | Mountain Valley Pipeline (Mountain Valley Pipeline, LLC) | | Franklin County | R3 | 03010101 | 8/27/2021 S | S-GH6 | 20 | 1 | | |
| 1. CHANNEL | ALTERATION: Stream crossir | ngs, riprap, concret | e, gabions, or cor | ncrete blocks, stra | ightening of chann | el, channelization | , embankments, s | spoil piles, constricti | ions, livestock | | |
| | | Conditiona | | | | | | NOTES>> | | | |
| | Negligible | Mir | nor | 40 - 60% of reach | erate | Sev | /ere | | | | |
| Channel Alteration | Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized. | Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines. | 20-40% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines. | | 60 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered. | by any of the chann | | | | СІ | |
| Scores | 1.5 | 1.3 | 1.1 | 0.9 | 0.7 | 0 | .5 | | | 1.50 | |
| | REACH | CONDITION | INDEX and S | STREAM CO | NDITION UN | ITS FOR THI | S REACH | | | | |
| NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number. THE REACH CONDITION INDEX (RCI) >> | | | | | | | | | 1.09 | | |
| | | | | | | RCI= (Sum of | all Cl's)/5, exce | ept if stream is ep | ohemeral RCI = (| Riparian CI/ | |
| | | | | | | | COMPENSA | TION REQUIRE | MENT (CR) >> | 22 | |
| | | | | | | | | | | | |




PROVIDED UNDER SEPARATE COVER



| CL STAKEOUT POINTS: S-GH6 CROSS SECTION A (PIPE CL) | | | | | | | | |
|---|---------------|---------------|-----------|-------|-------|--|--|--|
| | PI | POST-CROSSING | | | | | | |
| PT. LOC. | NORTHING | FACTING | | VERT. | HORZ. | | | |
| | | EASTING | ELEV | DIFF. | DIFF. | | | |
| TS-L | 13469806.2870 | 1936888.0850' | 1263.781' | | | | | |
| BS-L | 13469805.0980 | 1936883.7770' | 1260.265' | | | | | |
| THW | 13469807.1770 | 1936880.6740' | 1260.371' | | | | | |
| BS-R | 13469805.8570 | 1936879.2810' | 1260.658' | | | | | |
| TS-R | 13469805.4640 | 1936876.6080' | 1263.143' | | | | | |





- LOCATIONS WERE COMPLETED ON SEPTEMBER 13, 2021.
- PIPELINE.
- GENERATE A CLEAN PRE-CROSSING SURFACE.

