Reach S-KL17 (Timber Mat Crossing) Intermittent Spread I Franklin County, Virginia

| Data | Included |
|---|-----------------|
| Photos | \checkmark |
| SWVM Form | \checkmark |
| FCI Calculator and HGM Form | N/A – Slope <4% |
| RBP Physical Characteristics Form | \checkmark |
| Water Quality Data | \checkmark |
| RBP Habitat Form | \checkmark |
| RBP Benthic Form | \checkmark |
| Benthic Identification Sheet | N/A – Low flow |
| Wolman Pebble Count | \checkmark |
| RiverMorph Data Sheet | \checkmark |
| USM Form (Virginia Only) | \checkmark |
| Longitudinal Profile and Cross Sections | \checkmark |

Stream S-KL17 (Timber Mat) Franklin County



Photo Type: DS VIEW Location, Orientation, Photographer Initials: Downstream view of ROW/LOC looking NE, JM



Photo Type: US VIEW Location, Orientation, Photographer Initials: Upstream view of ROW/LOC looking S, JM

DEQ Permit #21-0416

Spread I

Stream S-KL17 (Timber Mat) Franklin County



Photo Type: LB CL Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking SE, JM



Photo Type: RB CL Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking NW, JM

Spread I

Stream S-KL17 (Timber Mat) Franklin County



```
Photo Type: DS COND
Location, Orientation, Photographer Initials: Downstream conditions outside of ROW/LOC looking NE, JM
```

West Virginia Stream and Wetland Valuation Metric (SWVM) Version 2.1, September 2017

| USACE FILE NO./ Project Name: (v2.1, Sept 2015) | | Mountain | Valley Pipeline | | COORDINATES: imal Degrees) | Lat. | 37.031011 | Lon. | -79.778435 | WEATHER: | | Sunny | DATE: | August 1 | 17, 2021 |
|---|---------------------|------------------------|--|----------------------|-------------------------------|------|---|--|-------------|--|-----------------------------|---------------|--|-----------------------|-------------|
| IMPACT STREAM/SITE ID (watershed size (acreage), | | | S-ł | KL17 | | | MITIGATION STREAM CLAS (watershed size {acre | SS./SITE ID AND S age}, unaltered or impair | | | | | Comments: | | |
| STREAM IMPACT LENGTH: | 20 | FORM OF MITIGATION: | RESTORATION (Levels I-III) | | ORDINATES: imal Degrees) | Lat. | | Lon. | | PRECIPITATION PAST 48 HRS: | | Yes | Mitigation Length: | | |
| Column No. 1- Impact Existing | Condition (Del | bit) | Column No. 2- Mitigation Existing C | ondition - Basel | ine (Credit) | | Column No. 3- Mitigation Post Comple | | ears | Column No. 4- Mitigation Proj Post Completion (| ected at Ten Yea Credit) | ars | Column No. 5- Mitigation Project | ed at Maturity (C | redit) |
| Stream Classification: | Intern | nittent | Stream Classification: | | | | Stream Classification: | | 0 | Stream Classification: | | 0 | Stream Classification: | 0 | |
| Percent Stream Channel SI | оре | 3.03 | Percent Stream Channel Sl | ope | | | Percent Stream Channe | I Slope | 0 | Percent Stream Channel Si | lope | 0 | Percent Stream Channel S | lope | 0 |
| HGM Score (attach d | ata forms): | | HGM Score (attach | data forms): | | | HGM Score (atta | ch data forms): | | HGM Score (attach d | ata forms): | | HGM Score (attach o | ata forms): | |
| | | Average | | | Average | | | | Average | | | Average | | | Average |
| Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling | | 0 | | Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling | | 0 | Hydrology Biogeochemical Cycling | - | 0 |
| Habitat PART I - Physical, Chemical and | Biological India | cators | Habitat PART I - Physical, Chemical an | d Biological Indi | icators | | Habitat PART I - Physical, Chemica | I and Biological Ind | icators | Habitat PART I - Physical, Chemical and | Biological India | cators | Habitat PART I - Physical, Chemical and | d Biological Indic | ators |
| | Points Scale Range | Site Score | | Points Scale Range | Site Score | | | Points Scale Range | Site Score | | Points Scale Range | Site Score | | Points Scale Range | Site Score |
| PHYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all streams | classifications) | | | PHYSICAL INDICATOR (Applies to all stre | ams classifications) | | PHYSICAL INDICATOR (Applies to all streams | s classifications) | | PHYSICAL INDICATOR (Applies to all stream | is classifications) | |
| USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (Low Gradient Data Sheet) | | | | USEPA RBP (High Gradient Data Shee | n | | USEPA RBP (High Gradient Data Sheet) | | | USEPA RBP (High Gradient Data Sheet) | | |
| 1. Epifaunal Substrate/Available Cover | 0-20 | 6 | 1. Epifaunal Substrate/Available Cover | 0-20 | | | 1. Epifaunal Substrate/Available Cover | 0-20 | | 1. Epifaunal Substrate/Available Cover | 0-20 | | 1. Epifaunal Substrate/Available Cover | 0-20 | |
| 2. Embeddedness | 0-20 | 2 | 2. Pool Substrate Characterization | 0-20 | | | 2. Embeddedness | 0-20 | | 2. Embeddedness | 0-20 | | 2. Embeddedness | 0-20 | |
| 3. Velocity/ Depth Regime | 0-20 | 5 | 3. Pool Variability 4. Sediment Deposition | 0-20 | | | 3. Velocity/ Depth Regime | 0-20 | | 3. Velocity/ Depth Regime | 0-20 | | 3. Velocity/ Depth Regime 4. Sediment Deposition | 0-20 | |
| 4. Sediment Deposition 5. Channel Flow Status | 0-20 | 5 | 4. Sediment Deposition 5. Channel Flow Status | 0-20 | | | 4. Sediment Deposition 5. Channel Flow Status | 0-20 | | 4. Sediment Deposition 5. Channel Flow Status | 0-20 | | 4. Sediment Deposition 5. Channel Flow Status | 0-20 | |
| 6. Channel Alteration | 0-20 0-1 | 11 | 6. Channel Alteration | 0-20 0-1 | | | 6. Channel Alteration | 0-20 0-1 | | 6. Channel Alteration | 0-20 0-1 | | 6. Channel Alteration | 0-20 0-1 | |
| 7. Frequency of Riffles (or bends) | 0-20 | 1 | 7. Channel Sinuosity | 0-20 | | | 7. Frequency of Riffles (or bends) | 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | | 7. Frequency of Riffles (or bends) | 0-20 | |
| 8. Bank Stability (LB & RB) | 0-20 | 5 | 8. Bank Stability (LB & RB) | 0-20 | | | 8. Bank Stability (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) | 0-20 | | 8. Bank Stability (LB & RB) | 0-20 | |
| 9. Vegetative Protection (LB & RB) | 0-20 | 6 | 9. Vegetative Protection (LB & RB) | 0-20 | | | 9. Vegetative Protection (LB & RB) | 0-20 | | 9. Vegetative Protection (LB & RB) | 0-20 | | Vegetative Protection (LB & RB) | 0-20 | |
| 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | 4 | 10. Riparian Vegetative Zone Width (LB & RB) | 0-20 | | | Riparian Vegetative Zone Width (LB & RB | | | Riparian Vegetative Zone Width (LB & RB) | | | Riparian Vegetative Zone Width (LB & RB) | | |
| Total RBP Score Sub-Total | Poor | 48 0.24 | Total RBP Score Sub-Total | Poor | 0 | | Total RBP Score Sub-Total | Poor | 0 | Total RBP Score Sub-Total | Poor | 0 | Total RBP Score Sub-Total | Poor | 0 |
| CHEMICAL INDICATOR (Applies to Intermitter | nt and Perennial St | | CHEMICAL INDICATOR (Applies to Intermitten | t and Perennial Stre | eams) | | CHEMICAL INDICATOR (Applies to Interm | ittent and Perennial Str | eams) | CHEMICAL INDICATOR (Applies to Intermitte | nt and Perennial S | treams) | SUD- I otal CHEMICAL INDICATOR (Applies to Intermitte | ant and Perennial Str | • |
| WVDEP Water Quality Indicators (General | 1) | | WVDEP Water Quality Indicators (General) | | | | WVDEP Water Quality Indicators (Gen | eral) | | WVDEP Water Quality Indicators (Genera | I) | | WVDEP Water Quality Indicators (General |) (IL | |
| Specific Conductivity | | | Specific Conductivity | | | | Specific Conductivity | | | Specific Conductivity | | | Specific Conductivity | | |
| <=99 - 90 points | 0-90 | 54.7 | | 0-90 | | | | 0-90 | | | 0-90 | | | 0-90 | |
| N=99 - 90 points | - | | nH | | | | nH | | | nH | - | | pH | | |
| 20 | 0-80 | 7.55 | 21 | 5-90 0-1 | | | p., | 5-90 0-1 | | 20 | 5-90 0-1 | | p., | 5-90 0-1 | |
| 6.0-8.0 = 80 points | 0-80 | 7.55 | | P-80 | | | | 5-90 | | | P-A0 | | | 2-90 | |
| DO | | | DO | | | | DO | | | DO | | | DO | | |
| >5.0 = 30 points | 10-30 | 6.2 | | 10-30 | | | | 10-30 | | | 10-30 | | | 10-30 | |
| Sub-Total | | 1 | Sub-Total | | 0 | | Sub-Total | | 0 | Sub-Total | | 0 | Sub-Total | | 0 |
| BIOLOGICAL INDICATOR (Applies to Intermit | tent and Perennial | Streams) | BIOLOGICAL INDICATOR (Applies to Intermitte | ent and Perennial S | Streams) | | BIOLOGICAL INDICATOR (Applies to Int | ermittent and Perennia | al Streams) | BIOLOGICAL INDICATOR (Applies to Intern | nittent and Perenr | nial Streams) | BIOLOGICAL INDICATOR (Applies to Interr | nittent and Perenni | al Streams) |
| WV Stream Condition Index (WVSCI) | 1 1 | | WV Stream Condition Index (WVSCI) | | | | WV Stream Condition Index (WVSCI) | 1 1 | | WV Stream Condition Index (WVSCI) | 1 1 | | WV Stream Condition Index (WVSCI) | | |
| 0 | 0-100 0-1 | | | 0-100 0-1 | | | 1 | 0-100 0-1 | | | 0-100 0-1 | | | 0-100 0-1 | |
| Sub-Total | | 0 | Sub-Total | | 0 | | Sub-Total | | 0 | Sub-Total | | 0 | Sub-Total | | 0 |
| PART II - Index and U | Init Score | | PART II - Index and | Unit Score | | | PART II - Index a | and Unit Score | | PART II - Index and U | Init Score | | PART II - Index and | Jnit Score | |
| Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score | | Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score | Index | Linear Feet | Unit Score |
| 0.620 | 20 | 12.4 | 0 | 0 | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0.020 | | | ů. | L Č | | | | | ÷ | , , , , , , , , , , , , , , , , , , , | <u> </u> | , v | | | |

| PART II - Index and only score | | | | | |
|--------------------------------|-------------|------------|--|--|--|
| Index | Linear Feet | Unit Score | | | |
| 0.620 | 20 | 12.4 | | | |

| WVDEP Water Quality Indicators (General) | | | | WVD |
|---|-----------|------------|------------|-------|
| Specific Conductivity | | | | Spec |
| | 0-90 | | | |
| pH | | I | | pH |
| | 5-90 | 0-1 | | |
| DO | | t i | | DO |
| | 10-30 | | | |
| Sub-Total | | | 0 | Sub-1 |
| BIOLOGICAL INDICATOR (Applies to Intermitte | nt and Pe | rennial \$ | Streams) | BIOL |
| WV Stream Condition Index (WVSCI) | | | | wvs |
| | 0-100 | 0-1 | | |
| Sub-Total | | | 0 | Sub-1 |
| PART II - Index and | Jnit Sco | re | | |
| | | | | |
| Index | Linear | Feet | Unit Score | |

| PART II - Index and Unit Score | | | | | |
|--------------------------------|---|---|--|--|--|
| Index Linear Feet Unit Score | | | | | |
| 0 | 0 | 0 | | | |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME | REASON FOR SURVEY |

| WEATHER CONDITIONS | Now Past 24 hours Has there been a heavy rain in the last 7 days? Storm (heavy rain) rain (steady rain) showers (intermittent) Yes No % cloud cover clear/sunny Mas there been a heavy rain in the last 7 days? |
|----------------------------|---|
| SITE LOCATION/MAP | Draw a map of the site and indicate the areas sampled (or attach a photograph) |
| | Bridge |
| | LOD |
| STREAM CHARACTERIZATION | Stream Subsystem Perennial Stream Type Coldwater Warmwater Stream Origin Glacial Spring-fed Mixture of origins Swamp and bog Catchment Area_km ² |

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

| WATERSHED FEATURES RIPARIAN VEGETATION (18 meter buffer) | Predominant Surrounding Landuse Forest Commercial Field/Pasture Industrial Agricultural Other Residential Inductrial Indicate the dominant type and record the domin Trees | Local Watershed NPS Pollution No evidence Some potential sources Obvious sources Local Watershed Erosion None Moderate Heavy ant species present Grasses Herbaceous |
|--|--|---|
| INSTREAM FEATURES | Dominant species present | Canopy Cover Partly open Partly shaded Shaded High Water Mark m Proportion of Reach Represented by Stream Morphology Types Riffle % Run % Pool % Channelized Yes No Dam Present Yes |
| LARGE WOODY DEBRIS | LWDm ² Density of LWDm ² /km ² (LWD/ reac | h area) |
| AQUATIC VEGETATION | Indicate the dominant type and record the domin Rooted emergent Floating Algae Rooted submergent Attached Algae Dominant species present Portion of the reach with aquatic vegetation | Rooted floating Free floating |
| WATER QUALITY | Temperature0 C Specific Conductance Dissolved Oxygen pH Turbidity WQ Instrument Used | Water Odors Sewage Normal/None Sewage Petroleum Chemical Fishy Other Water Surface Oils Slick Slick Sheen Globs Flecks None Other Turbidity (if not measured) Clear Slightly turbid Clear Slightly turbid Turbid Opaque Stained Other |
| SEDIMENT/ SUBSTRATE | Odors Petroleum Normal Sewage Petroleum Chemical Anaerobic None Other | Deposits Sludge Sawdust Paper fiber Sand Relict shells Other |

| INC | ORGANIC SUBSTRATE (should add up to | | ORGANIC SUBSTRATE COMPONENTS (does not necessarily add up to 100%) | | | | |
|-------------------|--|------------------------------------|---|----------------------------|-----------------------------------|--|--|
| Substrate Type | Diameter | % Composition in Sampling Reach | Substrate Type | Characteristic | % Composition in Sampling Area | | |
| Bedrock | | | Detritus | sticks, wood, coarse plant | | | |
| Boulder | > 256 mm (10") | | | materials (CPOM) | | | |
| Cobble | 64-256 mm (2.5"-10") | | Muck-Mud | black, very fine organic | | | |
| Gravel | 2-64 mm (0.1"-2.5") | | | (FPOM) | | | |
| Sand | 0.06-2mm (gritty) | | Marl | grey, shell fragments | | | |
| Silt | 0.004-0.06 mm | |] | | | | |
| Clay | < 0.004 mm (slick) | | | | | | |

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

| STREAM NAME | LOCATION | |
|---------------------|--------------------|-------------------|
| STATION # RIVERMILE | STREAM CLASS | |
| LAT LONG | RIVER BASIN | |
| STORET # | AGENCY | |
| INVESTIGATORS | | |
| FORM COMPLETED BY | DATE TIME AM PM | REASON FOR SURVEY |

| | Habitat | | Condition | ı Category | |
|--|---|---|---|---|---|
| | Parameter | Optimal | Suboptimal | Marginal | Poor |
| | 1. Epifaunal Substrate/ Available Cover | Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are <u>not</u> new fall and <u>not</u> transient). | 40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale). | 20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed. | Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| n sampling reach | 2. Embeddedness | Gravel, cobble, and boulder particles are 0- 25% surrounded by fine sediment. Layering of cobble provides diversity of niche space. | Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment. | Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment. | Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment. |
| ted i | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| Parameters to be evaluated in sampling reach | 3. Velocity/Depth Regime | All four velocity/depth regimes present (slow- deep, slow-shallow, fast- deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.) | Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes). | Only 2 of the 4 habitat regimes present (if fast- shallow or slow-shallow are missing, score low). | Dominated by 1 velocity/ depth regime (usually slow-deep). |
| uram | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| P | 4. Sediment Deposition | Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition. | Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools. | Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent. | Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| | 5. Channel Flow Status | Water reaches base of both lower banks, and minimal amount of channel substrate is exposed. | Water fills >75% of the available channel; or <25% of channel substrate is exposed. | Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed. | Very little water in channel and mostly present as standing pools. |
| | SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |

Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates, and Fish, Second Edition - Form 2

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

| Habitat | | Condition | ı Category | |
|---|--|--|--|---|
| Parameter | Optimal | Suboptimal | Marginal | Poor |
| 6. Channel Alteration | Channelization or dredging absent or minimal; stream with normal pattern. | Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present. | Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted. | Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely. |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| 7. Frequency of Riffles (or bends) | Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important. | Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15. | Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25. | Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25. |
| SCORE | 20 19 18 17 16 | 15 14 13 12 11 | 10 9 8 7 6 | 5 4 3 2 1 0 |
| SCORE 8. Bank Stability (score each bank) Note: determine left or right side by facing downstream. SCORE (LB) SCORE (RB) 9. Vegetative Protection (score each bank) | Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected. | Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion. | Moderately unstable; 30- 60% of bank in reach has areas of erosion; high erosion potential during floods. | Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| 9. Vegetative Protection (score each bank) | More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally. | 70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well- represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining. | 50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining. | Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE (RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| 10. Riparian Vegetative Zone Width (score each bank riparian zone) | Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone. | Width of riparian zone 12-18 meters; human activities have impacted zone only minimally. | Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal. | Width of riparian zone <6 meters: little or no riparian vegetation due to human activities. |
| SCORE (LB) | Left Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |
| SCORE(RB) | Right Bank 10 9 | 8 7 6 | 5 4 3 | 2 1 0 |

Total Score _____

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

| STREAM NAME | | LOCATION | |
|----------------------|---|--|-------------------|
| STATION # | _ RIVERMILE | STREAM CLASS | |
| LAT | LONG | RIVER BASIN | |
| STORET # | | AGENCY | |
| INVESTIGATORS | | | LOT NUMBER |
| FORM COMPLETED | BY | DATE TIME | REASON FOR SURVEY |
| HABITAT TYPES | Indicate the percentage of Cobble% Sn Submerged Macrophytes | ags% Vegetated B | anks% Sand%)% |
| SAMPLE COLLECTION | Indicate the number of jab | lected? wading fi ps/kicks taken in each habitat ty lags Vegetated B | anks Sand |
| GENERAL COMMENTS | | | |

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3= Abundant, 4 = Dominant

| Periphyton | 0 | 1 | 2 | 3 | 4 | Slimes | 0 | 1 | 2 | 3 | 4 |
|-------------------|---|---|---|---|---|--------------------|---|---|---|---|---|
| Filamentous Algae | 0 | 1 | 2 | 3 | 4 | Macroinvertebrates | 0 | 1 | 2 | 3 | 4 |
| Macrophytes | 0 | 1 | 2 | 3 | 4 | Fish | 0 | 1 | 2 | 3 | 4 |

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3= Abundant (>10 organisms), 4 = Dominant (>50 organisms)

| Porifera | 0 | 1 | 2 | 3 | 4 | Anisoptera | 0 | 1 | 2 | 3 | 4 | Chironomidae | 0 | 1 | 2 | 3 | 4 |
|-----------------|---|---|---|---|---|-------------|---|---|---|---|---|---------------|---|---|---|---|---|
| Hydrozoa | 0 | 1 | 2 | 3 | 4 | Zygoptera | 0 | 1 | 2 | 3 | 4 | Ephemeroptera | 0 | 1 | 2 | 3 | 4 |
| Platyhelminthes | 0 | 1 | 2 | 3 | 4 | Hemiptera | 0 | 1 | 2 | 3 | 4 | Trichoptera | 0 | 1 | 2 | 3 | 4 |
| Turbellaria | 0 | 1 | 2 | 3 | 4 | Coleoptera | 0 | 1 | 2 | 3 | 4 | Other | 0 | 1 | 2 | 3 | 4 |
| Hirudinea | 0 | 1 | 2 | 3 | 4 | Lepidoptera | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Oligochaeta | 0 | 1 | 2 | 3 | 4 | Sialidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Isopoda | 0 | 1 | 2 | 3 | 4 | Corydalidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Amphipoda | 0 | 1 | 2 | 3 | 4 | Tipulidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Decapoda | 0 | 1 | 2 | 3 | 4 | Empididae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Gastropoda | 0 | 1 | 2 | 3 | 4 | Simuliidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| Bivalvia | 0 | 1 | 2 | 3 | 4 | Tabinidae | 0 | 1 | 2 | 3 | 4 | | | | | | |
| | | | | | | Culcidae | 0 | 1 | 2 | 3 | 4 | | | | | | |

WOLMAN PEBBLE COUNT FORM

Basin:

County:Franklin CountyStream Name:UNT to Foul Ground CreekHUC Code:03010101Survey Date:8/17/2021Surveyors:SK, VM, RHType:Representative

Stream ID:

S-KL17

Upper Roanoke

| T I | DADTICLE | | LE COUNT | Dent 1 | T-4 1 // | T4 0/ | 0/ 5 |
|-------------|-------------|-------------|----------|-------------------|----------|--------|-------|
| Inches | PARTICLE | Millimeters | | Particle Count | Total # | Item % | % Cur |
| | Silt/Clay | < .062 | S/C | ^ | 12 | 12.00 | 12.00 |
| | Very Fine | .062125 | | ▲ ▼ | | 0.00 | 12.00 |
| | Fine | .12525 | | * * | | 0.00 | 12.00 |
| | Medium | .255 | SAND | ▲ ▼ | 1 | 1.00 | 13.00 |
| | Coarse | .50-1.0 | | ▲ ▼ | | 0.00 | 13.00 |
| .0408 | Very Coarse | 1.0-2 | | ▼ | 2 | 2.00 | 15.00 |
| .0816 | Very Fine | 2 -4 | | | 8 | 8.00 | 23.00 |
| .1622 | Fine | 4 -5.7 | | ▲ ▼ | 9 | 9.00 | 32.00 |
| .2231 | Fine | 5.7 - 8 | | ▲ ▼ | 4 | 4.00 | 36.00 |
| .3144 | Medium | 8 -11.3 | GRAVEL | * | 4 | 4.00 | 40.00 |
| .4463 | Medium | 11.3 - 16 | | ▼ | 7 | 7.00 | 47.00 |
| .6389 | Coarse | 16 -22.6 | | ▲ ▼ | 3 | 3.00 | 50.00 |
| .89 - 1.26 | Coarse | 22.6 - 32 | | ▲ ▼ | 7 | 7.00 | 57.00 |
| 1.26 - 1.77 | Vry Coarse | 32 - 45 | | ▲ ▼ | 15 | 15.00 | 72.00 |
| 1.77 -2.5 | Vry Coarse | 45 - 64 | | ▲ ▼ | 12 | 12.00 | 84.00 |
| 2.5 - 3.5 | Small | 64 - 90 | | ▲ ▼ | 12 | 12.00 | 96.00 |
| 3.5 - 5.0 | Small | 90 - 128 | | ▲ ▼ | 4 | 4.00 | 100.0 |
| 5.0 - 7.1 | Large | 128 - 180 | COBBLE | ▲ ▼ | | 0.00 | 100.0 |
| 7.1 - 10.1 | Large | 180 - 256 | | ▼ | | 0.00 | 100.0 |
| 10.1 - 14.3 | Small | 256 - 362 | | ↓ | | 0.00 | 100.0 |
| 14.3 - 20 | Small | 362 - 512 | 1 | • | | 0.00 | 100.0 |
| 20 - 40 | Medium | 512 - 1024 | BOULDER | ₹ | | 0.00 | 100.0 |
| 40 - 80 | Large | 1024 -2048 | 1 | ▲ ▼ | | 0.00 | 100.0 |
| 80 - 160 | Vry Large | 2048 -4096 | 1 | ₹ | | 0.00 | 100.0 |
| | Bedrock | | BDRK | ÷ | | 0.00 | 100.0 |
| | | | | Totals: | 100 | | |

| | UNT to Foul G G-KL17 Representative 08/17/2021 | | κ |
|---|---|--|--|
| Size (mm) | тот # | ITEM % | CUM % |
| 0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock | $ \begin{array}{r} 12 \\ 0 \\ 0 \\ 1 \\ 0 \\ 2 \\ 8 \\ 9 \\ 4 \\ 4 \\ 7 \\ 3 \\ 7 \\ 15 \\ 12 \\ 12 \\ 4 \\ 0 \\ $ | 12.00 0.00 0.00 1.00 0.00 2.00 8.00 9.00 4.00 4.00 7.00 3.00 7.00 15.00 12.00 12.00 12.00 4.00 0.00 | 12.00 12.00 12.00 13.00 13.00 13.00 23.00 32.00 36.00 40.00 47.00 50.00 57.00 72.00 84.00 96.00 100.00 100.00 100.00 100.00 100.00 100.00 100.00 |
| D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Gravel (%) Boulder (%) Bedrock (%) | 2.25 7.42 22.6 64 87.83 128 12 3 69 16 0 0 | | |

Total Particles = 100.

| | | | For use in wadea | ble channels cla | ssified as interm | ittent or perennia | | | | |
|----------------------|--|---|--------------------|---|---|---|---|--|---|---|
| Project # | Project Name (App | licant) | Locality | Cowardin Class. | HUC | Date | SAR # | Impact Length | Impact Factor | |
| 22865.06 | Mountain Valley Pipeline Valley Pipeline, L | • | Franklin County | R4 | 03010101 | 8/17/2021 | S-KL17 | 20 | 1 | |
| Nam | e(s) of Evaluator(s) | Stream Name | e and Informat | tion | | | | SAR Length | | |
| | JM, team of 8 | UNT to Foul | Ground Creek | ζ. | | | | 74 | | |
| . Channel C | condition: Assess the cross-section | on of the stream ar | | | , | | | | | |
| | Optimal | Subo | ptimal | Conditional Catego | ^{ry} ginal | Ba | or | Sev | oro | |
| Channel Condition | less than 10% of bottom. | erosion or unprotect of banks are st Vegetative protect prominent (60- Depositional feat stability. The ban channels are well de has access to bankfin developed floo | om. | Poor. Banks more st Poor due to low Erosion may be pre both banks. Veget 40-60% of banks. St vertical or unde 40-60% Sediment r transient, contri Deposition that cor may be forming/pre | er bank slopes. esent on 40-60% of ative protection on treambanks may be ercut. AND/OR may be temporary / bute instability. htribute to stability, esent. AND/OR V- have vegetative % of the banks and s which contribute to | laterally unstable further. Majority of vertical. Erosion pr banks. Vegetative on 20-40% of bank to prevent erosion. the stream is cove Sediment is temp nature, and contril AND/OR V-shap vegetative protect 40% of the banks a | ered by sediment. orary / transient in outing to instability. ed channels have ion is present on > and stable sediment | incision, flow contain Streambed below av majority of banks Vegetative protecti than 20% of banks erosion. Obvious present. Erosion/raw AND/OR Aggradin | stability. Severe ned within the banks. erage rooting depth, vertical/undercut. fon present on less s, is not preventing s bank sloughing v banks on 80-100%. g channel. Greater n bed is covered by uting to instability. channels and/or | |
| Scores | 2 | 2. | ٨ | | - - | 1 | .6 | | | 2 |

2. RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) NOTES>> **Conditional Category** Marginal Optimal **Suboptimal** Poor High Poor: Lawns Low Marginal: Non-maintained, mowed, and High Suboptimal: Low Suboptimal: **High Marginal:** dense herbaceous maintained areas, Low Poor: Riparian areas with Riparian areas with vegetation, riparian nurseries; no-till Impervious Non-maintained, tree stratum (dbh > tree stratum (dbh > surfaces, mine dense herbaceous areas lacking shrub cropland; actively 3 inches) present, 3 inches) present, Tree stratum (dbh > 3 inches) present, and tree stratum, grazed pasture, spoil lands, vegetation with Riparian with 30% to 60% with 30% to 60% with > 60% tree canopy cover. either a shrub layer hay production, sparsely vegetated denuded surfaces, tree canopy cover tree canopy cover **Buffers** Wetlands located within the riparian or a tree layer (dbh ponds, open water. non-maintained row crops, active and containing both and a maintained > 3 inches) feed lots, trails, or If present, tree area, recently areas. herbaceous and understory. Recent present, with <30% stratum (dbh >3 seeded and other comparable cutover (dense shrub layers or a inches) present, stabilized, or other conditions. tree canopy cover. non-maintained vegetation). with <30% tree comparable understory. canopy cover with condition. maintained understory. High High High Low Low Low 1.5 1.2 1.1 0.85 0.75 0.6 0.5 Scores Delineate riparian areas along each stream bank into Condition Categories and Condition Scores using the descriptors. Ensure the sums of % Riparian 2. Determine square footage for each by measuring or estimating length and width. Calculators are provided for you below. 3. Enter the % Riparian Area and Score for each riparian category in the blocks below. Blocks equal 100 Assessment is limited to % Riparian Area> 100% 100% areas within the temporary **Right Bank** 1.1 ROW Score > CI= (Sum % RA * Scores*0.01)/2 100% 100% CI % Riparian Area> Rt Bank CI > 1.10 Left Bank 1.1 1.10 Lt Bank CI > 1.10 Score > 3. INSTREAM HABITAT: Varied substrate sizes, water velocity and depths; woody and leafy debris; stable substrate; low embededness; shade; undercut banks; root mats; SAV; riffle/pool complexes, stable features. **Conditional Category** NOTES>> **Optimal Suboptimal** Marginal Poor Instream Habitat/

| Scores | 1.5 | 1.2 | 0.9 | 0.5 | High / Low | 1.20 |
|-----------|--------------------------------|---|---|--|-----------------|------|
| | | | | | Stream Gradient | CI |
| Cover | greater than 50% of the reach. | adequate for maintenance of populations. | adequate for maintenance of populations. | elements are typically present in less than 10% of the reach. | | |
| Available | 51 51 | • | present in 10-30% of the reach and are | 5 | | |
| | | Stable habitat elements are typically | Stable habitat elements are typically | Habitat elements listed above are | | |

Reach R3-R4

File: https://tetratechinc.sharepoint.com/teams/MVPStreamWetlandAssessment/Shared Documents/General/01. Virginia Field Data Management/03. Preliminary QAQC (working files)/S-KL17_20210915KEH/9. S-KL17_USM_MVP_20210917KEH.xlsx

| Project # | Project Name (App | licant) | Locality | Cowardin Class. | HUC | Date | SAR # / Data Point | Impact / SAR length | Impact Factor | |
|-----------------------|--|--|--------------------|---|------------------------------------|---|--|-------------------------|------------------|-------------|
| 22865.06 | Mountain Valley Pipeline Valley Pipeline, L | • | Franklin County | R4 | 03010101 | 8/17/2021 | S-KL17 | 20 | 1 | |
| . CHANNEI | ALTERATION: Stream crossin | gs, riprap, concret | e, gabions, or con | crete blocks, strai | ghtening of channe | el, channelization, | , embankments, sp | oil piles, constrictior | ns, livestock | |
| | | | Conditiona | al Category | | | | NOTES>> | | |
| | Negligible | Mir | nor | | erate | Sev | vere | | | |
| Channel Alteration | Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized. | Less than 20% of the stream reach is disrupted by any of the channel alterations listed in the parameter guidelines. | the channel | 40 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered. | is disrupted by any of the channel | by any of the chan in the parameter (80% of banks sh | of reach is disrupted nel alterations listed guidelines AND/OR hored with gabion, or cement. | | | CI |
| Scores | 1.5 | 1.3 | 1.1 | 0.9 | 0.7 | 0 | .5 | | | 1.50 |
| | REACH | CONDITION | INDEX and S | STREAM CO | NDITION UN | ITS FOR TH | IS REACH | | | |
| <i>OTE:</i> The Cls a | nd RCI should be rounded to 2 decir | nal places. The CR | R should be round | ed to a whole num | ıber. | | THE REACH | H CONDITION IND | DEX (RCI) >> | 1.24 |
| | | | | | | RCI= (Sum o | f all Cl's)/5, exce | ept if stream is eph | nemeral RCI = (F | Riparian Cl |
| | | | | | | | COMPENSA | ATION REQUIREM | /IENT (CR) >> | 25 |
| | | | | | | | CR = RC | | | |

INSERT PHOTOS:





DESCRIBE PROPOSED IMPACT:


PROVIDED UNDER SEPARATE COVER


Reach R3-R4

File: https://tetratechinc.sharepoint.com/teams/MVPStreamWetlandAssessment/Shared Documents/General/01. Virginia Field Data Management/03. Preliminary QAQC (working files)/S-KL17_20210915KEH/9. S-KL17_USM_MVP_20210917KEH.xlsx









BENCHMARK POINT (WSSI)