Baseline Assessment – Stream Attributes

Reach S-Q2 (Timber Mat Crossing) Perennial Spread I Pittsylvania County, Virginia

Data	Included
Photos	✓
SWVM Form	✓
FCI Calculator and HGM Form	N/A – Perennial stream (not shadeable, slope less
	than 4%)
RBP Physical Characteristics Form	✓
Water Quality Data	✓
RBP Habitat Form	✓
RBP Benthic Form	✓
Benthic Identification Sheet	N/A – No Riffles
Wolman Pebble Count	✓
RiverMorph Data Sheet	✓
USM Form (Virginia Only)	✓
Longitudinal Profile and Cross Sections	✓



Location, Orientation, Photographer Initials: Downstream view of ROW looking NE, CB/BH



Location, Orientation, Photographer Initials: Upstream view of ROW looking SW, CB/BH



Location, Orientation, Photographer Initials: Standing on LB looking at RB along pipe centerline looking N, CB/BH



Location, Orientation, Photographer Initials: Standing on RB looking at LB along pipe centerline looking N, CB/BH



Photo Type: DS COND Location, Orientation, Photographer Initials: Downstream conditions outside of ROW looking NE, CB/BH

USACE FILE NO / Project Name: Mountain Valley Pipeline (v2.1, Sept 2015)			cimal Degrees)	Lat.	36.884284	Lon.	-79.427914	WEATHER:		Sunny	DATE:	Augu	ust 19, 2021			
IMPACT STREAM/SITE I (watershed size (acreage				ZS-Q2; ·	107.3 ac			MITIGATION STREAM CLAS (watershed size (acres						Comments:		
STREAM IMPACT LENGTH:	20	FORM MITIGAT		RESTORATION (Levels I-III)		OORDINATES: cimal Degrees)	Lat.		Lon.		PRECIPITATION PAST 48 HRS:		Yes	Mitigation Length:		
Column No. 1- Impact Existing	ng Condition (Deb	oit)		Column No. 2- Mitigation Existing Co	ndition - Base	line (Credit)		Column No. 3- Mitigation Post Complet		Years	Column No. 4- Mitigation Proj Post Completion (ears	Column No. 5- Mitigation Pro	jected at Maturit	y (Credit)
Stream Classification:	Pere	nnial		Stream Classification:				Stream Classification:		0	Stream Classification:		0	Stream Classification:		0
Percent Stream Channel S	Slope	1.98		Percent Stream Channel Slo	ре			Percent Stream Channel	Slope	0	Percent Stream Channel S	lope	0	Percent Stream Chann	al Slope	0
HGM Score (attach	data forms):			HGM Score (attach d	ata forms):			HGM Score (attac	ch data forms):		HGM Score (attach d	ata forms):		HGM Score (attac	h data forms):	
		Average				Average				Average			Average			Average
Hydrology				Hydrology				Hydrology			Hydrology			Hydrology		
Biogeochemical Cycling Habitat		0		Biogeochemical Cycling Habitat		0		Biogeochemical Cycling Habitat	_	0	Biogeochemical Cycling Habitat		0	Biogeochemical Cycling Habitat		_ °
PART I - Physical, Chemical an	nd Biological Indic	ators		PART I - Physical, Chemical and	Biological Ind	licators		PART I - Physical, Chemical	and Biological I	ndicators	PART I - Physical, Chemical and	Biological Ind	icators	PART I - Physical, Chemical	and Biological In	ndicators
	Points Scale Range	Site Score			Points Scale Range	Site Score			Points Scale Range	ge Site Score		Points Scale Range	s Site Score		Points Scale R	Range Site Score
PHYSICAL INDICATOR (Applies to all stream	ms classifications)			PHYSICAL INDICATOR (Applies to all streams of	lassifications)			PHYSICAL INDICATOR (Applies to all stres	ıms classifications)		PHYSICAL INDICATOR (Applies to all stream	s classifications)		PHYSICAL INDICATOR (Applies to all str	ams classifications	·)
USEPA RBP (High Gradient Data Sheet)				USEPA RBP (Low Gradient Data Sheet)				USEPA RBP (High Gradient Data Sheet			USEPA RBP (High Gradient Data Sheet)			USEPA RBP (High Gradient Data She		
	0-20	15			0-20			Epifaunal Substrate/Available Cover	0-20		Epifaunal Substrate/Available Cover	0-20		Epifaunal Substrate/Available Cover		
Embeddedness Velocity/ Depth Regime	0-20	- 6		Pool Substrate Characterization Pool Variability	0-20			Embeddedness Velocity/ Depth Regime	0-20		Embeddedness Velocity/ Depth Regime	0-20		Embeddedness Velocity/ Depth Regime	0-20 0-20	
Sediment Deposition	0-20	6		4. Sediment Deposition	0-20			4. Sediment Deposition	0-20		Sediment Deposition	0-20		Sediment Deposition	0-20	
5. Channel Flow Status	0-20	6		5. Channel Flow Status	0-20			5. Channel Flow Status	0-20		5. Channel Flow Status	0-20		5. Channel Flow Status	0-20	0.4
6. Channel Alteration	0-20	11		6. Channel Alteration	0-20			Channel Alteration	0-20		Channel Alteration	0-20		6. Channel Alteration	0-20	0-1
Frequency of Riffles (or bends)	0-20	7		7. Channel Sinuosity	0-20			7. Frequency of Riffles (or bends)	0-20		Frequency of Riffles (or bends)	0-20		Frequency of Riffles (or bends)	0-20	
8. Bank Stability (LB & RB)	0-20	10 12		8. Bank Stability (LB & RB)	0-20			8. Bank Stability (LB & RB)	0-20		8. Bank Stability (LB & RB)	0-20		8. Bank Stability (LB & RB)	0-20	
Vegetative Protection (LB & RB) Regetative Zone Width (LB & RB)	0-20	12		Vegetative Protection (LB & RB) Repartan Vegetative Zone Width (LB & RB)	0-20			Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20		Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & RB)	0-20		Vegetative Protection (LB & RB) Riparian Vegetative Zone Width (LB & R	0-20 B) 0-20	
Total RBP Score	Marginal	92		Total RBP Score	Poor	0		Total RBP Score	Poor	0	Total RBP Score	Poor	0	Total RBP Score	Poor	0
Sub-Total	magna	0.46		Sub-Total	1 001	0		Sub-Total	1 001	Ö	Sub-Total	1 001	ŏ	Sub-Total	1 001	Ö
CHEMICAL INDICATOR (Applies to Intermitt	tent and Perennial Str	reams)		CHEMICAL INDICATOR (Applies to Intermittent	and Perennial Str	reams)		CHEMICAL INDICATOR (Applies to Intermi	ttent and Perennial	Streams)	CHEMICAL INDICATOR (Applies to Intermitte	ant and Perennial	Streams)	CHEMICAL INDICATOR (Applies to Inter	nittent and Perennia	al Streams)
WVDEP Water Quality Indicators (Gener	al)			WVDEP Water Quality Indicators (General)				WVDEP Water Quality Indicators (Gene	ral)		WVDEP Water Quality Indicators (General	al)		WVDEP Water Quality Indicators (Ger	neral)	
Specific Conductivity				Specific Conductivity				Specific Conductivity			Specific Conductivity			Specific Conductivity		
<=99 - 90 points	0-90	89.3			0-90				0-90			0-90			0-90	1
pH ~ 99 - 90 points				pH				pН			pH	-		рH	_	
	0-80	7.19			5-90				5-90 0-1		F.:	5-90 0-1			5-90	0-1
6.0-8.0 = 80 points		7.10			00							1				
DO				00				DO			DO	_		DO	$\overline{}$	
>5.0 = 30 points	10-30	7.75			10-30				10-30			10-30			10-30	1
Sub-Total		1		Sub-Total		0		Sub-Total		0	Sub-Total		0	Sub-Total		0
BIOLOGICAL INDICATOR (Applies to Intern	nittent and Perennial	Streams)		BIOLOGICAL INDICATOR (Applies to Intermitte	nt and Perennial	Streams)		BIOLOGICAL INDICATOR (Applies to Inte	rmittent and Peren	nnial Streams)	BIOLOGICAL INDICATOR (Applies to Intern	nittent and Peren	nnial Streams)	BIOLOGICAL INDICATOR (Applies to Ir	termittent and Per	ennial Streams)
WV Stream Condition Index (WVSCI)				WV Stream Condition Index (WVSCI)				WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)			WV Stream Condition Index (WVSCI)		
	0-100 0-1				0-100 0-1				0-100 0-1	1		0-100 0-1			0-100	0-1
Sub-Total		0		Sub-Total		0		Sub-Total		0	Sub-Total		0	Sub-Total	-	0
PART II - Index and	Unit Score		1	PART II - Index and U	Init Score			PART II - Index a	and Unit Score		PART II - Index and L	Jnit Score		PART II - Index a	nd Unit Score	
Index	Linear Feet	Unit Score		Index	Linear Feet	Unit Score		Index	Linear Feet	t Unit Score	Index	Linear Feet	Unit Score	Index	Linear Fe	eet Unit Score
0.730	20	14.6		0	0	0		0	0	0	0	0	0	0	0	0

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (FRONT)

STREAM NAME	LOCATION				
STATION # RIVERMILE	STREAM CLASS				
LAT LONG	RIVER BASIN				
STORET#	AGENCY				
INVESTIGATORS					
FORM COMPLETED BY	DATE TIME	REASON FOR SURVEY			

WEATHER CONDITIONS	Now storm (heavy rain) rain (steady rain) showers (intermittent) —% %cloud cover clear/sunny	Past 24 hours Has there been a heavy rain in the last 7 days? Yes No Air Temperature0 C 0 Other
SITE LOCATION/MAP	Te Stream	
STREAM CHARACTERIZATION	Stream Origin	Stream Type dal Coldwater Warmwater Catchment Areakm² ed of origins

PHYSICAL CHARACTERIZATION/WATER QUALITY FIELD DATA SHEET (BACK)

WATERS FEATURI		Fores Field Agric	Pasture Industri	ercial	No evidence Son Obvious sources Local Watershed Erosi None Moderate	ne potential sources		
RIPARIA VEGETA (18 meter	ΓΙΟΝ	Trees	e the dominant type and S ant species present	hrubs		rbaceous		
INSTREA FEATURI			ted Reach Length		Canopy Cover Partly open Part	ly shaded Shaded		
				m	High Water Mark	m		
					Proportion of Reach Re	epresented by Stream		
			km² (m²x1000) ted Stream Depth	km²	Morphology Types Riffle Pool %	Run%		
			Velocity		Channelized Yes	No		
		(111 11111			Dam Present Yes	No		
LARGE V DEBRIS	VOODY		m² of LWDn	n ² /km ² (LWD /	reach area)			
AQUATIO VEGETA		Indicate Roote Floati Domina	e the dominant type and demergent R ng Algae A	l record the do ooted submerge ttached Algae	minant species present nt Rooted floating	C		
		Portion	of the reach with aqua	tic vegetation _	%			
WATER (QUALITY	Specific	rature0 C Conductance	-	Water Odors Normal/None Sewage Petroleum Fishy	Chemical Other		
		рН	ed Oxygen		Water Surface Oils Slick Sheen None Other	Globs Flecks		
			strument Used		Turbidity (if not measu Clear ☐ Slightly tur Opaque Stained	r ed) rbid Turbid Other		
SEDIMEN SUBSTRA		Odors Norm Chem		Petroleum None	Deposits Sludge Sawdust Relict shells	Sludge Sawdust Paper fiber Sand		
		Oils Abser		te Profu	are the undersides blac	h are not deeply embedded, k in color?		
INC	ORGANIC SUBS		COMPONENTS (00%)		ORGANIC SUBSTRATE C (does not necessarily add			
Substrate Type	Diamete	er	% Composition in Sampling Reach	Substrate Type	Characteristic	% Composition in Sampling Area		
Bedrock				Detritus	sticks, wood, coarse plant			
Boulder	> 256 mm (10")				materials (CPOM)			
Cobble	64-256 mm (2.5	"-10")		Muck-Mud	black, very fine organic			

Gravel

Sand

Silt

Clay

2-64 mm (0.1"-2.5")

0.06-2mm (gritty)

< 0.004 mm (slick)

0.004-0.06 mm

grey, shell fragments

Marl

HABITAT ASSESSMENT FIELD DATA SHEET - HG - USE ON ALL STREAMS (FRONT)

STREAM NAME	LOCATION				
STATION # RIVERMILE	STREAM CLASS				
LAT LONG	RIVER BASIN				
STORET#	AGENCY				
INVESTIGATORS					
FORM COMPLETED BY	DATE AM PM	REASON FOR SURVEY			

	Habitat		Condition	ı Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor			
	1. Epifaunal Substrate/ Available Cover	Greater than 70% of substrate favorable for epifaunal colonization and fish cover; mix of snags, submerged logs, undercut banks, cobble or other stable habitat and at stage to allow full colonization potential (i.e., logs/snags that are not new fall and not transient).	40-70% mix of stable habitat; well-suited for full colonization potential; adequate habitat for maintenance of populations; presence of additional substrate in the form of newfall, but not yet prepared for colonization (may rate at high end of scale).	20-40% mix of stable habitat; habitat availability less than desirable; substrate frequently disturbed or removed.	Less than 20% stable habitat; lack of habitat is obvious; substrate unstable or lacking.			
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
n sampling reach	2. Embeddedness	Gravel, cobble, and boulder particles are 0-25% surrounded by fine sediment. Layering of cobble provides diversity of niche space.	Gravel, cobble, and boulder particles are 25- 50% surrounded by fine sediment.	Gravel, cobble, and boulder particles are 50- 75% surrounded by fine sediment.	Gravel, cobble, and boulder particles are more than 75% surrounded by fine sediment.			
ted in	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Parameters to be evaluated in sampling reach	3. Velocity/Depth Regime	All four velocity/depth regimes present (slow-deep, slow-shallow, fast-deep, fast-shallow). (Slow is < 0.3 m/s, deep is > 0.5 m.)	Only 3 of the 4 regimes present (if fast-shallow is missing, score lower than if missing other regimes).	Only 2 of the 4 habitat regimes present (if fast-shallow or slow-shallow are missing, score low).	Dominated by 1 velocity/depth regime (usually slow-deep).			
ıram	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Pa	4. Sediment Deposition	Little or no enlargement of islands or point bars and less than 5% of the bottom affected by sediment deposition.	Some new increase in bar formation, mostly from gravel, sand or fine sediment; 5-30% of the bottom affected; slight deposition in pools.	Moderate deposition of new gravel, sand or fine sediment on old and new bars; 30-50% of the bottom affected; sediment deposits at obstructions, constrictions, and bends; moderate deposition of pools prevalent.	Heavy deposits of fine material, increased bar development; more than 50% of the bottom changing frequently; pools almost absent due to substantial sediment deposition.			
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
	5. Channel Flow Status	Water reaches base of both lower banks, and minimal amount of channel substrate is exposed.	Water fills >75% of the available channel; or <25% of channel substrate is exposed.	Water fills 25-75% of the available channel, and/or riffle substrates are mostly exposed.	Very little water in channel and mostly present as standing pools.			
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			

HABITAT ASSESSMENT FIELD DATA SHEET—HIGH GRADIENT STREAMS (BACK)

	Habitat		Condition	n Category				
	Parameter	Optimal	Suboptimal	Marginal	Poor			
	6. Channel Alteration	Channelization or dredging absent or minimal; stream with normal pattern.	Some channelization present, usually in areas of bridge abutments; evidence of past channelization, i.e., dredging, (greater than past 20 yr) may be present, but recent channelization is not present.	Channelization may be extensive; embankments or shoring structures present on both banks; and 40 to 80% of stream reach channelized and disrupted.	Banks shored with gabion or cement; over 80% of the stream reach channelized and disrupted. Instream habitat greatly altered or removed entirely.			
	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
oling reach	7. Frequency of Riffles (or bends)	Occurrence of riffles relatively frequent; ratio of distance between riffles divided by width of the stream <7:1 (generally 5 to 7); variety of habitat is key. In streams where riffles are continuous, placement of boulders or other large, natural obstruction is important.	Occurrence of riffles infrequent; distance between riffles divided by the width of the stream is between 7 to 15.	Occasional riffle or bend; bottom contours provide some habitat; distance between riffles divided by the width of the stream is between 15 to 25.	Generally all flat water or shallow riffles; poor habitat; distance between riffles divided by the width of the stream is a ratio of >25.			
samp	SCORE	20 19 18 17 16	15 14 13 12 11	10 9 8 7 6	5 4 3 2 1 0			
Parameters to be evaluated broader than sampling reach	8. Bank Stability (score each bank) Note: determine left or right side by facing downstream.	Banks stable; evidence of erosion or bank failure absent or minimal; little potential for future problems. <5% of bank affected.	Moderately stable; infrequent, small areas of erosion mostly healed over. 5-30% of bank in reach has areas of erosion	Moderately unstable; 30-60% of bank in reach has areas of erosion; high erosion potential during floods.	Unstable; many eroded areas; "raw" areas frequent along straight sections and bends; obvious bank sloughing; 60-100% of bank has erosional scars.			
e eva	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
to be	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
Parameters	9. Vegetative Protection (score each bank)	More than 90% of the streambank surfaces and immediate riparian zone covered by native vegetation, including trees, understory shrubs, or nonwoody macrophytes; vegetative disruption through grazing or mowing minimal or not evident; almost all plants allowed to grow naturally.	70-90% of the streambank surfaces covered by native vegetation, but one class of plants is not well-represented; disruption evident but not affecting full plant growth potential to any great extent; more than one-half of the potential plant stubble height remaining.	50-70% of the streambank surfaces covered by vegetation; disruption obvious; patches of bare soil or closely cropped vegetation common; less than one- half of the potential plant stubble height remaining.	Less than 50% of the streambank surfaces covered by vegetation; disruption of streambank vegetation is very high; vegetation has been removed to 5 centimeters or less in average stubble height.			
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			
	10. Riparian Vegetative Zone Width (score each bank riparian zone)	Width of riparian zone >18 meters; human activities (i.e., parking lots, roadbeds, clear-cuts, lawns, or crops) have not impacted zone.	Width of riparian zone 12-18 meters; human activities have impacted zone only minimally.	Width of riparian zone 6- 12 meters; human activities have impacted zone a great deal.	Width of riparian zone <6 meters: little or no riparian vegetation due to human activities.			
	SCORE (LB)	Left Bank 10 9	8 7 6	5 4 3	2 1 0			
ĺ	SCORE (RB)	Right Bank 10 9	8 7 6	5 4 3	2 1 0			

Total	Caama	
i otai	Score	

BENTHIC MACROINVERTEBRATE FIELD DATA SHEET

STREAM NAME		LOCATION					
STATION #	_ RIVERMILE	STREAM CLASS					
LAT	LONG	RIVER BASIN					
STORET#		AGENCY					
INVESTIGATORS			LOT NUMBER				
FORM COMPLETED BY		DATE REASON FOR SURVEY TIME					
HABITAT TYPES	Indicate the percentage of each habitat type present						

HABITAT TYPES	Indicate the percentage of each habitat type present Cobble% Snags% Vegetated Banks% Sand% Submerged Macrophytes% Other ()%
SAMPLE COLLECTION	Gear used D-frame kick-net Other
	How were the samples collected? wading from bank from boat
	Indicate the number of jabs/kicks taken in each habitat type. Cobble Snags Vegetated Banks Sand
	Cobble Snags Vegetated Banks Sand Submerged Macrophytes Other ()
GENERAL COMMENTS	

QUALITATIVE LISTING OF AQUATIC BIOTA

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare, 2 = Common, 3 = Abundant, 4 = Dominant

Periphyton	0	1	2	3	4	Slimes	0	1	2	3	4
Filamentous Algae	0	1	2	3	4	Macroinvertebrates	0	1	2	3	4
Macrophytes	0	1	2	3	4	Fish	0	1	2	3	4

FIELD OBSERVATIONS OF MACROBENTHOS

Indicate estimated abundance: 0 = Absent/Not Observed, 1 = Rare (1-3 organisms), 2 = Common (3-9 organisms), 3 = Abundant (>10 organisms), 4 = Dominant (>50 organisms)

Porifera	0	1	2	3	4	Anisoptera	0	1	2	3	4	Chironomidae	0	1	2	3	4
Hydrozoa	0	1	2	3	4	Zygoptera	0	1	2	3	4	Ephemeroptera	0	1	2	3	4
Platyhelminthes	0	1	2	3	4	Hemiptera	0	1	2	3	4	Trichoptera	0	1	2	3	4
Turbellaria	0	1	2	3	4	Coleoptera	0	1	2	3	4	Other	0	1	2	3	4
Hirudinea	0	1	2	3	4	Lepidoptera	0	1	2	3	4						
Oligochaeta	0	1	2	3	4	Sialidae	0	1	2	3	4						
Isopoda	0	1	2	3	4	Corydalidae	0	1	2	3	4						
Amphipoda	0	1	2	3	4	Tipulidae	0	1	2	3	4						
Decapoda	0	1	2	3	4	Empididae	0	1	2	3	4						
Gastropoda	0	1	2	3	4	Simuliidae	0	1	2	3	4						
Bivalvia	0	1	2	3	4	Tabinidae	0	1	2	3	4						
						Culcidae	0	1	2	3	4						

WOLMAN PEBBLE COUNT FORM

Stream ID: S-Q2

County: Pittsylvania
Stream Name: UNT to Pole Bridge Branch
HUC Code: 03010105 Basin: Banister

Survey Date: 8/19/2021 Surveyors: SK VM Type: Representative

			LE COUNT			I:	0/ =
Inches	PARTICLE	Millimeters		Particle Count	Total #	Item %	% Cum
	Silt/Clay	< .062	S/C	A	30	30.00	30.00
	Very Fine	.062125		•		0.00	30.00
	Fine	.12525		•		0.00	30.00
	Medium	.255	SAND	A		0.00	30.00
	Coarse	.50-1.0		*		0.00	30.00
.0408	Very Coarse	1.0-2		A	33	33.00	63.00
.0816	Very Fine	2 -4		A		0.00	63.00
.1622	Fine	4 -5.7		A		0.00	63.00
.2231	Fine	5.7 - 8		^	15	15.00	78.00
.3144	Medium	8 -11.3		^		0.00	78.00
.4463	Medium	11.3 - 16	GRAVEL	•		0.00	78.00
.6389	Coarse	16 -22.6		A	6	6.00	84.00
.89 - 1.26	Coarse	22.6 - 32		A		0.00	84.00
1.26 - 1.77	Vry Coarse	32 - 45		•		0.00	84.00
1.77 -2.5	Vry Coarse	45 - 64		-		0.00	84.00
2.5 - 3.5	Small	64 - 90		-		0.00	84.00
3.5 - 5.0	Small	90 - 128	COBBLE	-	9	9.00	93.00
5.0 - 7.1	Large	128 - 180	COBBLE	-	1	1.00	94.00
7.1 - 10.1	Large	180 - 256		-	6	6.00	100.00
10.1 - 14.3	Small	256 - 362		A		0.00	100.00
14.3 - 20	Small	362 - 512		▲		0.00	100.00
20 - 40	Medium	512 - 1024	BOULDER	A		0.00	100.00
40 - 80	Large	1024 -2048]	•		0.00	100.00
80 - 160	Vry Large	2048 -4096]	•		0.00	100.00
	Bedrock		BDRK	A		0.00	100.00
				Totals:	100		

RIVERMORPH PARTICLE SUMMARY

UNT to Pole Bridge Branch

River Name: Reach Name: Sample Name: Survey Date: S-Q2 Representative 08/19/2021

Size (mm)	TOT #	ITEM %	CUM %
0 - 0.062 0.062 - 0.125 0.125 - 0.25 0.25 - 0.50 0.50 - 1.0 1.0 - 2.0 2.0 - 4.0 4.0 - 5.7 5.7 - 8.0 8.0 - 11.3 11.3 - 16.0 16.0 - 22.6 22.6 - 32.0 32 - 45 45 - 64 64 - 90 90 - 128 128 - 180 180 - 256 256 - 362 362 - 512 512 - 1024 1024 - 2048 Bedrock	30 0 0 0 0 0 33 0 0 15 0 0 6 0 0 0 9 1 6 0 0	30.00 0.00 0.00 0.00 0.00 0.00 33.00 0.00	30.00 30.00 30.00 30.00 30.00 63.00 63.00 78.00 78.00 78.00 84.00 84.00 84.00 84.00 93.00 94.00 100.00 100.00 100.00 100.00
D16 (mm) D35 (mm) D50 (mm) D84 (mm) D95 (mm) D100 (mm) Silt/Clay (%) Sand (%) Gravel (%) Cobble (%) Boulder (%) Bedrock (%)	0.03 1.15 1.61 22.6 192.67 256 30 33 21 16 0		

Total Particles = 100.

Stream Assessment Form (Form 1) Unified Stream Methodology for use in Virginia le channels classified as intermittent or perennia Cowardin **Impact** Impact Project # t Name (Applicant) Locality HUC SAR# Date Class Length Factor Mountain Valley Pipeline (Mountain 22865.06 Pittsylvania R3 03010105 8-19-21 S-Q2 20 1 Valley Pipeline, LLC) Stream Name and Information SAR Length Name(s) of Evaluator(s) **UNT to Pole Bridge Branch** 114 SK, VM 1. Channel Condition: Assess the cross-section of the stream and prevailing condition (erosion, aggradation) Optimal Suboptimal Marginal Poor Severe Very little incision or active erosion; 80-Slightly incised, few areas of active Often incised, but less than Severe or Overwidened/incised. Vertically / Deeply incised (or excavated) oor. Banks more stable than Severe of oor due to lower bank slopes. Erosion may be present on 40-60% of 100% stable banks. Vegetative surfact protection or natural rock, prominent or unprotected banks. Majorit vertical/lateral instability. Severe sion, flow contained within the bank laterally unstable. Likely to wide of banks are stable (60-80%) Majority of both banks are ne Channel (80-100%). AND/OR Stable point bars Vegetative protection or natural rock vertical. Erosion present on 60-80% of Streambed below average rooting depth Condition bankfull benches are present. Access to their original floodplain or fully prominent (60-80%) AND/OR Depositional features contribute to both banks. Vegetative protection on 40-60% of banks. Streambanks may be panks. Vegetative protection present or 20-40% of banks, and is insufficient to prevent erosion. AND/OR 60-80% of majority of banks vertical/undercut. Vegetative protection present on less than 20% of banks, is not preventing developed wide bankfull benches. Midstability. The bankfull and low flow vertical or undercut. AND/OR channel bars and transverse bars few. Transient sediment deposition covers channels are well defined. Stream like has access to bankfull benches,or new 40-60% Sediment may be temporary transient, contribute instability. the stream is covered by sediment. Sediment is temporary / transient in erosion. Obvious bank sloughing sent. Erosion/raw banks on 80-100% less than 10% of bottom developed floodplains along Deposition that contribute to stability nature, and contributing to instability AND/OR Aggrading channel. Greater portions of the reach. Transient diment covers 10-40% of the stream may be forming/present. AND/OR V-shaped channels have vegetative AND/OR V-shaped channels have vegetative protection is present on > than 80% of stream bed is covered by deposition, contributing to instability. bottom protection on > 40% of the banks and 40% of the banks and stable sediment Multiple thread channels and/or epositional features which contribute to stability. deposition is absent subterranean flow. CI 2.4 Scores 3 1.6 1 2.40 NOTES>> 2. RIPARIAN BUFFERS: Assess both bank's 100 foot riparian areas along the entire SAR. (rough measurements of length & width may be acceptable) **Conditional Category** NOTES>> Optimal Suboptimal Marginal Poor Low Marginal: Non-maintained, High Poor: Lawn mowed, and High Suboptima Low Suboptimal Low Poor High Marginal dense herbaceous maintained areas Riparian areas with Riparian areas with Impervious surfaces, mine Non-maintained egetation, riparia nurseries: no-till ee stratum (dbh: ree stratum (dbh > eas lacking shrul nse herbaceo cropland; actively 3 inches) present 3 inches) present Tree stratum (dbh > 3 inches) present, vegetation with and tree stratum grazed pasture, spoil lands, Riparian with 30% to 60% with 30% to 60% either a shrub laye or a tree layer (dbl hay production, onds, open water parsely vegetate non-maintained with > 60% tree canopy cover. nuded surfaces tree canopy cover and containing bot tree canopy cover and a maintained **Buffers** Wetlands located within the riparian areas 3 inches) present If present, tree area, recently feed lots, trails, or herbaceous and nderstory. Recent cutover (dense with <30% tree stratum (dbh >3 seeded and other comparable shrub layers or a non-maintained inches) present, tabilized, or other conditions canopy cover vegetation). with <30% tree comparable understory. canopy cover with maintained condition. understory. High Low High Low High Low 1.5 0.5 Scores 1.2 1.1 0.85 0.75 0.6 1. Delineate riparian areas along each stream bank into Condition Categories and Condition Scores using the descriptors Ensure the sums 2. Determine square footage for each by measuring or estimating length and width. Calculators are provided for you below of % Riparian Enter the % Riparian Area and Score for each riparian category in the blocks below. Blocks equal 100 % Riparian Area> 100% 100% Right Bank 0.85 Score > CI= (Sum % RA * Scores*0.01)/2 % Riparian Area> 100% 100% Rt Bank CI > CI Left Bank 0.85 0.85 0.85 3. INSTREAM HABITAT: Varied substrate sizes, water velocity and depths; woody and leafy debris; stable substrate; low embededness; shade; undercut banks; root mats; SAV; riffle/pool complexes, stable features **Conditional Category** NOTES>> Optimal Suboptimal Marginal Poor Instream Habitat/ nents are typically Habitat elements listed above are Available esent in 10-30% of the reach and are Habitat elements are typically present i resent in 30-50% of the reach and are lacking or are unstable. Habitat greater than 50% of the reach adequate for maintenance of adequate for maintenance of elements are typically present in less than 10% of the reach. Cover populations. populations Stream Gradient CI

Scores

1.5

1.2

0.9

0.5

High / Low

1 50

	S	tream li	mpact A	ssessn	nent For	m Pag	e 2		
Project #	Project Name (App	licant)	Locality	Cowardin Class.	нис	Date	SAR # / Data Point	Impact / SAR length	Impact Factor
22865.06	Mountain Valley Pipeline (Mountain Valley Pipeline, LLC)		Pittsylvania	R3	03010105	8-19-21	S-Q2	20	1
. CHANNEL	. ALTERATION: Stream crossing	gs, riprap, concret			ghtening of channe	el, channelization	, embankments, sp		s, livestock
	Negligible	NA:	nor	al Category	lerate		evere	NOTES>>	
Channel Alteration	Channelization, dredging, alteration, or hardening absent. Stream has an unaltered pattern or has naturalized.	the channel		40 - 60% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	60 - 80% of reach is disrupted by any of the channel alterations listed in the parameter guidelines. If stream has been channelized, normal stable stream meander pattern has not recovered.	by any of the cha in the parameter 80% of banks s	of reach is disrupted nnel alterations listed guidelines AND/OR hored with gabion, or cement.		
Scores	1.5	1.3	1.1	0.9	0.7	(0.5		
	REACH	CONDITION	INDEX and	STREAM CO	NDITION UN	ITS FOR TH	IS REACH		

THE REACH CONDITION INDEX (RCI) >> 1.25

RCI= (Sum of all Cl's)/5, except if stream is ephemeral RCI = (Riparian Cl/2)

COMPENSATION REQUIREMENT (CR) >> 25

CR = RCI X L_i X IF

INSERT PHOTOS:

NOTE: The CIs and RCI should be rounded to 2 decimal places. The CR should be rounded to a whole number.



DESCRIBE PROPOSED IMPACT:

PROVIDED UNDER SEPARATE COVER

